



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

January 28, 2026

Billie Espejo
Regulatory Fellow
Janssen PMP
1125 Trenton-Harbourton Road
Titusville, NJ 08560-0200

Subject: Label Amendment - Registration Review Mitigation for Imazalil and Imazalil Sulfate
Product Name: FUNGAFLOR 750 WSP
EPA Registration Number: 43813-59
Case Number: 476809
Application Dates: June 22, 2021

Dear Billie Espejo:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Imazalil and Imazalil Sulfate Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must

submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Kelsi Grogan by phone at 202-566-2228, or via email at grogan.kelsi@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Kevin Costello".

Kevin Costello, Branch Chief
Risk Management and Implementation Branch 2
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER PELIGRO

Corrosive. Causes irreversible eye damage. May be fatal if swallowed or inhaled. Harmful if absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Do not breathe dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Handlers must wear goggles or face shield, long-sleeved shirt and long pants, shoes, socks, chemical resistant gloves made of any waterproof material, such as natural rubber, polyethylene, PVC, neoprene, rubber, nitrile rubber, or butyl rubber. Wear a minimum of a NIOSH-approved particulate respirator facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washable exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This chemical is toxic to fish and may not apply directly to water, to grass, when surface water is present or to intertidal areas below the mean high-water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application.

ACCEPTED
Jan 28, 2026
Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under
EPA Reg. No. 43813-59

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool dry place in the original container away from food and feed.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows:

[For plastic containers that are small enough to shake] – Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ½ full with water and recap. Shake for 10 seconds.

Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

[For fiber drums or paperboard containers with plastic liners] – Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment.

Then offer for recycling if available or dispose of in a sanitary landfill, or by incineration. If drum is contaminated and cannot be reused, dispose of it in a manner required for its liner.

FUNGICIDE RESISTANCE MANAGEMENT

IMAZALIL GROUP 3 FUNGICIDE

FUNGAFLOR 750 WSP is an imidazole fungicide in the Group 3 class of fungicides that inhibits sterol biosynthesis which results in disruption of membrane structure and function, thus affecting cell wall permeability. Because resistance development cannot be predicted, postharvest resistance management strategies and sanitation practices should be followed. Fungicides with acquired resistance to Group 3 fungicides, may eventually dominate the fungal population if Group 3 fungicides are used repeatedly or in successive years as the primary method of disease control for the

IMAZALIL GROUP 3 FUNGICIDE
FUNGAFLOR® 750 WSP

Water Soluble Powder

For postharvest treatment of Citrus Fruit postharvest: Calamondin, Citrus citron, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin, Orange sweet, Orange sour, Pummelo, Satsuma mandarin.

ACTIVE INGREDIENT: (W/W) IMAZALIL SULFATE: 1-2-(2,4-dichlorophenyl)-2-(2-propenyl)oxy-ethyl)-1H-imidazole sulfate* 100%

(*Equivalent to 75% imazalil base)

**KEEP OUT OF REACH OF CHILDREN
DANGER – PELIGRO**

FIRST AID

If in eyes	- Hold eye open and rinse slowly and gently with water for 15-20 minutes. - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. - Call a poison control center or doctor for treatment advice.
If swallowed	- Call a poison control center or doctor immediately for treatment advice. - Have person sip a glass of water if able to swallow. - Do not induce vomiting unless told to do so by a poison control center or doctor. - Do not give anything by mouth to an unconscious person.
If inhaled	- Move person to fresh air. - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. - Call a poison control center or doctor for further treatment advice.
If on skin or clothing	- Take off contaminated clothing. - Rinse skin immediately with plenty of water for 15-20 minutes. - Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency assistance call CHEMTREC at 1-800-424-9300. For non-emergency information concerning this product call the National Pesticide Information Center (NPIC) at 1-800-858-7378 (NPIC Web site: www.npic.orst.edu).

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

NET Contents:

EPA Reg. No. 43813-59
EPA EST. NO.

JANSSEN PMP
PRESERVATION AND MATERIAL PROTECTION
1125 Trenton-Harbourton Rd.
Titusville, NJ 08560-0200, USA

targeted pathogens. This may result in partial or total loss of control of those fungal species with imazalil or other Group 3 fungicides. A disease management program that includes alternation or tank mixes between FUNGAFLOR 750 WSP and other labeled fungicides that have a different mode of action, may prevent pathogen populations from developing resistance.

To delay fungicide resistance, take one or more of the following steps:

• Rotate the use of imazalil or other Group 3 fungicides, during a growing season sequence, such as long-term storage and prior to extended transport conditions, with different fungicide groups that control the same pathogens and in which the pathogen population is still highly sensitive to that fungicide or fungicide Group.

• Use tank mixtures or alternate with combination products containing fungicides from different fungicide groups that are equally effective when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.

• Adopt an integrated disease management program for fungicide use that includes monitoring fungal populations in the packinghouse and fruit storage rooms for resistance development, using different available sensitivity monitoring techniques; harvest information related to pesticide use, impact of packaging storage conditions on disease development and thresholds, as well as cultural, biological and other chemical control practices.

• Where possible and practical, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.

• Contact your local service company, Janssen PMP, or your local extension specialist for information on packinghouse spore monitoring programs and for any additional resistance management and/or Integrated Pest Management (IPM) recommendations for specific crops and pathogens.

• For further information, or to report suspected resistance contact Janssen PMP at info@janssennpm.com. You can also contact your pesticide distributor service company or university extension specialist to report resistance.

Intended for post-harvest control of *Penicillium digitatum* (green mold), *Penicillium italicum* (blue mold), *Diaporthe citri* (phomopsis stem end rot), *Diplodia natalensis* (diplodia rot) and *Alternaria citri* on citrus fruits.

FUNGAFLOR 750 WSP is formulated for aqueous application only; not compatible with storage or shipping waxes. Used as directed, FUNGAFLOR 750 WSP has curative and preventative action as well as anti-sporulation properties. Use of FUNGAFLOR

750 WSP should be integrated into an overall disease management program within each packinghouse.

DOSE REQUIREMENTS:

In dips, wash tanks, and drenches: Weigh out 9 - 13.5 ounces of imazalil or equivalent imazalil of FUNGAFLOR 750 WSP and dissolve in 100 gallons of water to give a treating rate of 500-750 ppm of active ingredient. The treatment solutions must be monitored to insure the proper concentration of FUNGAFLOR 750 WSP. Length of dip time must not exceed two minutes.

In non-recovery spray & water spray brush application: Dissolve 18 ounces (0.84 lbs. active ingredient imazalil) of FUNGAFLOR 750 WSP in 100 gallons of water to achieve a concentration of 1000 ppm FUNGAFLOR 750 WSP active ingredient. Apply to freshly cleaned citrus prior to waxing.

Remarks

• Combinations of above-listed treatments are possible, but never more than two (2) treatments may be carried out on the same citrus lot.

• Application restrictions for citrus drench: Stay outside of treatment area until citrus is treated and drained. The windows and doors of the transport vehicle must be closed prior to treatment.

NOTICE OF WARRANTY:

Janssen PMP, a Division of Janssen Pharmaceutica NV warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the presence of other pests. The application of which may be beyond the control of Janssen PMP. To the extent consistent with applicable law, Janssen PMP disclaims any liability for consequential, special or indirect damages resulting from the use or handling of this product. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, JANSSEN PMP MAKES NO WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

©Trademark of Janssen Pharmaceutica NV