LOCUCIDE BAIT

Locucide Bait contains a microsporidial pathogen of grasshoppers. It infects the fat bodies of most species of grasshoppers and some crickets. Mortality is greatest to young grasshoppers. Honey bees are not infected by this pathogen.

PRECAUTIONARY STATMENTS

Hazards to humans:
Avoid inhalation or contact with eyes or open wounds. Wash hands after using.

STORAGE and DISPOSAL

Storage:
Store in a cool dry location if not used immediately. Product must be used within two weeks of formulation.

Disposal:
Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

INSTRUCTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Locucide Bait is for use in controlling grasshoppers and Morman crickets. Apply to rangelands at a rate of 1.5 pounds per acre. A higher rate of application may be used on severe infestations. Application may be accomplished by hand, seed spreaders, or turbine spreaders.

Grasshoppers are most effectively controlled when they are young. For best results apply Locucide Bait when grasshoppers are ½ to ¼ inch long. Grasshopper death from this pathogen will begin in 3 to 6 weeks, and can remain active in controlling grasshoppers for several years.

CODE #   Date formulated

Nosema locustae, a biological insecticide for use in controlling grasshoppers and crickets.

ACTIVE INGREDIENT
Nosema locustae Canning
At least 600 million spores per lb. 0.05%

INERT INGREDIENT 99.95%

100.00%

CAUTION
KEEP OUT OF REACH OF CHILDREN
PRACTICAL TREATMENT

IF IN EYES — Flush with plenty of water. Get medical attention if irritation develops.

IF ON SKIN OR CLOTHING — Remove contaminated clothing. Wash skin with plenty of soap and water.

See side panel for additional Precautionary Statements.

Manufactured by LAPORTE INSECTARIES
2220 N. Hwy. U.S. 287
Fort Collins, CO 80524

EPA Est. 43915-CO-01
EPA Reg. No. 43915-2

Net Wt.   10 lbs.

43915-2

15

LOCUCIDE BAIT

Locucide Bait contains a microsporidial pathogen of grasshoppers. It infects the fat bodies of most species of grasshoppers and some crickets. Mortality is greatest to young grasshoppers. Honey bees are not infected by this pathogen.

PRECAUTIONARY STATMENTS

Hezards to humans:
Avoid inhalation or contact with eyes or open wounds Wash hands after using.

STORAGE and DISPOSAL

Storage:
Store in a cool dry location if not used immediately. Product must be used within two weeks of formulation.

Disposal:
Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

INSTRUCTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Locucide Bait is for use in controlling grasshoppers and Morman crickets. Apply to rangelands at a rate of 1.5 pounds per acre. A higher rate of application may be used on severe infestations. Application may be accomplished by hand, seed spreaders, or turbine spreaders.

Grasshoppers are most effectively controlled when they are young. For best results apply Locucide Bait when grasshoppers are ½ to ¾ inch long. Grasshopper death from this pathogen will begin in 3 to 6 weeks, and can remain active in controlling grasshoppers for several years.

CODE #   Date formulated

LOCUCIDE BAIT™

Nosema locustae, a biological insecticide for use in controlling grasshoppers and crickets.

ACTIVE INGREDIENT
Nosema locustae Canning
At least 600 million spores per lb. 0.05%

INERT INGREDIENT 99.95%

100.00%

CAUTION
KEEP OUT OF REACH OF CHILDREN
PRACTICAL TREATMENT

IF IN EYES — Flush with plenty of water. Get medical attention if irritation develops.

IF ON SKIN OR CLOTHING — Remove contaminated clothing. Wash skin with plenty of soap and water.

See side panel for additional Precautionary Statements.

Manufactured by LAPORTE INSECTARIES
2220 N. Hwy. U.S. 287
Fort Collins, CO 80524

EPA Est. 43915-CO-01
EPA Reg. No. 43915-2

Net Wt   50 lbs.