

**HAZAROUS TO HUMANS AND DOMESTIC ANIMALS**

**PHYSICAL HAZARDS: CANCER.**

Corrosive; may cause severe skin irritation or chemical burns to broken skin. Causes eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling this product. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until odors have dissipated.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish. Keep out of lakes, streams or ponds. Treated effluent may not be discharged into lakes, streams, ponds or public waters without a valid discharge permit. For guidance, contact the regional office of the Environmental Protection Agency.

**CHEMICAL HAZARDS: STRONG OXIDIZING AGENT.**

Mix only with water according to label directions. Mixing this product with gross filth such as feces, urine, etc., or with ammonia, acids, detergents or other chemicals may release hazardous gases irritating to eyes, lungs and mucous membranes.

**STORAGE AND DISPOSAL:**

Store in a cool, dry area away from direct sunlight. In case of spill, flood with large quantities of water. Rinse empty container thoroughly with water and return to manufacturer or place in trash collection for burying in an approved landfill. Product or rinsate that cannot be used should be diluted with water and disposed of in a sanitary sewer. Do not contaminate food or feed by storage, disposal or cleaning of equipment.

"This pesticide is toxic to fish and aquatic organisms. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or public waters unless this product is specifically identified and addressed in an NPDES permit. Do not discharge effluent containing this product to sewer systems without previously notifying the sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA."

**BEST AVAILABLE COPY**



NO DEPOSIT  
NO RETURN

Active Ingredients Sodium Hypochlorite . . . 10%  
Inert Ingredients . . . . . 90%


**WARNING:**  
**KEEP OUT OF REACH OF CHILDREN**

**FIRST AID:**  
For Eyes, flush liberally with water for at least 15 minutes. Seek prompt medical attention.  
For Skin, wash with soap and water. If swallowed, drink large amounts of water. Do NOT induce vomiting. Do NOT give vinegar or other acids. Do NOT induce vomiting. Get prompt medical attention.

See additional precautions on side panel.

**1 Gallon (128 Fl. Oz.) 3.785 Liters**

NO "S" IN  
the word  
"Ingredient"



"IF SWALLOWED, drink large amounts of water. DO NOT induce vomiting. Call a physician or poison control center immediately."

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

#### DIRECTIONS FOR SWIMMING POOL CHLORINATION

For a new pool or for spring start-up: superchlorinate with one pint (16 fl. oz.) of this sodium hypochlorite solution for each 2500 gallons of water. This dosage is equivalent to 5 ppm available chlorine by weight.

For pool maintenance:

- A) Adjust pool water pH to 7.2-7.8 range and maintain.
- B) Add manually or by a feeder device this sodium hypochlorite solution at a rate to maintain an available chlorine residual of 0.6 to 1.0 ppm. One-half pint (8 fl. oz.) of this solution will provide 1.0 ppm available chlorine for each 6000 gallons of water. Frequency of additions to maintain available chlorine level will depend upon temperature and number of swimmers. Use a reliable test kit daily to make certain the pH and chlorine residual are in the proper range.

NOTE: This product degrades with age. Use a reliable chlorine test kit and increase dosage, as necessary, to obtain the required level of available chlorine.

**CAUTION:** Keep upright, this Cap is Vented.

EPA Reg. No. 43922-2-37312

EPA Est. No. 43922-MI-1

### DIRECTIONS FOR DISINFECTION OF POTABLE WATER FOR HOME WELL WATER SYSTEMS

Dilute this sodium hypochlorite solution in the ratio of one part solution to nine parts softened water. Mix thoroughly and begin feeding with a hypochlorinator (metering pump). Maintain a free available chlorine residual of at least 0.2 ppm and not more than 0.8 ppm throughout the distribution system, as determined by a DPD chlorine test kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed by the National Interim Primary Drinking Water Regulations. Check water frequently with a DPD chlorine test kit. For further details, contact your local Health Department.


### DIRECTIONS FOR SANITIZING FOOD PROCESSING OR DAIRY EQUIPMENT

Clean equipment in the normal manner. Just before using, rinse all surfaces thoroughly with a sodium hypochlorite solution containing 200 ppm available chlorine. One ounce (2 blaps.) of this solution per four gallons of water will provide approx. 200 ppm available chlorine. Maintain contact with disinfectant for a minimum of two minutes. Do not rinse with water after treatment. Do not soak overnight.

DISTRIBUTED BY

**LITHOUSE PRODUCTS, INC.  
STRONGSVILLE, OH 44138**

**UN1791**



**REPROducible COPY**