

CONTENTS

55 POUNDS



ACCREDITED
 March 27, 1973
 43981-6

ZINEB TECHNICAL

AGRICULTURAL FUNGICIDE

Active Ingredient:	
Zineb (Zinc ethylene bisdithiocarbamate)	93%
Inert Ingredients	7%
	<u>100%</u>

SOLD FOR MANUFACTURING PURPOSES ONLY
 FOR THE FORMULATION OF ECONOMIC POISONS

CAUTION

KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed. Do not take internally. Avoid skin contact. Do not breathe dust. May cause irritation of nose and throat. Wash thoroughly with plenty of water after handling. In case of swallowing, apply stomach drain (by doctors only), or induce vomiting. Do not reuse container. Destroy when empty.

KEEP AWAY FROM FOODSTUFFS AND FEEDS
 STORE IN COOL, DRY PLACES, IN CLOSED PACKAGES

Notice: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use/or handling is contrary to label instructions.

Manufactured for:

CHEMPAR CHEMICAL CO., INC.
 200 MADISON AVENUE
 NEW YORK, N. Y. 10016

EPA Reg. No. 43981-6

BEST DOCUMENT AVAILABLE

23

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR THE USE OF THE FORMULATION ZINEB 75% W.P.

FOR THE CONTROL OF FUNGUS DISEASES IN GARDEN,

FRUIT AND FIELD CROPS.

ZINEB IS COMPATIBLE WITH MOST INSECTICIDES,
FUNGICIDES, AND NUTRITIONAL SPRAYS AND MAY BE USED
IN COMBINATION WHEN CONDITIONS WARRANT.

WARNING

KEEP OUT OF THE REACH OF CHILDREN

May cause irritation of nose, throat and skin. Do not breathe dust or spray mist. In case of prolonged or frequent use, wear clean protective clothing and bathe after each exposure. Keep bag closed. Store in dry place. Keep away from fire and sparks.

Do not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.

Use Chempar Zineb 75% W.P. at the rate of 1-1/2 to 2 pounds per 100 gallons of water (1 level tablespoonful per gallon) except where otherwise noted. Partially fill spray tank with water, add desired amount of Chempar Zineb 75% W.P. slowly while agitating. If insecticides or other spray ingredients are to be added to solution, mix Chempar Zineb 75% W.P. with water first. After mixing thoroughly add remaining water to spray tank.

Spray at the rate of 100 to 125 gallons per acre for low growing crops, up to 200 gallons for heavy foliage crops such as tomato, and potato vines.

VEGETABLES

Blight
TOMATOES ----- Early and Late Blight : Anthracnose, Septoria Leaf Spot - Start sprays when disease first appears in area, repeat applications on 7 to 10 day schedule or more often with severe late blight conditions.

CANTALOUPE, CUCUMBERS, WATERMELONS - Downy Mildew, Alternaria Blight, Anthracnose : Apply on weekly schedule, direct spray to underside of leaves. Begin applications when plants appear above ground or when disease appears.

beans
BEANS (SNAP AND LIMA) - Downy Mildew, Rust, Anthracnose: Spray at 7 to 10 day intervals, on Rust, begin when disease threatens, on Anthracnose, begin after first blossom appears. Do not apply within 7 days of harvest.

EPA Reg. NO. 43981-6

over
18

FRUITS

BEST DOCUMENT AVAILABLE

Note: Follow state spray schedule in commercial orchards. In home plantings apply as indicated. Cover plants thoroughly.

APPLES - Scab, Cedar Rust, Frog-eye (Black Rot), sooty Blotch;
Use 1 to 2 pounds Chempar Zineb 75% W.P. per 100 gallons of water from delayed dormant through the remainder of season, depending upon the severity of the disease conditions.

APPLES AND PEARS : Fire Blight - Use 2 lbs. Chempar Zineb 75% W.P. per 100 gallons in 2 to 3 applications at 10% bloom, full bloom and petal fall. Maintain recommended sanitary pruning practices.

Apply just before bloom, just after bloom, 10 days after bloom and 2-3 weeks later for :

GRAPES - Downy mildew, Black ^{ROT} Spot, Ripe Rot : Use 1-1/2 lbs. per 100 gallons

Do not apply within 7 days of harvest on Grapes and pears

ORNAMENTALS

Note - Use Chempar Zineb 75% W.P. at 1-1/2 lbs. per 100 gallons of water or 1 level tablespoonful per gallon, spraying ~~the~~ plants thoroughly.

ROSES - Black Spot : Apply at weekly intervals, beginning when disease appears.

CHRYSANTHEMUMS, CARNATIONS - Rust Leafspot and Anthracnose: Apply at 7 to 10 day intervals as soon as disease condition threatens.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

CHEMPAR CHEMICAL CO. INC.
260 Madison Avenue
New York, N.Y. 10016

EPA Reg No. 43981-6

19