

Sanitizing of Food Processing Equipment and other Hard Non-Porous Surfaces in Good Contact Locations

For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sinks, countertops, floors, and other hard non-porous surfaces. No Potable water rinse is required.

Wash and rinse all articles thoroughly, then apply a solution of 1 oz PCI-DISINFECTANT in 4 gallons of water (150 ppm active). Surfaces should remain wet for at least one minute followed by adequate draining and/or drying. Fresh solution should be prepared daily or when use solution becomes visibly dirty. For mechanical application, use solution may not be re-used for sanitizing purposes.

Dishes, silverware, glasses, cooking utensils and other similar size food processing equipment can be sanitized by immersion in a 1 oz / 4 gallon dilution of PCI-DISINFECTANT. No Potable water rinse is required.

At 1 oz / 4 gallons, PCI-DISINFECTANT fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinance 1978 Recommendations of the U.S. Public Health Service in waters up to 1000 ppm of hardness calculated as Ca CO₃ when evaluated by the AOAC Germicidal and Disinfectant Sanitizer Method against Escherichia coli and Staphylococcus aureus.

The udders, flanks, and teats of dairy cows can be sanitized by washing with a solution of 1 oz PCI-DISINFECTANT in 4 gallons of warm water. No Potable water rinse is required. Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

Precautionary Statements

Hazards to Humans and domestic animals

DANGER

Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

STORAGE AND DISPOSAL

—DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

—OPEN DUMPING IS PROHIBITED

—DO NOT REUSE EMPTY CONTAINER

PESTICIDE DISPOSAL

PESTICIDE OR RINSATE THAT CANNOT BE USED OR CHEMICALLY REPROCESSED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

CONTAINER DISPOSAL

TRIPLE RINSE (OR EQUIVALENT) AND DISPOSE IN AN INCINERATOR OR LANDFILL APPROVED FOR PESTICIDE CONTAINERS, OR BURY IN A SAFE PLACE.

FOR HOME USE: DO NOT REUSE EMPTY CONTAINER. WRAP CONTAINER AND PUT IN TRASH COLLECTION.

GENERAL

CONSULT FEDERAL, STATE OF LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES SUCH AS LIMITED OPEN BURNING.

PCI-DISINFECTANT

Disinfectant - Sanitizer - Fungicide
Deodorizer for Hospital, Institutional,
Industrial, School, Dairy and Other
Farm and Home Use

Active Ingredients

Didecyl dimethyl ammonium chloride 7.5%

Inert Ingredients

92.5%

100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE LEFT PANAL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Registration No. 43998-2

EPA Establishment No. 43998

Manufactured by:



Pickwick Chemical International
PO Box BC Florence Alabama 35630

ACCEPTED

MAR 1 1 1981

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended for the pesticide
registered under
EPA Reg. No. 43998-2

**Directions for Use
GENERAL CLASSIFICATION**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply PCI-DISINFECTANT with a cloth, mop or mechanical spray device. When applied with a mechanical spray device, surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the use solution becomes visibly dirty.

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions.

For disinfecting floors, walls, countertops, bathing areas, lockers, bedrooms, tables, chairs, garbage pails and other hard non-porous surfaces.

Add 3 1/2 oz PCI-DISINFECTANT to 4 gallons water, apply to previously cleaned hard non-porous surface with mop or cloth.

At this use level, PCI-DISINFECTANT is effective against Pseudomonas aeruginosa.

Disinfection in Institutions, Industry, Schools and Homes

For disinfecting floors, walls, bedrooms, countertops, tables, chairs, garbage pails, bathroom fixtures and other hard non-porous surfaces.

Add 2 oz of PCI-DISINFECTANT to 4 gallons of water. Apply to previously cleaned hard non-porous surface with mop or cloth.

At 2 oz / 4 gallon use level PCI-DISINFECTANT is effective against Staphylococcus aureus, Salmonella choleraesuis and Trichophyton interdigitale (the athlete's foot fungus).

Disinfection of Barber Tools

Pre-cleaned barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a 1/2 oz / gallon solution of PCI-DISINFECTANT.

Disinfection of Poultry Equipment, Animal Quarters and Kennels.

Poultry brooders, watering fountains, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 2 oz PCI-DISINFECTANT in 4 gallons of water with a mop, cloth or brush. Small utensils should be immersed in this solution.

Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals, or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

