

**PRECAUTIONARY  
STATEMENTS  
HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS  
WARNING**

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get on your skin or clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of children and pets. Do not use on humans, household pets, or livestock. Do not permit children or pets to go into sprayed area until spray has completely dried.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Do not apply directly to lakes, streams and ponds. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

**PHYSICAL OR CHEMICAL  
HAZARDS**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

Container Disposal: Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other approved State and local procedures.

# TOXAPHENE

## E-8

Contains 8 Lbs. Toxaphene Per Gallon

**ACTIVE INGREDIENTS:**

Toxaphene\* ..... 72%  
Petroleum Distillate ..... 20%

**INERT INGREDIENTS:** ..... 8%

**TOTAL** ..... 100%

\* Technical Chlorinated Camphene (Chlorine content 67-69%)

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

If swallowed- Call a physician immediately. Gastric lavage is indicated if material was taken internally. DO NOT INDUCE vomiting unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled- Remove victim to fresh air. Apply respiration if indicated.

If on skin- Wash immediately with soap and water.

If in eyes- flush with plenty of water.

NET CONTENTS GALLON

Gal. (Litres)

EPA Reg. No. 44215-52

EPA Est. No.

P. O. Box 285  
Lansing, Kansas 66043

**KAW  
VALLEY**

**FRUIT AND NUT CROPS** Rates given in pints per 100 gallons unless specified. Aircraft use equivalent rate in 10-50 gallons water per acre.

**APPLES** 1 pint, red banded leafroller. Do not apply after second cover spray or within 40 days of harvest. Do not exceed 16 pints per acre.

**CITRUS** (GRAPEFRUIT, KUMQUATS, LEMONS, LIMES, ORANGES, TANGELOES, TANGRINES) 3-4 pints, grasshopper. Apply to trunk and main branches. Do not apply within 30 days of harvest. Do not feed treated forage to dairy animals or animals being finished for slaughter. Do not use over 10 pints per acre.

**CRANBERRIES** 3 pints, grasshopper. 4 pints, armyworm. Do not apply after fruit begins to form.

**PEARS** 1 pint, red banded leafroller, 1.2 pints, pear psylla. Do not apply within 40 days of harvest or after second cover spray. Do not exceed 16 pints per acre.

**STRAWBERRIES** 2-3 pints per acre, cutworm (climbing only), strawberry crown borer, strawberry weevil. Do not apply after fruit begins to form.

**FORAGE, FIELD, SEED AND VEGETABLE CROPS.** Rates given in pints per acre. Do not exceed recommended rates. Ground rig use 15-100 gallons water per acre. Aircraft 5-50 gallons water per acre.

**ALFALFA** (Seed crop only): 2½-3½ pints, alfalfa looper, alfalfa weevil, armyworm, 1½-3½ pints, climbing cutworm, 3 pints, alfalfa caterpillar, pea aphids, webworm, 2 pints, clover leaf weevil, Lygus bug, grasshopper, 1½ pints, spittle bug. Do not feed treated forage to dairy animals or animals being finished for slaughter.

**BARLEY, OATS, WHEAT:** 2 pints, armyworms, cutworms, grasshopper. Do not feed treated forage to dairy animals or animals being finished for slaughter. No restriction on the use of the grain.

**BEANS** (Dry Green, Lima) 4 pints, armyworms, Mexican bean beetle, bean leaf beetle, cucumber beetle, blister beetle, flea beetle, cutworm, Lygus bug, bean leaf skeletonizer, garden webworm, pea weevil, 3 pints, corn earworm, grasshopper, serpentine leafminer, 1 pint, bean leafroller, bean leafhopper, 1.6 pints, thrips. No limitation on use of shelled beans as human food. Do not apply sprays to green or snap beans after pods begin to form. Do not feed treated vines to dairy animals or animals being finished for slaughter.

**BRUSSELS SPROUTS, CAULIFLOWER** 4 pints, cabbage loopers, cutworm, diamondback moth, 3 pints, flea beetle, 2 pints, green peach aphid, 1.6 pints, leafminer, 1.6 pints, climbing cutworms, fall armyworm, thrips. Do not apply within 30 days of harvest or after edible parts start to form.

**CABBAGE** 2-2½ pints to control insects listed under Broccoli. Do not apply after heads start to form.

**CARROTS** 3 pints, cutworm, flea beetle, 2 pints, grasshopper, 4 pints, Lygus bug. No time limitation. Do not feed treated foliage to dairy animals or animals being finished for slaughter.

**CELERY** 1.6 pints, cutworms, fall armyworms, green peach aphid, serpentine leafminer, thrips, 4.5 pints, celery leafhopper. Do not apply after celery plants start to bunch or after plants are half mature.

**CORN** 2-4 pints, fall armyworm and corn earworm feeding in whorl, 3 pints, chinch bug, cutworm, flea beetle, grasshopper, 2½ pints, webworm. Do not feed treated forage to dairy animals or animals being finished for slaughter in fields treated late in season. Do not graze dairy animals. Do graze meat animals on treated stover within four weeks of slaughter.

**COTTON** 2-4 pints, armyworm, boll weevil, bollworm, cotton leafworm, cutworm (climbing), flea beetle, grasshopper, Lygus bug, thrips, 4 pints, stink bug, 3 pints, flea hopper. Do not graze dairy animals or animals being finished for slaughter in fields treated late in season. Do not graze treated Cotton, or Feed gin waste to dairy animals or animals being finished for slaughter.

**EGGPLANT, PEPPERS:** 2-3 pints, armyworm, blister beetle, Colorado potato beetle, cutworm, flea beetle, leafhopper, serpentine leafminer, 3 pints, cucumber beetle. Do not apply within 5 days of harvest.

**LETTUCE** 4 pints, grasshopper, field cricket, 3 pints, cucumber beetle, flea beetle, 2 pints, armyworm, cutworm, 3-5 pints, cabbage looper. Do not apply after seedling stage-leaf lettuce. Do not apply after heads begin to form-head lettuce.

**ONIONS** 2 pints, thrips. Do not apply to green or spring onions.

**PARSNIPS** 1½ pints, grasshopper, striped flea beetle. Do not use tops for feed or food.

**RICE** 2 pints, armyworm, chinch bug, cutworm, grasshopper, rice stink bug. Do not feed treated forage to dairy animals or animals being finished for slaughter. No limitation on use of grain.

**RUTABAGAS** 2 pints, cutworm, 4 pints, flea beetle. Do not use treated tops for feed or food.

**SOYBEANS** 2-4 pints, armyworm, bean leaf beetle, blister beetle, corn earworm, grasshopper, velvetbean caterpillar, 2 pints, cricket, 1½ pints, flea beetle. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry. Do not apply within 21 days of harvest.

**SPINACH** 2 pints, armyworm, cutworm, green peach aphid, thrips, 1.6 pints, leafminer. Do not apply within 21 days of harvest. Do not apply more than once per season.

**TOMATOES** 2.5 pints, armyworm, blister beetle, cabbage looper, Colorado potato beetle, cutworm, flea beetle, pinworm, russet mite, serpentine leafminer, tomato pinworm, tomato hornworm, 3-5 pints, thrips, 4 pints, grasshopper. Do not apply within 30 days of harvest.