

1 of 1
 Page 2
 REG. # 44215-74

VITO SPOT FUNGICIDE

WETTABLE POWDER

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Do not apply directly to water or wetlands. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Cover or incorporate spills. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or waste is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide, or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

RESTRICTED USE PESTICIDE

Because of the high acute toxicity of triphenyltin hydrosulfide, and its potential for creating teratogenic effects, this product may be applied only by certified applicators or persons directly under their supervision.

A FUNGICIDE FOR CONTROL OF:

EARLY & LATE BLIGHT OF POTATOES LEAFSPOT OF SUGAR BEETS
 SCAB & CERTAIN OTHER DISEASES ON POTATOES

ANGER
PELIGRO



**DANGER: KEEP OUT OF REACH OF CHILDREN
 SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS**

POISON

STATEMENT OF PRACTICAL TREATMENT

- If on skin:** Wash with plenty of soap and water.
- If inhaled:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.
- If in eyes:** Flush with plenty of water. Call a physician.
- If swallowed:** Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician or Poison Control Center.

ACTIVE INGREDIENT:
 Triphenyltin hydrosulfide 47.5%
INERT INGREDIENTS: 52.5%
TOTAL 100.0%

*Equivalent to 15.5% Hezalic Tin

EPA Reg. No. 44215-74

EPA Est. No.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PREPARE: Before using VITO SPOT FUNGICIDE TO THE SPRAY TANK, PARTIALLY FILL WITH WATER. Cut or tear excess bagging along dotted line. Remove water soluble handle and place in partially filled tank while agitating. Empty the sprayer only after use. Do not reuse. After emptying thoroughly, and remaining water in spray tank. Continuous agitation within the spray tank is essential. Wash out spray tank thoroughly after use and dispose of washings by pouring away from water supplies.

PLANT	CONCENTRATION	CRASS RATE	SPRAYER	REMARKS
POTATOES	1 lb. per 100 gal. water	12 to 24 gal.	Apply to foliage during early stages of infection. Repeat 2 to 4 times at 7 to 10 day intervals. Do not use on plants with severe leaf damage. Do not use on plants with severe leaf damage.	Do not spray when plants have been treated in spray. Do not spray early or late relative to treated area.
SUGAR BEETS	1 lb. per 100 gal. water	4 to 10 gal.	Apply to foliage during early stages of infection. Repeat 2 to 4 times at 7 to 10 day intervals. Do not use on plants with severe leaf damage. Do not use on plants with severe leaf damage.	Do not spray when plants have been treated in spray. Do not spray early or late relative to treated area.
See precautions.				
Do not spray within 7 days of harvest when applied to produce which will be consumed in its natural state. Do not spray within 30 days when applied to produce which will be consumed after processing. Do not use on produce which will be consumed in its natural state. Do not use on produce which will be consumed after processing.				

ACCEPTED
 JUN 26 1985
 Federal Insecticide,
 Fungicide and
 Rodenticide Act
 Registered under
 Reg. No. 44215-74

BEST AVAILABLE COPY