

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
& DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing. Do not contaminate feed or foodstuffs.

ENVIRONMENTAL HAZARDS

Use of this product in a manner not in accordance with the information on this label may injure domestic animals, wildlife, aquatic life or produce other undesirable results.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with labeling.

PESTICIDE DISPOSAL

Pesticide, spray mixture or residue that cannot be used according to label instructions must be disposed of according to Federal, state or local procedures or for the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL

Triple rinse (or equivalent) and offer for recycling or reconditioning. Or dispose of in a sanitary and fill or by incineration if permitted by State and local authorities.

PRODUCT INFORMATION

STAY ON LIQUID SULFUR is a fixable sulfur formulation that may be applied as a general purpose application. Consult your State Agricultural Experiment Station or Extension Specialist for advice in selecting treatments from this label to best fit local conditions.

USE ONLY ON CROPS LISTED AND ONLY AS SPECIFIED ON THIS LABEL. EXCEPT ON CAUTION, do not apply when temperatures exceed or are likely to exceed 90° F. Do not use sulfur with oil or within 14 days of an oil spray (for citrus 21 days). Do not combine with emulsifiable salts.

NOTE: Sulfur will cause severe fruit and leaf injury to sulfur sensitive crops. Do not apply or allow to drift to apricots, Almond and Citrus peaches, strawberries, cucurbits, cucumber, cantaloupes, melons, squash, fibberts, spinach, tung trees, walnuts, or other sensitive plants. Sulfur may burn foliage and fruit during periods of high temperatures and under certain climate conditions.

STAY ON LIQUID SULFUR can be mixed and applied with equal parts of water.

DO NOT STOP AGITATION WITH SULFUR IN SPRAY EQUIPMENT. SULFUR IS HIGHLY CORROSIVE AND EQUIPMENT SHOULD BE CLEANED THOROUGHLY AFTER EACH DAY'S SPRAYING.

RECOMMENDATIONS

Unless otherwise specified for specific crop, dosage rates are given as parts of STAY ON LIQUID SULFUR per 100 gallons water for use with backpack sprayer. The total amount of STAY ON LIQUID SULFUR per acre should be 2.5-4 parts depending on the size of the trees. Because of variations in the types of spray equipment, label dosage rates are given in terms of spray volume. For example, a pump sprayer with a 100 gallon tank and a 1000 gpm nozzle should use 2.5 parts of STAY ON LIQUID SULFUR per 100 gallons of spray per acre. The dosage rate for backpack sprayers should be 20 to 40 parts per acre spray per acre, depending on the type of sprayer.

APPLE, PEAR

Scab: 1.5 parts per 100 gallons of water. 1.5 parts per 100 gallons of water. 1.5 parts per 100 gallons of water. 1.5 parts per 100 gallons of water.

PLUM, MANGO

Scab: 1.5 parts per 100 gallons of water. 1.5 parts per 100 gallons of water. 1.5 parts per 100 gallons of water. 1.5 parts per 100 gallons of water.

CHERRY, PLUM, PRUNE

Scab: 1.5 parts per 100 gallons of water. 1.5 parts per 100 gallons of water. 1.5 parts per 100 gallons of water. 1.5 parts per 100 gallons of water.

**STAY-ON
LIQUID SULFUR**

ACTIVE INGREDIENT

Sulfur as S

INERT INGREDIENTS

TOTAL

This product contains 6 lbs. of sulphur per gallon.

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

IF SWALLOWED Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Never give anything by mouth to an unconscious person.

IF ON SKIN - Wash thoroughly with soap and water.

IF IN EYES Wash eyes with plenty of clear water for at least 15 minutes.



PESTICIDE SERVICE CONSULTANTS

ALLIANCE, OHIO 44601

EPA EST NO 37429 GA 01

Net Volume

5 U.S. Gallons Liquid (1)

AC
Fung
regis
EPA

EPA REG

STAY-ON LIQUID SULFUR



INGREDIENTS	
ACTIVE INGREDIENTS	52%
INERTS	48%
TOTAL	100%

This product contains 6 lbs. of sulphur per gallon

KEEP OUT OF THE REACH OF CHILDREN
CAUTION
ALLOWED - Drink 1 or 2 glasses of water and induce vomiting by rubbing back of throat with finger. Never give anything by mouth to an unconscious person.
SKIN - Wash thoroughly with soap and water.
EYES - Wash eyes with plenty of clear water for at least 15 minutes.

PESTICIDE SERVICE CONSULTANTS
ALLIANCE, OHIO 44601

ACCEPTED
JUN 13 1981
U.S. Federal Insecticide, Fungicide, and Rodenticide Act
Registered at
EPA Reg. No.

37429 GA 01 EPA REG NO. 44283-3
Volume 5 U.S. Gallons Liquid (18.927L)

CITRUS - Rest Mite, cover mite.
Apply sprays November thru May.
THRIPS - Early spring treatment for the control of the pest on citrus. Apply 1 to 2 quarts per acre in sufficient water for thorough coverage. Make applications after sunrise and before 4 to 6 p.m. Repeat if necessary. Do not apply within 21 days of harvest.

PEACH, NECTARINE
Powdery Mildew, brown rot, fruit rot.
Pink and Brown Sprays.
Petiole Blight, leaf curl, silver leaf.
Application to mature trees: 1 to 3 quarts per acre.

FIELD AND VINE CROPS
Unless otherwise specified for specific crops, applications are made at 1 quart of STAY-ON LIQUID SULFUR per acre. It is applied by ground equipment at a rate of sufficient water to provide thorough coverage.

COTTON - Anilic Mite, 2 quarts per acre. Cotton Rootworm, 1 quart per acre. Cotton Leafhopper, 1 quart per acre.

GRAPES - Powdery Mildew. Use 1 to 2 quarts per 100 sq. ft. of water. For 1 acre, use 1 to 2 quarts of STAY-ON LIQUID SULFUR per acre. Make first application when shoots are 6 to 8 inches long, second when shoots are 12 to 15 inches long, and third at late bloom stage. The timing, rate and number of applications vary with soil conditions. Consult State Agricultural Extension Service or State Agricultural Experiment Station for detailed information. Follow the USE PRECAUTIONS on this label.

PEANUTS - For rust, use 1 to 4 quarts per acre of STAY-ON LIQUID SULFUR in sufficient water for good coverage. Begin 40 to 60 days after planting, use 1 to 4 quarts every 10 to 15 days as needed to control rust.

POTATOES - Powdery Mildew, use 3 to 4 quarts per acre. Repeat every 10 to 14 days. Apply at the first sign of the disease and repeat at 1 week intervals.

SUGAR BEETS - Early blight, late blight, downy mildew, root rot. Use 1 to 2 quarts of water per acre by air at first sign of infection. Use 4 quarts per acre of the product. For 1 acre, use 1 to 2 quarts per acre. Repeat applications should be made at 10 to 30 day intervals or as the necessary. The amount of water used should be 10 to 15 gallons per acre. Applications will be needed.

STRAWBERRIES - Powdery Mildew, use 1 quart. Apply at the first sign of infection and repeat at 1 week intervals. For Strawberry root rot, use 1 quart per acre at the first sign of infection and repeat at 1 week intervals with a fungicide containing chlorothaloxyl.

TOMATOES - Late blight, early blight, Septoria, stem rot, leaf mold, fruit rot. Use 1 to 2 quarts per acre.

RECOMMENDATIONS: DIRECTIONS FOR USE.

For a complete list of crops and pests controlled by this product, see the label on the container. For a complete list of crops and pests controlled by this product, see the label on the container.

ARTICHOKE	CABBAGE	DELPHINIUMS	LIMES	ROSES
ALMONDS	CALENDULA	EVERGREENS	ORANGES	SNAPDRAGONS
ASPARAGUS	CAMELIAS	FIGS	PEAS	STRAWBERRIES
AZALEAS	CARNATIONS	GOOSEBERRIES	PEPPERS	SUGAR BEETS
BEANS	CAULIFLOWER	GRAPEFRUIT	PHLOX	SWEET PEAS
BEETS	CHRYSANTHEMUMS	HEATHER	PLUMS	TANGELINES
BLACKBERRIES	COLLARDS	HOLLYHOCKS	POTATOES	TANGELINES
BLUEBERRIES	CORN	KALE	PRUNES	WALNUTS
BROCCOLI	CURRANTS	LEMONS	RASPBERRIES	
BRUSSELS SPROUTS	DAHLIAS	LILACS	RHODODENDRONS	

BEANS - For rust, use 1 to 4 quarts per acre of STAY-ON LIQUID SULFUR in sufficient water for good coverage. Begin 40 to 60 days after planting, use 1 to 4 quarts every 10 to 15 days as needed to control rust.

ROSES - Powdery Mildew, use 1 quart. Apply at the first sign of infection and repeat at 1 week intervals.

CHIGGERS - For chigger control, use 1 to 2 quarts per acre of STAY-ON LIQUID SULFUR in sufficient water for good coverage.

SOIL PH CORRECTIVE - For soil pH correction, use 1 to 2 quarts per acre of STAY-ON LIQUID SULFUR in sufficient water for good coverage.

WARRANTY STATEMENT

Buyer agrees to pay for the product and to use it in accordance with the label directions.