

BEEBLE BAIT



*Kills Darkling Beetles
and Hide Beetles
and Their Larva Stages*

EPA REG. NO. 44313-18

EPA EST NO. 2393-IL-3

NET WT. 30 lbs.

ACCEPTED

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the purposes
registered under
EPA Reg. No. 44313-18

ACTIVE INGREDIENT	
ORTHO BORIC ACID	5%
INERT INGREDIENTS	95%
TOTAL	100%

SPECIMEN

KEEP OUT OF THE REACH OF CHILDREN

— CAUTION —

STATEMENT OF PRACTICAL TREATMENT
IMMEDIATELY CONTACT PHYSICIAN OR POISON CONTROL CENTER IF
SWALLOWED. IF EYE CONTACT OCCURS, FLUSH EYES WITH PLENTY
OF WATER. GET MEDICAL ATTENTION IF IRRITATION PERSISTS.
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS PRECAUTIONARY STATEMENTS

CAUTION

HARMFUL IF SWALLOWED. WASH HANDS THOROUGHLY AFTER HANDLING. MAY CAUSE
EYE IRRITATION. AVOID EYE CONTACT. WEAR A DUST MASK OR RESPIRATOR.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry place. Do not store where children or animals may gain access.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Manufactured For

R Value, Inc.

P.O. Box 1144

Decatur, GA 30030-1144

(404) 371-0820

APPLICATION INSTRUCTIONS

For Darkling and Lesser Mealworm Beetle and
Hide Beetle Control in Poultry Houses
(Adult and Larvae)

For Poultry Houses in which birds are grown on litter.

Prior to application of bait, remove birds. Apply bait uniformly to floor of poultry house or to old litter with a fertilizer/seed spreader, at the rate of 4-6 per 1000 square feet, in bands along feeder lines. Spread fresh litter of at least four inches uniformly over all treated area (floor and old litter). Then introduce birds. Reapply after each grow-out, if needed, in the sequence outlined above.

Example: A poultry house measures 30' by 500'. There are two feeder lines per house. A seed spreader has a ten foot pattern. 500' x 10' = 5,000 square feet per feeder line or 10,000 square feet per house. At the rate of 6 pounds per thousand square feet, multiply 6 times 10 or 60 pounds of Beetle Bait per house.

For Poultry Houses in which birds are grown in cages [layer or high-rise (pit type) houses]:

Birds do not have to be removed prior to application of bait. Apply bait uniformly to floor of poultry house with a fertilizer/seed spreader, at the rate of 4-6 pounds per 1000 square feet, as needed, to maintain control. If facility is washed down and sanitized using standard cleaning and disinfecting procedures, bait must be reapplied to continue beetle control.

Suggested 5% Boric Acid Bait Label, EPA Reg. No. 44313-18
Poultry House Use
(revised 12-19-89)

APPLICATION INSTRUCTIONS¹

For Darkling and Lesser Mealworm Beetle and Hide Beetle
Control in Poultry Houses (Adults and Larvae)

For Poultry Houses in which birds are grown on litter:

Prior to application of bait, remove birds. Apply bait uniformly to floor of poultry house or to old litter with a fertilizer or seed spreader, at the rate of 4-6 pounds per 1000 square feet², in bands along feeder lines. Spread fresh litter of at least four inches uniformly over all treated area (floor or old litter). Then introduce birds. Reapply after each grow-out, if needed, in the sequence outlined above.

For Poultry Houses in which birds are grown in cages [layer or high-rise (pit type) houses]:

Birds do not have to be removed prior to application of bait. Apply bait uniformly to floor of poultry house with a fertilizer or seed spreader, at the rate of 4-6 pounds per 1000 square feet², as needed, to maintain control. Do not apply directly to manure. If facility is washed down and sanitized using standard cleaning and disinfecting procedures, bait must be reapplied to continue beetle control.

¹This text allows application of product on litter without removal of old litter. Since new litter separates the birds from the product, the use is non-food. Without this intervening layer of new litter, the applicant must petition the Agency for an exemption from a tolerance.

²Rate may have to be adjusted, depending upon the effectiveness review.

Poultry House 30 x 500 • Two 8' Bands x 500 =
1000 x 2 = 2000 sq ft. at 6 lb per 1000 =
6 x 2 = 12 lbs bait