

TIFCHEM SEVIN* 10 Dust

(INSECTICIDE)

NET WEIGHT 50 LBS.

Active Ingredient	By Wt.
Carbaryl (1-naphthyl N-methylcarbamate)	10%
Inert Ingredients	90%

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER
BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE

May be applied up to harvest unless otherwise noted. This product is formulated for application by ground or air equipment.

ALFALFA CLOVERS: COWPEAS PEANUTS SOYBEANS B. Star Beetle, Mexican Bean Beetle—5 to 10 lbs. per acre. Alfalfa Caterpillar, Bean Leaf Beetle, Cucumber Beetle, Green Cloverworm, Japanese Beetle, Leafhoppers, Three-cornered Alfalfa Hopper, Thrips, Velvetbean Caterpillar—10 lbs. per acre. Armyworms, Corn Earworm, Shink Bugs, Webworms—10 to 15 lbs. per acre. Some leaf injury may occur if applications are made when tender foliage is wet.

BEANS: Mexican Bean Beetle—5 lbs. per acre. Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Velvet Bean Caterpillar, Flea Beetles, Japanese Beetle—10 lbs. per acre. Corn Earworm, Armyworms, Tarnished Plant Bug, Shink Bugs—10 to 15 lbs. per acre. In California for Corn Earworm, Lima Bean Pod Borer, Lygus and Shink Bugs—20 lbs. per acre.

BLUEBERRIES: Blueberry Maggot, Cherry and Cranberry Fruitworms, Japanese Beetles—15 to 20 lbs. per acre. Apply 3 weeks before harvest and repeat 10 days later or as necessary.

CABBAGE BROCCOLI BRUSSELS SPROUTS CALL FLOWER KOHLEWART: Flea Beetles, Harlequin Bug—5 to 10 lbs. per acre. Armyworms, Corn Earworm, Yellow Turnip Moth—10 to 20 lbs. per acre. Do not apply within 3 days of harvest.

CORN: Corn Earworm—Application of this product to the tassels, beginning during the pollen shed period will seriously reduce bee populations. Corn Rootworm, Anjou, European Corn Borer, Fall Armyworm, Sap Beetles, Japanese Beetle, Leafhoppers, Flea Beetles—15 to 20 lbs. per acre. For insects attacking silks and ears apply at 2 to 3-day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. May be applied up to harvest.

COTTON: Early-Season—Thrips, Cotton Fleahopper—8 to 15 lbs. per acre, as soon as insects appear. Mid and Late-Season—Boll Weevil, Bollworm, Pink Bollworm, Cotton Leafworm, Lygus Bugs, Cotton Leaf Perforator—20 lbs. per acre. Apply when insects appear and repeat at 5 to 7 day intervals. Aphid populations will usually be suppressed when this insecticide is used on a regular schedule of treatments.

CUCUMBERS MELONS PUMPKIN SQUASH: Pickworm, Melonworm—10 to 15 lbs. per acre. Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs—10 lbs. per acre. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use SEVIN on watermelons in Florida.

FILBERTS: Filbert Aphid, Filbert Leaf Roller—40 lbs. per acre. Apply when Leaf Roller eggs are hatching. Filbertworm—40 lbs. per acre. Apply when adult moths first appear and repeat 3 to 4 weeks later. Do not apply within 60 days of harvest.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid Breathing Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Contact. Wash Hands and Face Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily. Note for Physician: Carbaryl is a moderate reversible Cholinesterase inhibitor. Atropine is antidotal. Do not store or transport near feed or food. AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

GRAPES: Grape Leaf Folder, Grape Leafhopper, Grape Leaf Skeletonizer—10 to 20 lbs. per acre. Apply just before first brood leaf folder larvae emerge from rolls and as needed for leafhoppers. Grape Berry Moth, Japanese Beetle, June Beetles, Red-Banded Leaf Roller—20 lbs. per acre.

GRASSHOPPERS: Alfalfa, Clovers, Cowpeas, Peanuts, Soybeans, Rice, Forage Grasses, Pasture, Milg and Grain Sorghum, Sugar Beets—5 to 15 lbs. per acre. Apply 5 to 10 lbs. for nymphs on small plants or sparse vegetation in wasteland, rangeland, ditch-banks and borders. Use 10 to 15 lbs. when grasshoppers are mature or material is applied to crops requiring greater coverage. Do not apply to Rice or Sugar Beets within 14 days of harvest, and to Sorghum Grain within 21 days of harvest.

LETTUCE CARROTS: Flea Beetles, Harlequin Bug, Leafhoppers—5 to 10 lbs. per acre. Six-spotted Leafhopper—10 to 15 lbs. in a 5 to 7-day schedule. Armyworms, Cabbage Caterpillars, Corn Earworm, Spittlebugs, Tarnished Plant Bugs, Lygus Bugs, Shink Bugs—10 to 20 lbs. per acre. For best results in the control of Corn Earworm after etuce heads begin to form use 20 lbs. in a 5 to 7-day schedule. May be used up to harvest on carrots. Do not use on Head Lettuce within 3 days of harvest and to Leaf Lettuce within 14 days.

PEACHES: Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Cat-Facing Insects, Periodical Cicada, Codling Moth—35 to 40 lbs. per acre. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. May be applied the day before harvest.

PLUMS PRUNES: Fruit Tree Leaf Roller, Peach Twig Borer, Plum Curculio, Red-Banded Leaf Roller, Codling Moth, Eastern Tent Caterpillar, Orange Tortix, Western Tussock Moth—20 to 30 lbs. per acre. Apply when insects appear and repeat as necessary. Make full coverage application. Do not apply within 1 day of harvest.

POTATO EGGPLANT PEPPER: Colorado Potato Beetle, Flea Beetles, Leafhoppers—5 to 10 lbs. per acre. Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Lygus Bugs, Shink Bugs, Lace Bugs, Tarnished Plant Bugs—10 to 20 lbs. per acre.

*Trade Mark of Union Carbide Corporation for the active ingredient 1-naphthyl N-methylcarbamate

RICE FORAGE GRASSES PASTURE: Armyworms, Shink Bugs, Thrips—10 to 15 lbs. per acre. For optimum thrip control in grasses grown for seed, direct dust into soil. Do not apply to Rice within 14 days of harvest. May be applied up to harvest on Pasture and Forage Grasses.

SORGHUMS: (Including Milg and Grain Sorghum, Annual Sorghum, Corn Earworm, Shink Bugs, Webworms—10 to 20 lbs. per acre. Sorghum Midge—15 lbs. per acre. Direct to the forming heads. Treatment for sorghum midge control should be made 3 to 4 days after heads have emerged from boot. Do not apply to Sorghum Grain within 21 days of harvest.

STRAWBERRIES: Meadow Spittlebug, Strawberry Leaf Roller, Strawberry Weevil—10 to 20 lbs. per acre. Do not apply within 1 day of harvest.

TOBACCO: Hand Dusting for Budworms and Tobacco Flea Beetles—4 to 6 lbs. per acre when hand dust applications are made on individual plants. Direct dust to the bud and top leaves only. Heavier application may cause plant injury. Rotary dusters are recommended. Shake cans and cloth bags usually result in excessive applications. Broadcast Application for Hornworms, Budworms, Japanese Beetles, Grasshoppers, Flea Beetles—15 to 25 lbs. per acre depending on size of plants. Use highest rate only on large plants. Full coverage is essential. Begin treatment when worms are small and repeat as needed. Heavier applications may cause plant injury. Late applications are necessary, treat immediately after pruning. Do not apply within 5 days of pruning or 15 days of cutting. Avoid getting wet with plant juices when cutting the crop. Direct dust to the bud and top leaves only.


TOMATO: Flea Beetles, Leafhoppers, Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer—20 to 40 lbs. per acre.

DOGS CATS: Brown Dog Ticks, Fleas—Dust liberally over the animal and rub thoroughly into the skin. Apply also to the feet, between the toes and on the legs. Apply at weekly intervals if needed. Avoid treatment of kittens and puppies less than 4 weeks of age.

IN OR AROUND DOG AND CAT QUARTERS: Fleas—Dust liberally in and around sleeping quarters, kennels and dog houses by means of a clean, hand-operated duster. Force dust into cracks and crevices. Repeat as necessary, but not more often than once weekly.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



EPA REG. NO. 10873-23
EPA EST. NO. 10873-GA-1

TIFTON CHEMICAL COMPANY

TIFTON, GEORGIA