

# STAF

## HOSPITAL SPRAY DISINFECTANT

**Bactericidal—Staphylocidal  
Pseudomonocidal  
Eliminates Odors—Fungicidal  
Virucidal against HIV-1  
(Aids Virus)\***

**ACTIVE INGREDIENTS:**

n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chloride.....0.1%  
n-Alkyl (68% C12, 32% C14) dimethyl ethylbenzyl ammonium chlorides .....0.1%  
Isopropanol .....50.7%

**INERT INGREDIENTS:**.....49.1%

EPA Reg. No. 44446-20

EPA Est. No. 44446-TX-2

**KEEP OUT OF REACH OF CHILDREN**

**WARNING:**

See Additional Precautionary Statements on Back Panel

**NET WT. 11 OZ. (312 g)**

Manufactured By  
Quest Chemical Corporation  
Houston, Tx. 77041

**ACCEPTED  
APR 26 1993**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 44446-20

**STAF**

**HOSPITAL SPRAY DISINFECTANT**

This product has been designed specifically for Homes, Hospitals, Nursing Homes, Institutions, Offices, Schools, Motels & Hotels as a disinfectant deodorant & air refresher. It is recommended for disinfection of telephones, bed springs, door knobs, linen carts, hampers, wheelchairs, light switches, garbage pails & other clean hard non-porous inanimate surfaces. It will also remove foul musty odors from automobile airconditioning systems caused by mildew and other odor producing organisms. Powerful deodorizers keep the automobile interiors smelling fresh and clean.

In addition, this product deodorizes those areas which generally are hard to keep fresh smelling (such as garbage storage areas, empty garbage bins & cans, restrooms, shower stalls & other areas which are prone to odors caused by micro-organisms). This product may also be applied directly to ash trays, food spills, etc., in automobiles to eliminate odors from these sources. When used as directed, this product demonstrates effective disinfection against Staphylococcus aureus, Salmonella choleraesuis, & Pseudomonas aeruginosa & Trichophyton mentagrophytes.

This product is fungicidal against Pathogenic fungi (Athletes Foot Fungus), when used as directed on hard surfaces found in the bathroom, shower stalls & locker rooms or other clean hard non-porous surfaces commonly contacted by the bare feet.

**MOLD & MILDEWSTAT:** This product controls mold & mildew & the odors they cause on hard non-porous inanimate surfaces when used as directed.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Wrap container & put in trash collection.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**SURFACE DISINFECTION & DEODORIZATION:** (1) First remove soil deposits from surface to be treated. (2) Hold can six inches from surface & spray until completely wet, allow the surface to remain wet for 10 minutes.

**AIR DEODORIZER & AIR FRESHENER:** Point upwards to center of room & spray for three seconds.

**MILDEWSTAT:** To control mold and mildew on hard non-porous surfaces (such as floors, walls, table tops) spray the surface to be treated, making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth reappears.

For use in automobile air conditioning systems, with engine running and air conditioner on maximum speed and cooling, spray the product into the air intake vent. Hold can eight to twelve inches above the vent while spraying four to six seconds. Make sure the spray tip is pointed directly at the intake vent before spraying. The recirculating system will pick up the product and the entire system will be deodorized in seconds. Shut off engine and air conditioner after applying the product.

\*Kills HIV-1 (AIDS virus) on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS).

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS) OF SURFACES/ OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

**Personal Protection:** Disposable latex or vinyl gloves, gowns, face masks, or eye coverings as appropriate, must be worn during all cleaning of body fluids, blood and decontamination procedures.

**Cleaning procedures:** Blood and body fluids must be thoroughly cleaned from surfaces and objects before application of disinfectant.

**Contact Time:** Effective against HIV-1 (AIDS virus) on hard non-porous surfaces in the presence of a moderate amount of organic soil (5 % blood serum) providing 1800 ppm of active quaternary water for a contact time of 10 minutes at room temperature (20-25° C). Use a ten (10) minute contact time for disinfection against all other bacteria and fung. claimed (listed on labeling).

**Disposal of Infectious Materials:** blood and other body fluids should be autoclaved and disposed of according to federal, state, and local regulations for infectious waste disposal.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**WARNING:** Keep out of reach of children. Causes eye & skin irritation. Do not get in eyes, skin or on clothing. Harmful if swallowed. Flammable. Contents under pressure. Do not use near fire, sparks or flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

**STATEMENT OF PRACTICAL TREATMENT**

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove & wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution or if these things are not available, drink large quantities of water. Call a physician. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Messed against circulatory shock, respiratory depression and convulsion may be needed.

ITEM 328