

44446-22 PM23 1062  
DIRECTIONS FOR USE (Cont'd)

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**CAUTION**

**HARMFUL IF SWALLOWED.** Do not inhale spray drift. This liquid can cause skin irritation. Avoid contact with skin, eyes and clothing. Do not wear contaminated clothing.

In case of contact wash with soap and water. For eyes, flush with large amounts of water immediately and get medical attention.

**ENVIRONMENTAL HAZARDS**

Do not apply this liquid to areas where the roots of desirable plants such as trees or ornamentals will come into contact with it as injury might result. Do not use on lawns, driveways, walks or similar areas. Avoid spray drift to susceptible plants, as this product may injure cotton, beans, peas, etc. Do not contaminate domestic water or irrigation water and keep from contact with fertilizers, insecticides, fungicides and seeds. Vapors from this product may injure susceptible plants in the immediate vicinity. Coarse sprays are less likely to drift.

This product is toxic to fish and wildlife. Do not apply directly to water. Birds and other wildlife in treated areas may be killed. Apply only as specified on the label.

Do not contaminate water by cleaning of equipment, or disposal of wastes.

Thoroughly clean spray equipment with a suitable chemical cleaner such as a butyl cleaner before using for other purposes.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is a liquid, non-selective, persistent herbicide designed for use where control of all vegetation is desired. It is recommended for non-crop areas such as fence rows, parking lots around buildings, loading ramps, storage yards, tank farms, along pipelines, industrial plant sites and other similar areas where grasses and weeds interfere with drainage, create a fire hazard or cause unsightly appearance.

**DOOM RTU  
WEED KILLER**

**ACTIVE INGREDIENTS:**

Petroleum Distillate ..... 95.39%  
Isooctyl ester of 2,4-dichlorophenoxyacetic acid\* ..... 1.09%

Bromacil (5-bromo-3-sec-butyl-6-methyluracil) ..... 0.98%

**INERT INGREDIENTS** ..... 2.54%

**TOTAL** ..... 100.00%

\*2,4-D equivalent 0.70%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See Side Panel For Additional Precautionary Statements

Manufactured By

**QUEST**

QUEST CHEMICAL CORPORATION  
15102 SOMMERMEYER  
HOUSTON, TEXAS 77041

Net Contents:      Gallon(s)

EPA Reg No. 44446-22

EPA Est. 44446 TX-1

This liquid weed killer is composed of 5-bromo-3-sec-butyl-6-methyluracil (bromacil) and a low volatile 2,4-D ester in a phytotoxic oil. The oil carrier combines the properties of low volatility with rapid penetration and translocation in the plant. The insolubility of the herbicide in water assures full effect regardless of rainfall after application or stage of growth at the time of application.

This liquid produces rapid top kill of all succulent vegetation. Translocation and root absorption of component herbicides produce extensive root kill of weeds and provide long-term residual control for one year.

This liquid may be applied at any time of the year except when the ground is frozen. Best results are obtained after predominant weeds have emerged and developed to a point where they present a problem. The product should be applied as is. No agitation or dilution is necessary and any equipment from sprinkler can to complicated sprayer may be used. Equipment should be calibrated so as to assure application of the recommended amount of material.

This liquid weed killer when used according to directions will provide low cost, convenient and complete weed control for a season or more. Sprayed vegetation is no more flammable than vegetation destroyed by mowing or other agrichemical means. Because of the taste and odor, it is unlikely that sprayed vegetation will be eaten by livestock.

**Recommendations for Use:** For annual weeds and grasses such as crabgrass, foxtail, barnyard grass, lambsquarter and pigweed use 1/2 to 1 gal. per 500 sq. ft. or 47 to 83 gallons per acre.

Perennial weeds and grasses such as smooth broome, quackgrass, Canada bluegrass, broomsedge, aster, dandelion, goldenrod, platan, wild carrot and purple top use 1/2 to 1 gal. per 250 sq. ft. or 87 to 174 gallons per acre.

**STORAGE AND DISPOSAL**

See attached  
Statement

**ACCEPTED**  
JUN 7 1985  
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 44446-22

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Do not contaminate water, food or feed by storage or disposal.

TRIPLE rinse or equivalent. Then, offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.