

44446-62

6/19/2001

1/1

READ ENTIRE LABEL BEFORE EACH USE FOR PET USES; USE ONLY ON DOGS PRECAUTIONARY STATEMENTS Hazards To Humans & Domestic Animals CAUTION

Keep children and/or pets out of treated areas during treatment. Do not apply this product to or near children's toys. Avoid spraying in eyes. Avoid breathing spray mist. Avoid contact with skin. Wash hands thoroughly after use. Vacate room after treatment and ventilate before reoccupying. Avoid contamination of feed and foodstuffs. Cover or remove fishbowls. This product is not for use on humans. ANIMALS: Do not spray directly in/on eyes, mouth or genitalia of pets. Use only on dogs. Do not use on cats or kittens. Do not use on puppies less than twelve (12) weeks old. Consult a veterinarian before using this product on debilitated, aged, pregnant, medicated or nursing animals. Sensitivities may occur after using any pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Environmental Hazards

This product is toxic to fish. Do not apply to any body of water. Do not contaminate water when disposing of equipment washwaters.

Physical Or Chemical Hazards

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Storage: Store in a cool, dry area. Do not transport or store below 32°F.

Disposal: Wrap container in several layers of newspaper and dispose of in trash. Can may be disposed of with other recyclables where local law permits. Do not incinerate or puncture.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not use in the edible product areas of food/feed processing plants, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held. Not for use in Federally Inspected Meat and Poultry Plants.

In the home, cover all food handling surfaces and cover or remove all food and cooking utensils, or wash thoroughly after treatment. Exposed food should be covered or removed.

NONFOOD AREAS: Multi-Purpose Insect Spray may be applied in nonfood/feed areas. Nonfood/feed establishment areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, mop closets and storage (after canning or bottling).

SHAKE WELL BEFORE USING

Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches. Remove birds and cover fish aquariums before spraying.

Spray until surface is just damp, but not to the point of wetness or run-off. Avoid excess application.

HOUSEFLIES, FRUIT FLIES, MOSQUITOES, GNATS, WASPS, HORNETS, BEES, BLACK FLIES, SKIPPER FLIES, BLOW FLIES AND SMALL FLYING MOTHS: Close all doors and windows and spray room at the rate of 2 to 3 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Upon reentry, open windows and ventilate for at least 2 hours. Repeat treatment once a month.

COCKROACHES, WATERBUGS, PALMETTO BUGS, SILVER-FISH, FIREBRATS, CRICKETS, SCORPIONS, SPIDERS, MILLIPEDES, CENTIPEDES, SOWBUGS AND PILLBUGS: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray when ever possible. Upon reentry, open windows and ventilate for at least 2 hours. Repeat treatment once a month.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat treatment once a month.

INDIAN MEAL MOTHS, ANGOUIMOIS GRAIN MOTHS: Close doors and windows, direct spray into all parts of room especially around stored food product containers, pellets and damaged areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Upon reentry, open windows and ventilate for at least 2 hours. Repeat treatment once a month.

EXPOSED STAGES OF CONFUSED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, RED FLOUR BEETLES, RICE WEEVILS, GRAMMARY WEEVILS, GRAIN MITES: Spray into cracks and crevices, processing and handling equipment, around cartons or containers or other areas where these insects tend to congregate. Spray 30 seconds for each

100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Upon reentry, open windows and ventilate for at least 2 hours. Repeat treatment once a month. Infested stored food products in the home should be disposed of.

CARPET BEETLES: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment once a month.

BEDBUGS: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Allow all sprayed articles to dry thoroughly before reuse. Repeat treatment once a month.

TO CONTROL CLOTHES MOTHS AND CARPET BEETLES on wool and other proteinaceous fibers: Remove infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated by spraying lightly, keeping nozzle at least three feet from fabric to avoid staining. Allow all sprayed articles to dry thoroughly before use.

IN ANIMAL QUARTERS AND MILK ROOMS: For rapid control of House Flies, Mosquitoes, Gnats, Wasps, Hornets, Bees and Small-Flying Moths. Remove animals prior to application. Close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not reenter treated area and ventilate after treatment is completed. Upon reentry, open windows and ventilate for at least 2 hours. Repeat treatment once a month.

TO TREAT LIVESTOCK: To protect beef cattle, dairy cattle, goats, sheep, hogs and horses from attacks of Stable Flies, Horse Flies, Deer Flies, Face Flies, House Flies, Horn Flies, Mosquitoes, and Gnats. Spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from Face Flies, spray the face and head, but do not spray into eyes. Repeat treatment when flies are troublesome.

FOR SWINE: Do not ship animals for slaughter within 5 days of last treatment.

TO CONTROL BLOOD SUCKING LICE: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

TO CONTROL POULTRY LICE, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

FOR CONTROL OF BEDBUGS AND MITES IN POULTRY HOUSES, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

FLEAS AND TICKS: (ADULTS AND LARVAE) Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed, entire floor and floor covering, with special attention to cracks and crevices around baseboards, and window and door frames. Infested upholstered furniture may also be treated with this product. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use.

TO CONTROL FLEAS AND TICKS ON DOGS. Starting at the tail, stroke against the lay of the hair, spraying the parted fur in short bursts, directly behind the hand to insure penetrating the coat. Dampen the fur up to the head. Repeat application every two weeks if necessary. TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8 to 10 inches. Spray each square foot for three seconds. Spray only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all sprayed articles to dry thoroughly before use.

FOR THE CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray, as well as Whiteflies, Aphids, exposed Thrips, Spiders, Red Mites and Leafminers. May be used on plants, as well as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwood, English Ivy, euonymus, fuchsia, geraniums, crassula, grape vine, Kerkia palm, laurel, marigolds, rhododendrons, roses, rubber plants, serapongans, stocks, wandering Jew and zinnias.

Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lady Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Army Worms. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

Notice: Buyer assumes all responsibility for safety and use not in accordance with directions.

INGREDIENTS: Water (CAS# 7732-18-5), Propano-Isobutane (CAS# 68476-85-7), Petroleum Distillate (CAS# 8002-05-9), Permethrin (CAS# 52645-53-1) HHS RATINGS: HEALTH-1, FLAMMABILITY-2, REACTIVITY-1, PERSONAL-1

Multi-Purpose Insect Spray

- \* CONTAINS THE SYNTHETIC PYRETHROID PERMANONE®
\* CONTROLS NAMED RESIDENTIAL, INDUSTRIAL AND GARDEN INSECTS
\* FOR USE ON DOGS AND ON PREMISES TO CONTROL FLEAS AND TICKS.
\* AN AQUEOUS STABLE MICROEMULSION
\* CONTROLS PESTS FOR UP TO FOUR WEEKS
\* KILLS LICE AND THEIR EGGS
\* FOR USE ON BEDDING, FURNITURE AND OTHER INANIMATE OBJECTS INFESTED WITH LICE.
\* NOT FOR USE ON HUMANS
\* FOR USE ON LIVESTOCK AND THEIR PREMISES
\* CONTROLS FLEAS AND TICKS ON DOGS FOR 14 DAYS

ACTIVE INGREDIENT:

\*Permethrin 0.25%

OTHER INGREDIENTS: 99.75%

100.00%

1(3-phenoxyphenyl)methyl (+/-) cis/trans 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate
Cis/Trans ratio: Min. 35% (+/-) cis and max. 65% (+/-) trans.
PERMANONE - Registered Trademark of AgrEvo Environmental Health, Inc.

EPA REG. NO. 44446-62 EPA EST. NO. 44446-TX-1

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID: IF INHALED: Remove affected person to fresh air. Apply artificial respiration if indicated. IF IN EYES: Flush with plenty of water. Contact a physician if irritation persists. IF ON SKIN: Wash affected areas immediately with soap and water. Get medical attention if irritation persists.

See Side Panel for Additional Precautionary Statements

NET WT. 11 OZ. (312 g)

Manufactured by QUEST Chemical Corporation Houston, Texas 77041

ACCEPTED JUN 19 2001 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 44446-62