

44446-70

05-17-2004

44

MAY 17 2004

QUEST CHEMICAL CORPORATION
12255 FM 529
Northwoods Industrial Park West
Houston, TX 77041

Dear Ms Sharp:

Subject: Label Amendment
Product Name: QUEST WILD GAME INSECTICIDE
EPA Registration No. 44446-70
Application date: March 18, 2004


The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable, and a stamped copy is enclosed for your records.

Sincerely,



Richard Gebken
Acting Product Manager 10
Insecticide Branch
Registration Division 7505

2/4



Houston, Texas 77041

QUEST WILD GAME INSECTICIDE

- A CONCENTRATED PYRENONE® MICRO EMULSION
- CLEAR – STABLE – LEAVES NO OILY OR OBJECTIONABLE RESIDUES
- EFFECTIVE, FAST ACTING INSECTICIDE
- KILLS AND REPELS BITING FLIES
- FOR USE IN HOMES, INSTITUTIONS, RESTAURANTS, FOOD PROCESSING PLANTS, DAIRIES, LIVESTOCK AND OUTDOORS FOR MOSQUITO CONTROL FOR CONTROL OF FLEAS AND BROWN DOG TICKS ON PREMISES AND ON PETS
- APPLY WITH CONVENTIONAL MECHANICAL OR COMPRESSED AIR EQUIPMENT
- MULTI-PURPOSE
- MULTI-BUG KILLER
- KILLS CRAWLING & FLYING INSECTS ON CONTACT
- CONTAINS [PYRETHRIN] [INSECTICIDE] DERIVED FROM FLOWERS
- WATER BASED
- PLANT SPRAY BUG KILLER
- HOUSE & GARDEN INSECT KILLER
- FLYING & CRAWLING INSECT KILLER
- KILLS FLEAS, TICKS & OTHER LISTED INSECTS ON GAME ANIMALS

ACTIVE INGREDIENTS:

Pyrethrins	0.10%
*Piperonyl Butoxide, Technical	1.00%
OTHER INGREDIENTS	98.90%
	100.00%

*Equivalent to min. 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds.

EPA REG. NO. 44446-70 EPA Est. No. 44446-TX-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

If inhaled: <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing: <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes: <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed: <ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Do not give any liquid to the person. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

In case of medical emergencies or health and safety inquiries or in case of fire, leaking or damaged containers, information may be obtained by calling 1-877-315-9819.

READ ENTIRE LABEL BEFORE EACH USE
PET USE APPLIES TO USE ONLY ON DOGS AND CATS

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contamination of feed and foodstuffs. Remove pets, birds and cover fish aquaria before space spraying or surface applications.

Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing animals. Do not use on dogs or cats under 12 weeks.

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. For terrestrial use, do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

CONTAINERS LARGER THAN ONE GALLON:
Do not contaminate water, food or feed by storage or disposal.
Pesticide Storage and Spill Procedures: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste. Protect from freezing.
Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or an approved waste disposal facility.
Container Disposal: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other approved state and local procedures.

See Side Panel For Additional Precautions.

CONTAINERS ONE GALLON OR LESS:

STORAGE AND DISPOSAL

STORAGE: Store in a cool dry area away from heat or open flame.
DISPOSAL: Do Not Puncture or Incinerate! If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.

ACCEPTED

MAY 7 7 2004

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 44446-70

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL USE

INDOOR APPLICATION:

The use of this product in processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned with potable water before using.

Vacate treated areas and ventilate before reoccupying.

For maximum effectiveness, a combination of spot, surface, treatment and fogging is recommended.

SURFACE SPRAYING: To control *Cockroaches, Palmetto Bugs, Ants, Silverfish, Spiders, Crickets, Clovermites, Cheese Mites, Earwigs, Sowbugs, Boxelder Bugs, Centipedes, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Sawtoothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drugstore Beetles, Meal Worms, Grain Mites and Cadellles:* Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers, behind and beneath sinks, stoves, refrigerators and cabinets, along baseboards, around door and window frames, into dark corners and in and around garbage compactors, garbage chutes and cans. For *Silverfish*, spray bookcases. For *Ants*, spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

In case of infested foods in cabinets, dispose of the infested material. Remove all contents and clean thoroughly. Spray interior surfaces of empty cabinets thoroughly. After spray has dried, cover shelves with paper or otherwise protect foods from direct contact with sprayed surfaces.

TO CONTROL CARPET BEETLES: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

TO CONTROL BEDBUGS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

PET AREAS: To control *Fleas* and *Brown Dog Ticks* on premises: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals is recommended. Repeat treatment as needed.

TO CONTROL CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

SPACE SPRAYING: Close doors, windows and transoms and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of *House Flies, Fruit Flies, Gnats, Mosquitoes, Skipper Flies, Wasps, Hornets, Angoumois Grain Moths, Mediterranean Flour Moths, Indianmeal Moths* and *Tobacco Moths*, direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3-6 ounces per 1000 cubic feet of space. Do not remain in treated areas. If insects have not dropped to floor in 3 minutes, repeat spraying as quantity sprayed was insufficient. Ventilate the area after treatment is completed. To aid in the prevention of breeding, spray on standing water and in garbage pails.

FOGGING: To kill exposed or accessible stages of the above named crawling insects, use a rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

ON LIVESTOCK: To protect cattle, horses and hogs from *Horn Flies, House Flies, Mosquitoes* and *Gnats* apply a light mist sufficient to wet the surface of the hair.

TO CONTROL STABLE FLIES, HORSE FLIES AND DEER FLIES, apply at a rate of 2 ounces per adult animal sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

TO CONTROL BLOOD-SUCKING LICE, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

TO CONTROL POULTRY LICE, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

FOR CONTROL OF BEDBUGS AND MITES in poultry houses, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

FLEAS AND TICKS ON DOGS AND CATS: Start spraying at the animal's head, avoiding the eyes, nose and mouth as you spray. Make sure the animal's entire body is covered including the legs, under body and tail. While spraying, fluff the hair so that the spray will penetrate the skin. Make sure spray wets thoroughly. Avoid contact with genitalia. Repeat application every week, if necessary.

ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where *Mosquitoes* may rest (low, marshy places, stagnant pools or ponds and shore lines). Apply at a rate of 2.5 pints per acre (0.1 ounce per 100 square feet). Using truck-mounted conventional misting equipment such as Rotomist® models 51C, 91 or 100HT traveling at 5 m.p.h. to treat a 300 foot swath calibrate equipment to deliver 0.9 gallon per minute.

OUTDOORS: (Camping, fishing, hunting) - Spray outer clothing and surroundings. Repels *Biting Flies, Mosquitoes, Midges* and *Gnats*. If sprayed every hour or so, will aid in keeping surroundings free from *Mosquitoes*. If in rare cases skin irritation results, avoid continued contact. Do not pour or squirt - use a good sprayer which will throw a heavy fine mist.

FOR CONTROL OF Fleas, Ticks, Biting Flies, Horse Flies, Deer Flies, Stable Flies, Houseflies, Gnats, Mosquitoes and Lice on carcasses of wild game animals including but not limited to deer, feral hogs, javelina, mule deer and elk. Apply product with fine spray mist wetting but not saturating hair on the carcass. Begin at the head of the animal and apply over the entire body covering all exposed areas. Allow 5 to 10 minutes for control of listed insects to occur. Repeat as necessary to kill listed insects prior to field dressing carcass.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CONSUMER LABELING:

SHAKE WELL BEFORE USE

TO KILL-

CRAWLING INSECTS: *Cockroaches, Silverfish, Sowbugs, Ants, Crickets, Earwigs, Grain Mites, Clovermites, Spiders, Boxelder Bugs, Centipedes.* Turn sprayer nozzle to stream position and spray directly into cracks, crevices and other hiding places along baseboards, hitting as many bugs as possible. Repeat as necessary. Keep about three feet away from walls, fabrics and furniture.

FLYING INSECTS: *Flies, Houseflies, Mosquitoes, Gnats, Mediterranean Flour Moths, Small Flying Moths.* Turn sprayer nozzle to spray position. Keep about three feet away from walls, fabrics and furniture.

WASPS & HORNETS: Turn sprayer nozzle to stream position. Spray directly on insects whenever possible. Apply a wet spray after dark when all insects have returned to their nests.

4/4

GARDEN INSECTS ON HOUSE PLANTS AND ORNAMENTAL PLANTS OUTDOORS: Turn sprayer nozzle to spray position. Direct spray so that upper and lower leaf surfaces are contacted. To control *Aphids, Bagworms, Thrips, Flea Beetles, Japanese Beetles, Lace Bugs, Leaf Hoppers, Leafhoppers, Mimosa Webworm Larvae, Plant Bugs* and *Whiteflies* use fine mist spray about 18 to 24 inches from infested plants. Use caution on new growth and tender foliage. Repeat treatment as necessary. Avoid spraying plants when temperature exceeds 85°F. Spray when air is still. Avoid heavy wetting.

FLEAS AND TICKS ON DOGS AND CATS: Turn sprayer nozzle to spray position. Start spraying at the animal's head, avoiding the eyes, nose and mouth as you spray. Make sure the animal's entire body is covered including the legs, underbody and tail. While spraying, fluff the hair so that the spray will penetrate the skin. Make sure spray wets thoroughly. Avoid contact with genitalia. Repeat application every week if necessary.

PET AREAS: To control *Fleas* and *Brown Dog Ticks* on premises: Turn sprayer nozzle to spray position. Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals is recommended. Repeat treatment as needed.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Quest Chemical Corporation. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: QUEST CHEMICAL CORPORATION MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Quest Chemical Corporation is authorized to make any warranties beyond those contained herein. Quest Chemical Corporation disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT QUEST CHEMICAL CORPORATION'S ELECTION, THE REPLACEMENT OF PRODUCT.

- * **ALTERNATE FRONT PANEL PLUGS FOR CONSUMER TYPE LABELING:**
 - * **LEAVES NO [LINGERING] [OBJECTIONABLE] [INSECTICIDE] ODOR**
 - * **LEAVES NO OBJECTIONABLE OILY OR STICKY RESIDUE**
 - * **FOR CONTROL OF FLEAS AND BROWN DOG TICKS**
 - * **FOR USE ON PETS**
 - * **CAN BE USED ON PETS**
 - * **KILLS AND REPELS [BITING] FLIES**
 - * **EFFECTIVE**
 - * **WORKS FAST**
 - * **FAST ACTING**
 - * **LEAVES NO OILY STAINS**
 - * **[LEAVES] NO STICKY MESS**
 - * **UNSCENTED**
 - * **NON-AEROSOL**
 - * **EASY TO USE TRIGGER PUMP**
 - * **CONVENIENT TRIGGER PUMP**
 - * **NO AEROSOL FUMES**
 - * **KILLS WITH PYRETHRIN [INSECTICIDE]**
 - * **PYRETHRIN [INSECTICIDE] IS MADE FROM FLOWERS**
 - * **FOR USE ON PLANTS AND SHRUBS**
- BRACKETS "[]" DENOTE OPTIONAL COPY

Pyrenone is a registered trademark of the Aventis Group.

QUEST CHEMICAL CORPORATION
12255 FM 529
HOUSTON, TEXAS 77041

