

P. 15 44 116-15 1 of 11

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460	EPA REGISTRATION NO.	DATE OF ISSUANCE
	44711-10	AUG 28 1989
	TERM OF ISSUANCE	
NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input checked="" type="checkbox"/> REREГИSTRATION <i>(Under the Federal Insecticide, Fungicide and Rodenticide Act, as amended)</i>	NAME OF PESTICIDE PRODUCT	
	bearing insecticide	

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

1. The registrant shall submit to the Agency, in accordance with the Act, all data required by the Act for registration/reregistration of any pesticide under the Act, 3(e)(2) when the Agency requires all registrants of similar products to submit such data.

2. The registrant shall submit to the Agency, in accordance with the Act, all data required by the Act for registration/reregistration of any pesticide under the Act, 3(e)(2) when the Agency requires all registrants of similar products to submit such data.

3. The registrant shall submit to the Agency, in accordance with the Act, all data required by the Act for registration/reregistration of any pesticide under the Act, 3(e)(2) when the Agency requires all registrants of similar products to submit such data.

- a. The label shall include the registration no. 44711-10.
- b. The label shall include the statement "Not for Use in USA Meat and Poultry Plants." Therefore, either add the statement "Not for Use in USA Meat and Poultry Plants," or submit an application to USA requesting use in these areas. If granted, you can then indicate on the label that the product can be used in these areas.

BEST AVAILABLE COPY

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL _____ DATE 8/28/89

57484: I:CR-31:LeMaster:LR-15:KENCO:08/18/89:10/02/89:CL:sw:ek:de
 R:57487:CR-31:LeMaster:LR-15:KENCO:08/23/89:10/05/89:AS:SN:VO:aw

- c. Standardize the first two paragraphs under Indoor Use Areas [i.e., "Do not use in edible . . . (after canning or bottling)"] as follows:

Do not use in food areas of food handling establishments, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Nonfood areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

- d. Change the phrase "Active Ingredients" to the singular.
- e. Add a footnote to the label ingredient statement indicating:
- Contains petroleum distillates.
- f. Change "Manufactured By" to "Manufactured For."
- g. Since your data matrix does not give a dielectrical breakdown voltage, you must add the following to the use directions:
- Do not use this product in or on electrical equipment due to the possibility of shock hazard.
- h. Add the following statement to the bedbug directions:
- Allow mattresses and other articles to dry thoroughly before use.
- i. Under the "As a Space Spray" heading, include the following statement:

EXTINGUISH ALL FLAMES AND PILOT LIGHTS.

- j. Under Precautionary Statements. replace "Full-faced gas mask with canister is recommended for . . ." with the following:

For space spraying/fogging indoors, wear a NIOSH or MSHA approved respiratory protection device.

3. Storage stability studies must be performed using containers similar to the ones you will be using in the trade. You indicate on the application form that you will be using plastic containers in the trade. However, the proposed storage stability study indicates that "amber glass containers" will be used. The proposed study is therefore unacceptable.

4. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosures

Fearing Insecticide Concentrate 10% E.C.

A long lasting livestock and premise spray that provides knock down, broad spectrum insecticidal effectiveness, and excellent residual activity for up to 28 days.

Kills Gypsy Moth Caterpillars, Japanese Beetles, Roaches, Ants, Fleas, Ticks, Lice and numerous other insects as listed on side panels.

For use outdoors and in homes and non-food areas of commercial buildings, institutions, warehouses, theaters, office buildings, schools, motels, hotels, restaurants food/feed processing establishments, hospitals, and kennels.

RECOMMENDED FOR AGRICULTURAL/COMMERCIAL USE

ACTIVE INGREDIENTS:

*Permethrin	10.0%
INERT INGREDIENTS:	90.0%
TOTAL	100.0%

* (3-phenoxyphenyl) methyl (±) cis/trans 3-(2,2-dichloroethenyl) 2,2-dimethyl cyclopropanecarboxylate

Cis/trans ratio: Max. 65% (±) trans and min. 35% (±) cis.
Contains 0.75 lbs. of permethrin per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Rear Panel for Additional Precautionary Statements

Manufactured By:
Fearing Manufacturing Co., Inc.
P. O. Box 708
So. St. Paul, Minnesota 55075

EPA Est. No. 44716-MN-2

EPA Reg. No. 44716-

ACCEPTED
with COMMENTS
in EPA Letter Dated:

NET CONTENTS _____ OZ.

AUG 28 1989

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide
registered under EPA Reg. No.

44716-10

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Keep container sealed when not in use. Protect from freezing. Shake or stir product thoroughly before using or reusing diluted emulsion.

To prepare dilutions, the concentrate should first be stirred or agitated well. Add the required amount of concentrate to water and blend thoroughly. Do not hold dilutions for more than 24 hours.

For maximum effectiveness, a combination of localized application and space treatment is recommended. Remove pets, birds and cover fish aquariums before spraying.

INDOOR USE AREAS


Do not use in edible product areas of food/feed processing plants, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving area while food is exposed.

NON FOOD/FEED AREAS include (but not limited to) garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling).

BAKERIES	HOSPITALS	RAILROAD CARS
BEVERAGE PLANTS	HOTELS, MOTELS	RESTAURANTS
CANNERIES	INDUSTRIAL INSTALL.	SCHOOLS
FLOUR MILLS	KENNELS	SHIPS' HOLDS
FOOD PROCESSING PLANTS	MEAT PACKING PLANTS	SUPERMARKETS
GRAIN ELEVATORS	HOMES	TRUCK TRAILER
GRANARIES	OFFICE BUILDINGS	WAREHOUSES

AS A SURFACE SPRAY TO CONTROL:

Cockroaches, Waterbugs, Palmetto Bugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Millipedes, Centipedes, Sowbugs, Pillbugs, Clover mites, Cheese mites, Granary weevils, Rice weevils, Confused flour beetles, Rust red flour beetles, Sawtoothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, mealworms, grain mites and cadelles:



For initial clean up of severe insect infestation, dilute at a rate of (1) part concentrate in 19 parts water (6.7 fluid ounces per gallon) (0.5%).

For normal infestations dilute (1) part concentrate in 100 parts water (1.33 ounces per gallon) (8 teaspoonful per gallon) (0.1%).

Use a good sprayer and adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, and behind shelves and drawers. Spray bookcases for silverfish. Spray ant trails, nests and points of entry. If surface application only is to be used, spray floors, walls and other surfaces applying at a rate of (1) gallon to 750 square feet of surface.

TO CONTROL: Carpet Beetles, spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.


TO CONTROL: Bedbug, spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

TO CONTROL: Fleas and ticks (adult and larvae), thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment.

AS A SPACE SPRAY:

For initial clean up of severe insect infestation, dilute at a rate of (1) part concentrate in 19 parts water (6.7 fluid ounces per gallon) (0.5%).

For normal infestations dilute (1) part concentrate in 100 parts water (1.33 ounces per gallon) (8 teaspoonful per gallon) (0.1%).



Apply with mechanical or compressed air equipment (non-thermal) adjusted to deliver a fine mist. Close doors and windows and shut off ventilating systems. For rapid control of house flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, bees, black flies, Angoumois grain moths and tobacco moths, direct spray at an upward angle distributing it uniformly through the entire area at a rate of (1) ounce per 1000 cubic feet of space. Keep area closed for at least 10 minutes. Vacate areas after treatment and ventilate and sweep up dead insects before occupying.

For rapid kill of exposed or accessible stages of other insects named on this label: Apply using conventional mechanical or compressed air equipment (non-thermal), following directions for space spraying. Apply the above dilutions at a rate of (1) ounce per 1000 cubic feet of space.

OUTDOOR USE AREAS

CORRALS
FEED LOTS
GARDENS
PARKS

DRIVE-IN THEATERS
DRIVE-IN RESTAURANTS
GOLF COURSES
ZOOS
PLAYGROUNDS


RECREATIONAL AREAS
SWINE LOTS
URBAN AREAS
HOMES

For initial clean up of severe insect infestation, dilute at a rate of (1) part concentrate in 19 parts water (6.7 fluid ounces per gallon) (0.5%).

For normal infestations dilute (1) part concentrate in 100 parts water (1.33 ounces per gallon) (8 teaspoonful per gallon) (0.1%).

TREATMENT OF FLYING INSECTS

TO AID IN CONTROL OF: Mosquitoes, Wasps, Gnats, Skippers, treat walls, ceilings, moldings, screens, door frames, window frames, beams, light cords and similar resting places.



FOR TEMPORARY REDUCTION OF ANNOYANCE
FLIES, GNATS, MOSQUITOES

In drive-in restaurants, drive-in theaters, golf courses, urban areas, parks, playgrounds and other recreational areas. Dilute (1) part concentrate in 40 parts water (3.2 fluid ounces per gallon of water) (0.25%). Direct the spray into tall grass, shrubbery and around lawns, where these pests may hover or rest. Apply while the air is still and avoid wetting foliage. Application should be made prior to occupancy. Repeat as necessary.

In the treatment of corrals, feedlots, swine lots and zoos, cover any exposed water, drinking fountains and animal feed before application. Fill the area with mist and direct the spray into tall grass, shrubbery and around lawns where the above pests may hover or rest. Apply while the air is still and avoid wetting foliage. In zoos, prevent exposure of reptiles to the product. Repeat as necessary.

GYPSY MOTH CATERPILLARS AND ADULTS, ADULT JAPANESE BEETLES,
EASTERN AND FOREST TENT CATERPILLARS

To kill on contact; Dilute (1) part concentrate in 40 parts water (3.2 fluid ounces per gallon of water) (0.25%), apply to insure thorough coverage of upper and lower surfaces.

ON TREES AND SHRUBS: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.


ON ROSES AND IN FLOWER GARDENS: Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

ANIMALS AND ANIMAL PREMISE USE

CATTLE BARNS
SWINE HOUSES
MILK ROOMS

POULTRY HOUSES
HORSE BARNS
KENNELS

DAIRIES
MILKING PARLORS



For initial clean up of severe insect infestation, dilute at a rate of (1) part concentrate in 19 parts water (6.7 ounces per gallon) (0.5%).

For normal infestations dilute (1) part concentrate in 100 parts water (1.33 ounces per gallon) (1/2 teaspoonful per

LIVESTOCK SPRAY

(Dairy & Beef Cattle, Sheep, Goats, Swine, Horses, Poultry)

For Swine: Do not ship animals for slaughter within 5 days of last treatment.

To Control; horn flies, house flies, mosquitoes and gnats: Apply to wet the hair thoroughly, with particular attention to topline, underline, flanks, withers and other infested areas. Repeat treatment at intervals of 5 to 12 days for small insect populations or as needed when flies are emerging in large numbers.

To Control; stable flies, horse flies and deer flies: Apply a quart per adult animal to wet the hair thoroughly with particular attention to the legs, flanks, barrel, topline and other body areas commonly attacked by these flies. Repeat daily as needed.


To Control; face flies: Apply using spray which produces large wetting droplets. Apply to the face of the animal in the morning before releasing to pasture. Apply sufficiently to wet the face but not more than 1 1/2 ounces per animal. Repeat daily as needed.

To Control; poultry lice: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

To Control; bedbugs and mites in poultry houses: Spray crevices of roost poles, cracks in walls and cracks in nests where the bedbugs and mites hide.

To Control; sheep "tick" or ked: Thoroughly wet all portions of the body by dipping or spraying with sufficient pressure and with a nozzle adjustment to give penetration of the wool. Treat at a rate sufficient to wet the animal.

To Control; fleas, ticks, lice and Mange mites (adults & larvae) on dogs, and livestock: Thoroughly wet the animal by dipping, sponging or spraying. Allow animal to dry in a warm place without rinsing or toweling. Gives long lasting protection against reinfestation and applications at intervals of two to three weeks. Repeat applications as necessary. DO NOT TREAT PUPPIES LESS THAN FOUR WEEKS OLD.



BEST AVAILABLE COPY

To Control; fleas, flies ticks lice and Mange mites (adults & larvae) in kennels, dog houses, runs, and yards: Thoroughly wet pest breeding or resting areas to run off. Most efficient pest control is accomplished when heavy pressure and good contact is achieved. Timing and frequency of application should be based on pest populations reaching nuisance levels, but accompanying manure removal and sanitation practices should precede sprays. Do not spray feed, food or water. Repeat as needed but not more often than once every two weeks.

IN CATTLE BARNS, HORSE BARNS, SWINE HOUSES, KENNELS, MILKING PARLORS, MILK ROOMS, DAIRIES AND POULTRY HOUSES: To control flies, mosquitoes and gnats.

For initial clean up of severe insect infestation, dilute at a rate of (1) part concentrate in 19 parts water (6.7 fluid ounces per gallon) (0.5%)

For normal infestations dilute (1) part concentrate in 100 parts water (1.33 ounces per gallon) (8 teaspoonful per gallon) (0.1%).

Apply as a fog or fine mist, directing the spray toward the ceiling and upper corners until the area is filled with mist. Use about 2 ounces per 1000 cubic feet of space. For best results, close doors and windows before spraying and keep closed for ten to fifteen minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

BEST AVAILABLE COPY


STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local



CONTAINERS ONE GALLON AND SMALLER: Do not reuse container. Wrap and discard in trash.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Remove and wash clothing before reuse. Avoid contamination of food and feed.

Full-faced gas mask with cannister is recommended for general protection when applying indoors as a space spray or fogger.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Contact a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove person to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with plenty of soap and water.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater/dip solution.

PHYSICAL HAZARDS

Do not use or store near heat or open flame.

BEST AVAILABLE COPY

