

DIRECTIONS FOR USE

TO BE USED IN ANIMAL FEED ONLY

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

PRESERVATION OF FEED GRAINS

NOTE TO USER: Grain may spoil if instructions are not followed carefully.

To insure complete coverage of the grain, apply the product with a pressure sprayer system. Determine the moisture level of the harvested feed grain, establish applicator grain throughput and set flow to apply the proper amount of Protex III[®] liquid preservative as follows. (Be sure to correct flow meter setting for viscosity.)

Grain Moisture %	Protex III [®] Weight %	Lbs Protex III [®] Per Ton of Grain
20 or less	0.7%	14
21 - 30	0.9%	18
31 - 35	1.2%	24
36 - 40	1.4%	28

If grain temperature is above 70 F when treated, increase the above rates by 0.1% or 2 pounds per ton for each 10 F above 70 F. Grain should be treated within 6 hours after harvest. Recommended rates are suitable for storage under roof for up to 1 year.

Do not mix treated grain with untreated grain in the same storage facility. Good harvesting and storage practices should always be followed in the handling, treatment and storage of Protex III[®] treated high moisture grain. Do not treat corn or other cereal grains which might be used for seed, planting purposes or human consumption. Treated corn and other cereal grains are to be used for animal feed only.

PRESERVATION OF HAY TO BE STORED IN BALES.

NOTE TO USER: Hay may spoil if instructions are not followed carefully.

Spray Protex III[®] completely over fresh forage prior to baling. Store in well ventilated barn or shed. The following application rates should be used.

Hay Moisture %	Lbs Protex III [®] Per Ton of Wet Hay
20 - 24	10
25 - 29	20
30 - 35	30

Do not store treated hay in contact with dry untreated hay.

PRESERVATION OF SILAGE OR HAY CROP SILAGE:

NOTE TO USER: Silage may spoil if instructions are not followed carefully.

Apply 20 lbs. Protex III[®] preservative per ton of fresh forage by metering the preservative into the blower housing continuously as the forage is blown into upright silos, or by metering the preservative into the blower housing of the forage harvester if silage is to be stored compacted in bunkers or pits. Follow customary best practices for moisture levels, compaction of forages.

In either of the above applications the Protex III[®] preservative should be diluted by adding an equal volume of water to improve coverage and make metering easier, but such dilution is not necessary for good results.

Consult your Protex III[®] Representative for specific recommendations on how to avoid spoilage.

PRO

LIQUID PRESERVATIVE
WHEAT, OATS, BARLEY,

ACTIVE INGREDIENTS:
Propionic Acid 67%
Formaldehyde 10%
INERT INGREDIENTS: 23%
Contains Methanol

TOTAL: 100%

This product contains 8.6 pounds of Protex III[®] per gallon.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
(& DOMESTIC ANIMALS)**

DANGER

Use in well ventilated area. Do not inhale vapor. Wear goggles, rubber gloves and protective clothing when handling Protex III.

Fatal if swallowed, inhaled or absorbed through skin. Corrosive, causes eye and skin damage. Do not breathe vapor, dust or spray mist. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling.

Protex III[®] is a Trademark of American Agricultural Products

AMERICAN AGRICULTURAL PRODUCTS
36 N. HURON ST. • YPSILANTI, MI 48197

PROTEX III[®]

LIQUID PRESERVATIVE for HIGH MOISTURE CORN, SORGHUM,
WHEAT, OATS, BARLEY, GRASS FORAGE and LEGUME FORAGE

ACTIVE INGREDIENTS:	
Propionic Acid	67%
Formaldehyde	10%
INERT INGREDIENTS:	23%
Contains Methanol	
TOTAL:	100%
This product contains 8.6 pounds of Protex III [®] per gallon.	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (& DOMESTIC ANIMALS)

DANGER

Use in well ventilated area. Do not inhale vapor. Wear goggles, rubber gloves and protective clothing when handling Protex III.

Fatal if swallowed, inhaled or absorbed through skin. Corrosive, causes eye and skin damage. Do not breathe vapors, dust or spray mist. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling.

Protex III[®] is a Trademark of American Agricultural Products Inc.

KEEP OUT OF REACH OF CHILDREN DANGER (CAUSES SEVERE BURNS)

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED:** May be fatal or cause blindness if swallowed. Do not induce vomiting. Give large amounts of water followed by milk or egg white beaten with water. Get immediate medical attention. Cannot be made nonpoisonous.
- IF INHALED:** Vapor harmful. Get medical attention.
- IF ON SKIN:** Immediately flush with plenty of water for at least 15 minutes.
- IF IN EYES:** Immediately flush with plenty of water for at least 15 minutes. Get immediate medical attention.

STORAGE AND DISPOSAL

STORAGE: Do not store, pour, spill near heat or open flame.

DISPOSAL:

- PROHIBITIONS:** Do not contaminate water, food, or feed by disposal. Open dumping is prohibited.
- PESTICIDE DISPOSAL:** Pesticide spray mixture or residue that cannot be used should be disposed of according to federal, state, or local procedure under the Federal Insecticide, Fungicide, and Rodenticide Act.
- CONTAINER DISPOSAL:** Triple rinse or equivalent and offer for recycling or reconditioning, or dispose of in a sanitary landfill or by other approved state and local procedure.
- GENERAL:** Consult federal, state or local disposal authorities for approved alternative procedures.

EPA Est. No. 44718-MI-1
EPA Reg No. 44718-3

AMERICAN AGRICULTURAL PRODUCTS, INC.
36 N. HURON ST. • YPSILANTI, MI 48197 • PHONE: (313) 484-1773

CORROSIVE

WARNING: Avoid contact with skin, eyes, or clothing. Do not breathe vapors. Wash thoroughly with soap and water. If you experience any symptoms, call your doctor immediately. For more information, call 1-800-4-A-AG-1.

TEX III

HIGH MOISTURE CORN, SORGHUM,
GRASS FORAGE and LEGUME FORAGE

KEEP OUT OF REACH OF CHILDREN
DANGER (CAUSES SEVERE BURNS)

STATEMENT OF PRACTICAL TREATMENT


- IF SWALLOWED:** May be fatal or cause blindness if swallowed. Do not induce vomiting. Give large amounts of water followed by milk or egg white beaten with water. Get immediate medical attention. Cannot be made nonpoisonous.
- IF INHALED:** Vapor harmful. Get medical attention.
- IF ON SKIN:** Immediately flush with plenty of water for at least 15 minutes.
- IF IN EYES:** Immediately flush with plenty of water for at least 15 minutes. Get immediate medical attention.

STORAGE AND DISPOSAL

STORAGE:
Do not store, pour, spill near heat or open flame.

DISPOSAL:

- 1. PROHIBITIONS**
Do not contaminate water, food, or feed by disposal. Open dumping is prohibited.
- 2. PESTICIDE DISPOSAL**
Pesticide spray mixture or residue that cannot be used according to label instructions must be disposed of according to federal, state, or local procedure or under the Resource Conservation & Recovery Act.
- 3. CONTAINER DISPOSAL**
Triple rinse or equivalent and offer for recycling or reuse. If not recycled, dispose of in a sanitary landfill or by other approved state and/or local procedure.
- 4. GENERAL**
Consult federal, state or local disposal authorities for approved alternative procedures.



**CORROSIVE CAUSES EYE DAMAGE
AND SKIN BURNS**

CULTURAL PRODUCTS, INC.
NTI, MI 48197 • PHONE: (313) 484-1773

EPA Est. No. 44718-MI-1
EPA Reg. No. 44718-3

WARRANTY: As with the other in the Pates, Tex III, there is NO WARRANTY, representation or statement of any kind, expressed or implied, concerning NO WARRANTY OF MERCHANTABILITY, fitness for a particular purpose, or any other matter. Buyer's use and use of this product USE PATES, TEX III, IN ACCORDANCE WITH THE DIRECTIONS AND WARNINGS GIVEN ON THE LABEL. BUYER WAIVES ALL CLAIMS AGAINST THE MANUFACTURER, DISTRIBUTOR, or other party, whether or not such party is a manufacturer, distributor, or other party, in the event of any breach of warranty, strict liability, or other cause of action, connected with the use of Pates, Tex III, other than that which is set forth in the directions and warnings given on the label.