PRECAUTIONARY STATEMENTS **HAZARDS TO HUMANS** AND DOMESTIC ANIMALS

DANGER: Corrosive, may cause severe skin irritation or chemical burns to broken skin. Causes eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling this product. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until coors have dissi-

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of takes, streams or ponds. Treated effluent may not be discharged into lakes, streams, ponds or public waters without a valid discharge permit. For guidance, contact the regional office of the Environmental Protection Agency.

PHYSICAL AND CHEMICAL HAZARDS: STRONG OXIDIZING AGENT: Mix only with water according to label directions. Mixing this product with gross filth such as feces, urine, etc., or with ammonia, acids, detergents or other chemicals may release hazardous gases irritating to eyes, lungs and mucous membranes.

manufactured by

PURE WATER 13 Sillcon Way Bradford, PA 16701

EPA REG. NO. 44899-1 EPA EST. NO. **NET CONTENTS.**

ACCEPTED

44899-1

DEC 0 1 1980

Under the Federal I 😁 Pungicido, and as amended, los ODIUM recistered under HYPOCHLORITE SOLUTION

FOR SWIMMING POOL CHLORINATION

ACTIVE INGREDIENT SODIUM HYPOCHLORITE INERT INGREDIENTS

87 5%

KEEP OUT OF REACH OF CHILDREN DANGER

FIRST AID: If on skin, wash with plenty of soap and water. If in eyes, flush with water for at least 15 minutes. Get medical attention. If swallowed, drink large quantities of milk, or gelatin solution or, if these are not available, drink large quantilies of water Do NOT give vinegar or other acids. Do NOT induce romiting Get prompt medical attention

See additional precautions on side panel

DIRECTIONS FOR USE GENERAL CLASSIFICATION

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

DIRECTIONS FOR SWIMMING POOL CHLORINATION For a new pool or for spring start-up, superchlorinate with one (1) pint (16 fiur ozs.) a sodium hypochlorite solution for each 3000 gations of water. This design is equivalent to 5 ppm available chlorine by weight.

5 ppm available chlorine by weight
r Pool Maintenance
Adjust pool water pH to 7.6 - 7.2 range and maintain
Add manually or by a feeder device this sodium hypochlorite solution at a rate tmaintain an available chlorine residual of 0.6 to 1.0 ppm. One-fourth punt (4.6)
outs) of this sodium hypochlorite solution for each 4000 gallions of water will provide 1.0 ppm available chlorine by weight. Frequency of additions to maintain 0.6 to 1.0 ppm available chlorine will depend on temperature and number of swimmers.
Use text to make certain the pH and chlorine residual are in the proper range.

DIRECTIONS FOR DISINFECTION OF POTABLE WATER
FOR HOME WELL WATER SYSTEMS.

This sodium hypochlorite solution in the ratio of one part sodium hypochlorite.

All a sodium hypochlorite solution and water thor

FOR HOME WELL WATER SYSTEMS

Dilute this sodium hypochlorite solution in the ratio of one part sodium hypochlorite solution to 11 parts softened water. Mix sodium hypochlorite solution and water thor oughly and begin feeding of solution with a hypochlorinator (metering pump). Maintain a free available chiforine residual of all least 0.2 ppm and no more than 0.6 ppm throughout the distribution system, as determined by a DPD chlorine test kit. Bacter iological sampling must be conducted at a frequency no less than that prescribed by the National Interim Primary Drinking Water Regulations. Check water frequently with a DPD chlorine test kit.

"Comtact your local Health Department for further details.

DIRECTIONS FOR SANITIZING FOOD PROCESSING
OR DAIRY EQUIPMENT

Clean equipment in the normal manner. Just before using innse all surfaces thorough by with this sodium hypochlorite solution containing 200 prima variable chlorine. Maintain contact with disinfectant for a minimum of two minutes. Do not rinse with water after treatment with sodium hypochlorite solution. Do not soak overright. Addition of one fit do at 20 black overright.

after treatment with sodium hypochlorite solution. Do not soak overright. Addition of one fluid or (2 libits) of this sodium hypochlorite solution per 5 gailons of waterwill provide approximately 200 ppm available chlorine by weight. PCTE, This product degrades with age. Use a chlorine lest kit and increase dosage as recessly, to other the gequired fevel of available chlorine.

STORAGE AND DISPOSAL: Store in a coot, dry area away from Greg sufficient to case of spill, flood area with large quantities of water. Rinse compty container more upfit with water and either return to manufacturer or discard by placing in trash folloction or burying in an approved landfull Product or misse that cannot be used, should be diluted with water and disposed of in a sanitary sewer. Do not contaminate food, or feed by storage, disposal or cleaning of equipment.

34