

**PRECAUTIONARY STATEMENTS
HAZARDOUS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Avoid breathing dust, fumes, or vapors from this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when dispensing of equipment washers. Do not apply when weather conditions favor drift from treated areas.

PHYSICAL HAZARDS

Sulfur dust suspended in air ignited easily. Keep away from heat, sparks, or flames. Do not smoke while applying this product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Sulfur may cause severe frost and foliage injury to certain crops. Sulfur sensitive plants include onion and onion potato, apricot, cherry, certain varieties of cucumber, alfalfa, spinach, long beans, and certain ornamentals. Apples, pears and strawberries are susceptible to foliage burn with sulfur under certain climatic conditions. Sulfur may burn foliage when temperatures are high. Do not apply if temperatures during or within 3 days after application are expected to exceed 90°F in the area to be treated. Do not use within 2 weeks of an oil spray treatment on citrus within 3-6 weeks. When growing crops for processing, consult the processor before applying sulfur.

Washers needed connect into bucket or other, stir to mix thoroughly. Add to tank when part full and before other materials are added. Add wetting agents last to avoid foaming.

RE-ENTRY STATEMENT

Do not enter treated areas without protective clothing for 24 hours after application. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The crop being treated must be protected by appropriate means. Certain states may require more restrictive re-entry intervals for various crops treated with this product. Consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area 24 hours after application. Written or oral warnings are given, warnings shall be given in a conspicuous, easily understood manner. Written or oral warnings shall include the following information: **CAUTION** Area treated with sulfur. Do not enter treated areas without appropriate protective clothing for 24 hours following application. **CAUTION** Sulfur dust suspended in air ignited easily. Wash thoroughly with soap and water after handling.

STORAGE AND DISPOSAL

RESTRICTIONS:
Do not contaminate water, food or feed by storage or disposal.
SPILLAGE:
Sulfur is a solid, dry area.
CONTAINER DISPOSAL:
Completely empty bag into application equipment, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.
EMPTYING DISPOSAL:
Do not reuse containers. Do not use this product any longer. Do not use this product any longer. Do not use this product any longer.

CUPROQUIM CORPORATION

**SUPER-SUL
MICRONIZED WETTABLE SULFUR**

W D G
WATER EMULSIFIABLE GRANULES

ACTIVE INGREDIENTS: Sulfur.....80%
INERT INGREDIENTS:.....20%
TOTAL:.....100%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

HARMFUL IF SWALLOWED. DO NOT SWALLOW OR INHALE

See additional precautions on side panel

PRACTICAL TREATMENT (First Aid):


IF LARGE QUANTITIES ARE SWALLOWED give water and induce vomiting. Get medical attention.

IF INHALED and overcome by vapor or dust remove immediately from exposure and call a physician. Administer artificial respiration if breathing is irregular or has stopped.

FOR SKIN CONTACT flush with water; use soap if available.

IF IN EYES immediately flush with plenty of water for at least 15 minutes. Call a physician.

EPA No. **ACCEPTED** FILE NO. 45002-10 CR-81



This product manufactured for
CUPROQUIM CORPORATION
Houston, Texas 77074

NET WT. 30 LBS.

DIRECTIONS FOR DILUTION

DILUTE APPLICATION: Ground Application Apply specified rate in 20-40 gallons of water per acre. Ground Application apply specified rate in 100-200 gallons of water per acre.

CONCENTRATE APPLICATION: Ground Application Apply specified rate in not less than 20 gallons of water per acre. **SPRAYER APPLICATION:** Apply specified rate in 20-100 gallons of water per acre. Special concentrate equipment is necessary for these uses.

AIR APPLICATION: Apply specified rate in 3-20 gallons of water per acre. Ground rates should not be applied in less than 10 gallons of water per acre.

Do not apply this product through any type of irrigation system.
Use high volume water when ground equipment is desired.

CROPS:

Almonds: Brown Rot Blossom and Twig Blight Leaf Spot. Apply 10-15 lbs. per acre. Apply at bloom or early petal fall. Repeat as necessary usually 10-14 days or after a period of wet weather.

Apples: Pests: Powdery Mildew, Leaf Miner, Spider Mite, Red Spider Mite. Apply 10-15 lbs. per acre. Do not apply to 20-30 lbs. per acre. Apply ground and petal fall. Apply with cover spray throughout the season.
Berries: Powdery Mildew, Red Spider Mite. Apply 1-5 lbs. per acre. Start in early leaf fall stage. Repeat application every 10-14 days, with or without pesticides, immediately after a rain.
Citrus: Citrus Blight, Citrus Leafminer, Citrus Leafminer, Citrus Leafminer. Apply 1-5 lbs. per acre. Start in early leaf fall stage. Repeat applications every 10-14 days, with or without pesticides, immediately after a rain.
Citrus: Citrus Blight, Citrus Leafminer, Citrus Leafminer. Apply 1-5 lbs. per acre. Start in early leaf fall stage. Repeat applications every 10-14 days, with or without pesticides, immediately after a rain.

Cotton: Good Alfalfa. Red Spider Mite. Apply 1-5 lbs. per acre.
Corn: Powdery Mildew, Red Spider Mite, Leaf Miner, Spider Mite. Apply 1-5 lbs. per acre. Apply in late bloom or with late spray. Early should be petal and through bloom on 14 day intervals or with pesticides.

Cucumbers: Brown Spot, Powdery Mildew, Leaf Miner, Spider Mite. Apply 10-15 lbs. per acre. Apply at bloom or early petal fall. Repeat as necessary, usually 10-14 days, or after a period of wet weather.

Peaches and Nectarines: Brown Spot, Blight and Leaf Blight, Powdery Mildew, Leaf Miner, Spider Mite. Apply 10-15 lbs. per acre. Apply at bloom or early petal fall. Repeat as necessary, usually 10-14 days, or after a period of wet weather.

Pears: Powdery Mildew, Red Spider Mite, Leaf Miner, Spider Mite. Apply 10-15 lbs. per acre. Start in early leaf stage. Repeat applications as necessary usually 10-14 days.
Pineapples: Brown Spot, Blight and Leaf Blight, Powdery Mildew, Leaf Miner, Spider Mite. Apply 10-15 lbs. per acre. Apply at bloom or early petal fall. Repeat as necessary, usually 10-14 days, or after a period of wet weather.

Strawberries: Powdery Mildew, Red Spider Mite. Apply 1-5 lbs. per acre. Start in early leaf stage. Repeat applications with or without pesticides, for Powdery Mildew begin at first signs of disease and repeat as necessary.

Tomatoes: Brown Spot, Powdery Mildew, Leaf Miner, Spider Mite. Apply 10-15 lbs. per acre. Apply at bloom or early petal fall. Repeat as necessary, usually 10-14 days, or after a period of wet weather.

Washers: Washers needed connect into bucket or other, stir to mix thoroughly. Add to tank when part full and before other materials are added. Add wetting agents last to avoid foaming.

Washers: Washers needed connect into bucket or other, stir to mix thoroughly. Add to tank when part full and before other materials are added. Add wetting agents last to avoid foaming.

Washers: Washers needed connect into bucket or other, stir to mix thoroughly. Add to tank when part full and before other materials are added. Add wetting agents last to avoid foaming.

Washers: Washers needed connect into bucket or other, stir to mix thoroughly. Add to tank when part full and before other materials are added. Add wetting agents last to avoid foaming.

Washers: Washers needed connect into bucket or other, stir to mix thoroughly. Add to tank when part full and before other materials are added. Add wetting agents last to avoid foaming.

Washers: Washers needed connect into bucket or other, stir to mix thoroughly. Add to tank when part full and before other materials are added. Add wetting agents last to avoid foaming.

BEST AVAILABLE COPY