

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of rinsate or equipment washwater. Do not contaminate water when disposing of equipment washwater.

PHYSICAL HAZARDS

Sulfur dust suspended in air ignites easily. Keep away from heat, sparks, or flames. Do not smoke while applying this product.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system. Sulfur may cause severe fruit and foliage injury to certain crops. Sulfur sensitive plants include Anjou and Comice pears, apricots, cranberries, certain varieties of cucurbits, filberts, spinach, Tung walnuts, and certain ornamentals. Apples, pears and strawberries are susceptible to foliage burn with sulfur under certain climatic conditions. Sulfur may burn foliage when temperature is high. Do not apply if temperatures during or within 3 days after application are expected to exceed 90°F in the area to be treated. Do not use within 2 weeks of an oil spray treatment (3 weeks for citrus) unless stated otherwise on this label. When growing crops for processing, consult the processor before applying sulfur.

Measure needed amount into bucket of water, stir to make thin slurry. Add to tank when part full and before other materials are added. Add wetting agents last to avoid foaming.

ACCEPTED

MAY 25 1993

Under the Federal Insecticide, Fungicide, and
Rodenticide Act, as amended, for the
pesticide registered under
EPA Reg. No. 45002-10

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter into treated area until at least 24 hours after the completion of application to the area unless wearing the protective clothing listed below. During early re-entry into treated areas (less than 24 hours after the completion of application) to perform hand labor tasks, wear long pants (or coveralls), long sleeved shirt, shoes, socks, chemical/water resistant gloves, goggles or face shield.

Pesticide Handlers: During mixing, loading and application of this product, wear long pants (or coveralls), long sleeved shirt, shoes, socks, chemical/water resistant gloves, and goggles or face shield.

Certain states may require more restrictive re-entry intervals for various crops treated with this product. Consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written and oral warnings must include the following information. CAUTION: Area treated with Sulfur on (date of application). Do not enter treated areas without appropriate protective clothing for 24 hours following application.

Required Protective Clothing: Long pants (or coveralls), long sleeved shirt, shoes, socks, chemical/water resistant gloves, goggles or face shield.

+-----+
STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal of this product.

STORAGE: Store in cool, dry area.

CONTAINER DISPOSAL: Completely empty bag into application equipment, then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

PRODUCT DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

+-----+

CUPROQUIM CORPORATION
SUPER-SUL
MICRONIZED WETTABLE SULFUR
WDG
WATER DISPERSIBLE GRANULE

ACTIVE INGREDIENTS	BY WT.
Sulfur.....	80%
INERT INGREDIENTS.....	20%
TOTAL.....	100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

HARMFUL IF SWALLOWED. DO NOT SWALLOW OR INHALE.

STATEMENT OF PRACTICAL TREATMENT (First Aid):

IF SWALLOWED: Give 1-2 glasses of water and induce vomiting. Get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED, overcome by vapor or dust: Remove victim to fresh air immediately and call a physician. Administer artificial respiration if breathing is irregular or has stopped.

IF ON SKIN: Wash with plenty of soap and water. Remove and wash contaminated clothing before reuse.

IF IN EYES: Immediately flush with plenty of water for at least 15 minutes. Call a physician.

SEE ADDITIONAL PRECAUTIONS ON SIDE PANEL

CUPROQUIM CORPORATION

9601 Katy Freeway, Suite 350
Houston, Texas 77024-1333 USA
TEL: (713) 464-1103 FAX: (713) 464-1421
TELEX II: 910-240-6712 PDC

MANUFACTURED FOR CUPROQUIM CORPORATION

EPA Reg. No.: 45002-10
EPA Est. No.: 39827-CH-01
Form. No.: 2-10-0891

NET WEIGHT 30 LBS.

DIRECTIONS FOR DILUTION

DILUTE APPLICATION: Ground Application Apply: specified rate in 20-60 gallons of water per acre. Orchard application: Apply specified rate in 100-800 gallons of water per acre.

CONCENTRATE APPLICATION: Ground Application: Apply specified rate in not less than 5 gallons of water per acre. Orchard Application: Apply specified rate in 20-100 gallons of water per acre. Special concentrate equipment is necessary for these uses.

AERIAL APPLICATION: Apply specified rate in 5-20 gallons of water per acre. Orchard rates should not be applied in less than 10 gallons of water per acre.

USE HIGH DOSAGE RATES WHEN DISEASE PRESSURE IS SEVERE.

ALMONDS: Brown Rot, Blossom and Twig Blight, Leaf Spot, Red Spider Mites. - Apply 10-30 lbs. per acre. Apply at bloom or early petal fall. Repeat as necessary, usually 10-14 days or after a period of wet weather.

APPLES, PEARS: Powdery Mildew, Scab, Blister Mites, Red Spider Mites. - Apply 10-30 lbs. per acre. Do not apply to D'Anjou pears. Apply pre-bloom and petal fall. Apply with cover sprays throughout the season. Apply with spray oil during the dormant and delayed dormant periods of bud development using 10-20 lbs./A. Do not apply to sensitive varieties of pears except at these stages.

ASPARAGUS: Rust - Apply 10-30 lbs. per acre. Use after cutting stops. Irrigate and cultivate before applying. Repeat at 7-10 day intervals through the fern growth season.

AVOCADOS: Brown Mite - Apply 10-30 lbs. per acre when mites are first noticed.

BEANS, PEAS, PEPPERS, POTATOES: Powdery Mildew, Red Spider Mites. - Apply 3 to 5 lbs. per acre. Start in early leaf stage. Repeat application every 10-14 days, with or without pesticides, immediately after a rain.

BROCCOLI, BRUSSELS SPROUT, CABBAGE, CAULIFLOWER, COLLARDS, KALE: Powdery Mildew, Septoria Leaf Spot, Red Spider Mites - Apply 3-5 lbs. per acre. Start in early leaf stage. Repeat applications every 10-14 days, with or without pesticides, immediately after a rain.

BLACKBERRY (Rubis Spp): Powdery Mildew - Apply 6-15 lbs. per acre. Begin when new shoots are 6 inches long and before blossoms open. Repeat at 10 day intervals or as necessary.

CARROTS, GARLIC, ONIONS: Petrobia mite and powdery mildew. - Apply 3-10 lbs. per acre when the disease first appears and repeat as needed.

CHERRIES: Brown Rot, Blossom and Twig Blight, Powdery Mildew, Leaf Spot, Silver Mites, Flat Mites, Red Spider Mites - Apply 10-30 lbs. per acre. Apply at bloom or early petal fall. Repeat as necessary, usually 10-14 days or after a period of wet weather.

CITRUS: Flat Mites, Silver Mites, Red Spider Mites - Apply 8-20 lbs. per acre. Apply in any spray - NOT CONTAINING OIL - or follow oil by 21 days.

COTTON, SEED ALFALFA: Red Spider Mites - Apply 3-5 lbs. per acre.

CUCURBITS: Powdery Mildew - Apply 4-6 lbs. per acre when the disease first appears and repeat as necessary. Application intervals throughout the growing season 10-14 days.

GRAPES: Powdery Mildew, Bug Mites, Blister Mites, Red Spider Mites - Apply 2-10 lbs. per acre. Apply in late dormant or with cutworm spray, early shoot period and through season on 14-day schedule or with pesticides.

HOPS: To aid in the suppression of Red Spider Mites, use 2-6 lbs. per acre by air or ground. Do not apply in less than 10 gallons of water per acre. Mix 2-6 lbs. per 100 gallons of water.

MELONS: Powdery Mildew - Apply 20-25 lbs. per acre. Treat when the disease first appears and repeat as necessary. Sulfur can injure plants, especially when the temperature reaches 95°F. Do not use on sulfur sensitive varieties.

DELETED MINT

PEACHES and NECTARINES: Brown Rot, Blossom and Twig Blight, Powdery Mildew, Silver Mites, Red Spider Mites - Apply 10-30 lbs. per acre. Apply at bloom or early petal fall. Repeat as necessary, usually 10-14 days, or after a period of wet weather.

PEANUTS: Powdery Mildew, Red Spider Mites, Leaf Spot - Apply 3-5 lbs. per acre. Start in early leaf stage. Repeat applications as necessary, usually 10-20 days.

PEPPERMINT, SPEARMINT: Powdery Mildew - Apply 4-6 lbs. per acre when mint is 5-6" tall or when disease appears. Repeat twice, at 30 day intervals. Do not apply within 30 days of harvest.

PLUMS and PRUNES: Brown Rot, Blossom and Twig Blight, Powdery Mildew, Silver Mites, Red Spider Mites - Apply 10-30 lbs. per acre. Apply at bloom or early petal fall. Repeat as necessary, usually 10-14 days, or after a period of wet weather.

RASPBERRY (Rubis Spp.): Powdery Mildew, (Sphaerotheca humuli) - Apply 6-15 lbs. per acre. Spray each week starting at first bloom color. Continue until all fruit is set, then repeat as weather requires.

STRAWBERRIES: Powdery Mildew, Red Spider Mites - Apply 5-10 lbs. per acre. Start in early leaf stages repeat application with or without pesticides. For Powdery Mildew, begin at first signs of disease and repeat as necessary.

SUGAR BEETS: Red Spider Mites (Spp.), Powdery Mildew - Apply 10-20 lbs. per acre. Apply at first indication of problem with or without other pesticides. Repeat as necessary.

TOMATOES: Russet Mites - Apply 10-20 lbs. per acre. Thorough coverage is required.

WALNUTS: European Red Mites, Pacific Mites, Almond Mites, Two-spotted Mites, Red Spider Mites - Apply 10-20 lbs. per acre. Apply with mite sprays during season to increase the effectiveness of miticides. CAUTION: Certain varieties of walnuts are sensitive to sulfur sprays under certain conditions. Do not apply unless the varieties are known to be tolerant to sulfur in your area.

SUPER-SUL: Applications will provide sulfur for the plant's nutrient requirements and should be considered in the total plant nutrient applications. SUPER-SUL can be used as a micronutrient supplement by either foliar or soil application. SUPER-SUL is recommended on the following crops: alfalfa, almonds, avocados, apples, barley, all beans, broccoli, cabbage, cauliflower, carrots, celery, citrus, corn, grapes, lawns, lettuce, melons, milo, nectarines, rice, pears, peaches, pecans, peppers, plums, prunes, potatoes, peanuts, sorghum, soybeans, sugar beets, sweet corn, strawberries, tomatoes, turnips, walnuts, watermelons, wheat and most other crops. Consult your State Agricultural Experimental Station or Extension Specialist for advice in selecting treatments to best fit local conditions.

-----+-----
 | NOTE: Seller warrants that the product conforms to its |
 | chemical description and is reasonably fit for the purpose |
 | stated on this label when used in accordance with direc- |
 | tions under normal condition of use: but neither this |
 | warranty or any other warranty of merchantability of fit- |
 | ness for a particular purpose, expressed or implied, |
 | extends to the use of this product contrary to label |
 | instructions not reasonably foreseeable to seller: the |
 | the buyer assumes the risk of any such use. |
 |-----+-----

.....

