

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Avoid breathing spray mist. Avoid contact with eyes, skin and clothing. Harmful if swallowed, inhaled, or absorbed through skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL HAZARDS

Sulfur dust suspended in air ignites easily. Keep away from heat, sparks, or flame. Do not smoke while applying this product.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

NOTE TO USER: Some plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proven that sulfur is safe in that locality.

During periods of high temperature sulfur may burn foliage and fruit. Do not make sulfur applications at excessively high temperatures. Do not use with oil or within 4 weeks of an oil application. When growing crops for processing, consult the processor before applying sulfur.

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing for 24 hours after application. Certain states may require more restrictive re-entry intervals for various crops treated with this product. Consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written and oral warnings must include the following information: "CAUTION: Area treated with sulfur on (date of application). Do not enter treated areas without appropriate protective clothing for 24 hours following application." Required Protective Clothing: long-sleeved shirt, long-legged pants and gloves.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal of this product.

STORAGE: Store in cool, dry area.

CONTAINER DISPOSAL: Completely empty container into application equipment, then dispose of empty container in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

PRODUCT DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CUPROQUIM CORPORATION

**SULFUR 6
FLOWABLE**

| | |
|-------------------------|--------|
| ACTIVE INGREDIENTS: | BY WT. |
| Sulfur..... | 52% |
| INERT INGREDIENTS:..... | 48% |
| TOTAL..... | 100% |

Contains 6 lbs. Sulfur per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

HARMFUL IF SWALLOWED. DO NOT SWALLOW OR INHALE

STATEMENT OF PRACTICAL TREATMENT (First Aid):

IF SWALLOWED: Give 1-2 glasses of water and induce vomiting. Get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED, overcome by vapor or dust: Remove victim to fresh air immediately.

IF ON SKIN: Wash with plenty of soap and water. Remove and wash contaminated clothing before reuse.

IF IN EYES: Immediately flush with plenty of water for at least 15 minutes.

SEE ADDITIONAL PRECAUTIONS ON SIDE PANEL

ACCEPTED



SPECIMEN LABEL

Under the provisions of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this pesticide is registered under...

CUPROQUIM CORPORATION

9601 Katy Freeway, Suite 350
Houston, Texas 77024-1333 USA
TEL: (713) 464-1103 FAX: (713) 464-1421
TELEX II: 910-240-6712 PDC

MANUFACTURED FOR CUPROQUIM CORPORATION

EPA Reg. No. 45002-18
EPA Est. No. 45002-MX-02
FORM #: 218-1191

DIRECTIONS FOR DILUTION

Stir until smooth before using. Wash recommended amount through screen into partially filled spray tank or pre-mix in a small amount of water before pouring into tank. Do not use a screen finer than 50 mesh in entire system. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Failure to maintain agitation will cause sulfur to settle and necessitate manual removal of the deposit. This product should be applied by ground equipment or aircraft in sufficient water to provide thorough coverage.

Do not apply this product through any type of irrigation system.

Tree Crops: Unless otherwise stated for specific crop, dosage rates are given as pints of SULFUR 6 FLOWABLE per 100 gal. water for use in a thorough coverage spray. The total amount of SULFUR 6 FLOWABLE per acre should be 2 - 4 gallons depending on size of trees. Because of spray equipment variations and type used, a range of low and high rates are listed. For high volume sprayers (output 800 - 1000 gals. spray per acre) use the low rate. For low volume sprayers (output 20 - 300 gals. spray per acre) use high rate.

Apples: Apple cedar rust, black rot, frog-eye leaf spot, powdery mildew, quince rust, scab, sooty blotch - Apply 1 1/2 - 3 1/2 pts. from prepink through calyx periods and 3/4 - 2 pts. in cover sprays. **NOTE:** McIntosh, Golden Delicious, Jonathan and certain other varieties may be injured by sulfur application under certain conditions.

Cherries: Brown rot, leaf spot, powdery mildew, rust, scab and stem blight - Apply 1 1/2 - 3 1/2 pts. through bloom stage and 3/4 - 1 3/4 pts. from petal fall through cover sprays for brown rot and leaf spot.

Citrus: Rust mite, clover mite - Apply 3/4 - 4 1/4 pts. in November through May. Thrips (Early spring treatment for the reduction of thrip population) - Apply 1 - 2 gallons per acre in sufficient water for thorough coverage. Make application after spring flush has made 3 to 4 inch growth. Repeat if necessary. Do not apply within 21 days of an oil application.

Grapes: Powdery mildew, rust (physcopella) - Apply 1 - 2 pts per 100 gal. water using up to 8 pints SUPER 6 FLOWABLE per acre. Begin when new shoots are 6 to 10 inches long. Repeat before blossoms open and continue at 10 to 14 day intervals as necessary. **NOTE:** Concord and other Labrusca type grapes may be injured.

Nectarines, Peaches: Brown rot, leaf spot, powdery mildew, peach canker, rust, scab, shot hole - Apply 1 1/2 - 3 1/2 pts. through bloom and 3/4 - 1 1/4 pts. in cover sprays. Apply from early petal fall through cover periods as necessary.

Pears: Brown rot, leaf spot, powdery mildew, rust, scab - Apply 1 1/2 - 3 1/2 pts. through calyx and 3/4 - 2 pts. in cover sprays. Apply from prebloom through calyx periods as necessary.

Plums, Prunes: Brown rot, leaf spot, powdery mildew, rust, scab, shot hole - Apply 1 1/4 - 7 1/4 pts. through bloom and 3/4 - 2 pts. through cover sprays for brown rot and leaf spot.

Strawberries: Powdery mildew - Apply 2 pts. per acre beginning at first sign of disease and repeat as necessary. **NOTE:** Sulfur may injure strawberries under certain climatic conditions.

NOTE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal condition of use; but neither this warranty or any other warranty of merchantability of fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions not reasonably foreseeable to seller; the buyer assumes the risk of any such use.