



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

45002-38

Date of Issuance:

4/21/22

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

DICAMBA DMA SALT 5

Name and Address of Registrant (include ZIP Code):

ALBAUGH LLC
1525 NE 36th Street
Ankeny, IA 50021

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Emily Schmid, Product Manager 25
Herbicide Branch, Registration Division (7505P)

Date:

4/21/22

2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 45002-38.”
3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 6/25/2021

If you have any questions, please contact Lydia Crawford by phone at 202-566-2575, or via email at Crawford.Lydia@epa.gov.

Enclosure

DICAMBA DMA SALT 5


For Weed Control in Corn, Sorghum, Small Grains (wheat, barley, and oats), Pasture, Hay, Rangeland, General Farmstead (Non-Cropland), Fallow croplands, Cotton, Sugarcane, Asparagus, Turf, and Grass Seed Crops

ACTIVE INGREDIENT:

Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)* 60.63%
 OTHER INGREDIENTS: 39.37%
TOTAL: 100.0%

*This product contains 50.36% 3,6-dichloro-o-anisic acid (dicamba)
 (5 lbs Dicamba acid equivalent (a.e.) per gallon)

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**



| FIRST AID | |
|--|---|
| IF IN EYES: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| <p>HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.</p> | |

See inside booklet for complete First Aid, Precautionary Statements, Directions For Use, State-Specific Crop and/or Use Site Restrictions and Conditions of Sale and Warranty.

EPA Reg. No. 45002-____

EPA Est. No. _____

NET CONTENTS: _____

MANUFACTURED BY:
 1525 36th Street
 Albaugh, LLC
 Ankeny, Iowa 50021

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear appropriate protective eyewear such as goggles, face shield, or safety glasses. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT

All mixers, loaders, and applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Chemical-resistant gloves (except for applicators using ground boom equipment, pilots and flaggers) including Barrier Laminate, Butyl Rubber ≥14 mils, Nitrile Rubber ≥14 mils, Neoprene Rubber ≥14 mils, Natural Rubber ≥14 mils, Polyethylene, Polyvinyl Chloride (PVC) ≥14 mils or Viton ≥14 mils.
- Shoes plus socks, and protective eyewear.

See engineering controls for additional requirements and exceptions.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use cockpits in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d) (4-6)).

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

Groundwater Advisory: Dicamba is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow coming in contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

1. Coveralls worn over short-sleeved shirt and short pants.
2. Chemical resistant footwear plus socks
3. Chemical resistant gloves made of any waterproof material.
4. Chemical resistant headgear for overhead exposure.
5. Protective eyewear

Notify workers of application by warning them orally and by posting warning signs at entrances to treated area.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

WEED RESISTANCE MANAGEMENT

| | | | |
|---------|-------|---|-----------|
| DICAMBA | GROUP | 4 | HERBICIDE |
|---------|-------|---|-----------|

For resistance management, this product is a group 4 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.

- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method including hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, consult an Albaugh LLC representative at 1-800-247-8013.

PRODUCT INFORMATION

Mode of Action: This product is readily absorbed by plants through shoot and root uptake, translocates throughout the plant's system, and accumulates in areas of active growth. This product interferes with the plant's growth hormones (auxins) resulting in death of many broadleaf weeds.

The following directions apply to all uses of DICAMBA DMA SALT 5. Additional precautions and restrictions will be found in each specific use section. Do not treat irrigation ditches or water used for crop irrigation or domestic uses. Do not apply this product through any type of irrigation system.

Rates of use of this product must not exceed 25.6 fl. oz. per acre, per application; and 51.2 fl. oz. per acre, per year. Rates of use of dicamba acid equivalent must not exceed a maximum single application rate of 25.6 fl. oz. (1.0 lb) per acre; and an annual application rate of 51.2 fl. oz. (2.0 lbs) per acre. These restrictions apply to this product and possible mixtures with other products containing dicamba.

GROUND AND SURFACE WATERS PROTECTION

- 1) Point source contamination – To prevent point source contamination, do not mix or load this pesticide product within 50 feet of wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. Do not apply pesticide product within 50 feet of wells. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas as described below.

Mixing, loading, rinsing, or washing operations performed within 50 feet of a well are allowed only when conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be on or move across the pad. The pad must be self-contained to prevent surface water flow over or from the pad. The pad capacity must be maintained at 110% that of the largest pesticide container or application equipment used on the pad and have sufficient capacity to contain all product spills, equipment or container leaks, equipment wash waters, and rainwater that may fall on the pad. The containment capacity does not apply to vehicles delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Care must be taken when using this product to prevent:

- a) back-siphoning into wells,
- b) spills or
- c) improper disposal of excess pesticide, spray mixtures or rinsate. Check valves or anti-siphoning devices must be used on all mixing equipment.

- 2) Movement by surface runoff or through soil – Do not apply under conditions which favor runoff. Do not apply to impervious substrates including paved or highly compacted surfaces in areas with high potential for ground water contamination. Ground water contamination may occur in areas where soils are permeable or coarse and ground water is near the surface. Do not apply to soils classified as sand with less than 3% organic matter and where ground water depth is shallow. To minimize the possibility of ground water contamination, carefully follow application rate instructions as affected by soil type in the product information section of this label.
- 3) Movement by water erosion of treated soil – Do not apply or incorporate this product through any type of irrigation equipment nor by flood or furrow irrigation. Ensure treated areas have received at least one-half inch rainfall (or irrigation) before using tailwater for subsequent irrigation of other fields.

SENSITIVE CROP PRECAUTIONS

DICAMBA DMA SALT 5 may cause injury to desirable trees and plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes, and other broadleaf plants when contacting their roots, stems or foliage. These plants are most sensitive to DICAMBA DMA SALT 5 during their development or growing stage. Follow the precautions and restrictions listed below when using DICAMBA DMA SALT 5.

PRECAUTIONS WHEN APPLYING DICAMBA DMA SALT 5

- Use coarse sprays to avoid potential herbicide drift. Select nozzles which are designed to produce minimal amounts of fine spray particles. Examples of nozzles designed to produce coarse sprays via ground applications are Delavan Raindrops, Spraying Systems XR flat fans or large capacity flood nozzles including D10, TK10, or greater capacity tips. Keep the spray pressure at or below 20 psi and the spray volume at or above 20 GPA, unless otherwise required by the manufacturer of drift-reducing nozzles. Consult with your spray nozzle supplier concerning the choice of drift-reducing nozzles.
- Agriculturally approved drift-reducing additives may be used.
- To avoid injury to desirable plants, thoroughly clean the equipment used to apply Dicamba DMA salt before using the equipment to apply any other chemical. (see PROCEDURE FOR CLEANING SPRAY EQUIPMENT).
- Crops growing under stress conditions including drought, poor fertility, or foliar damage due to hail, wind or insects, can exhibit various injury symptoms that may be more pronounced if herbicides are applied.
- Consult your local or state authorities for possible application restrictions and advice concerning these and other special local use situations. Tank mix instructions are for use only in states where the tank mix product and application site are registered.

RESTRICTIONS WHEN APPLYING DICAMBA DMA SALT 5

- Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of DICAMBA DMA SALT 5 with the roots of desirable plants including trees and shrubs.
- Do not apply when spray particles maybe carried by air currents to areas where sensitive crops and plants are growing or when temperature inversions exist.
- Do not spray near sensitive plants if wind is gusty or in excess of 5 mph and moving in the direction of adjacent sensitive plants.
- Do not apply DICAMBA DMA SALT 5 adjacent to sensitive crops when the temperature on the day of application is expected to exceed 85°F as drift is more likely to occur.

All crop uses of DICAMBA DMA SALT 5 are intended for a normal growing interval between planting and harvest. No crop rotation restrictions exist if normal harvest of treated crop has occurred. If this interval is shortened, including in cover crops that will be plowed under, do not follow up with the planting of a sensitive crop.

MIXING AND APPLICATION

The applicator must follow the most restrictive use cautions to avoid drift hazards, including those found in this labeling, as well as state and local regulations and ordinances.

Unless otherwise specified under the individual use headings of this label, the following directions apply to all crop and noncrop uses of DICAMBA DMA SALT 5. Refer to individual use sections for additional precautions, restrictions, application rates and timings.

DICAMBA DMA SALT 5 is a water-soluble formulation that can be applied using water or sprayable fluid fertilizer as the carrier. If a fluid fertilizer is to be used, a compatibility test (see COMPATIBILITY TEST below) should be made prior to tank mixing.

Use ground or aerial application equipment that will give good spray coverage of weed foliage. However, do not use aerial application equipment if spray particles can be carried by wind into areas where sensitive crops or plants are growing or when temperature inversions exist.

Apply 3 to 50 gallons of diluted spray per treated acre when using ground application equipment, or 2 to 10 gallons of diluted spray per treated acre (2 to 20 gallons of diluted spray per acre for preharvest uses) in a water-based carrier when using aerial application equipment. Use the higher level of the listed spray volumes when treating dense or tall vegetation. Use coarse sprays.

Select nozzles designed to produce minimal amounts of fine spray particles. Spray with nozzles as close to the weeds as is practical for good weed coverage.

To avoid uneven spray coverage, DICAMBA DMA SALT 5 should not be applied during periods of gusty wind or when wind is in excess of 15 mph.

Avoid disturbing (e.g., cultivating or mowing) treated areas for at least 7 days following application.

BAND TREATMENTS

DICAMBA DMA SALT 5 may be applied as a band treatment. Use the formulas below to determine the appropriate rate and volume per treated acre.

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast rate per treated acre} = \text{Band rate per treated acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast volume per treated acre} = \text{Band volume per treated acre}$$

COMPATIBILITY TEST

Before mixing in the spray tank, it is advisable to test compatibility by mixing all components in a small container in proportionate quantities (see following table).

Amount of Herbicide to Add to One Pint of Spray Carrier
(Assuming Volume is 25 Gallons per Acre)

| HERBICIDE FORMULATIONS | RATE PER ACRE | LEVEL TEASPOONS |
|------------------------|---------------|-----------------|
| Dry | 1 lb. | 1 ½ |
| Liquid | 1 pt. | ½ |

If herbicide(s) do not ball-up or form flakes, sludge, gels, oily films or layers, or other precipitates, then the tested spray mix is compatible. Usually, incompatibility in any of the above described forms will occur within 5 minutes after mixing.

If components are incompatible, the use of a compatibility agent is required. Rerun the above COMPATIBILITY TEST with a suitable compatibility agent (1/4 teaspoon is equivalent to 2 pints per 100 gallons of fluid fertilizer).

PROCEDURE FOR CLEANING SPRAY EQUIPMENT

The steps listed below are suggested for thorough cleaning of spray equipment following applications of DICAMBA DMA SALT 5 or tank mixes of DICAMBA DMA SALT 5 plus 2,4-D amine:

1. Hose down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of the rinse water.
2. Fill tank with water while adding 1 quart of household ammonia for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
3. Flush the solution out of the spray tank through the boom.
4. Remove the nozzles and screens and flush the system with two full tanks of water.
5. The steps listed below are suggested for thorough cleaning of spray equipment used to apply DICAMBA DMA SALT 5 as a tank mix with wetttable powders (WP), emulsifiable concentrates (EC), or other types of water-dispersible

formulations. DICAMBA DMA SALT 5 tank mixes with water-dispersible formulations require the use of a water/detergent rinse:

6. Complete step 1.
7. Fill tank with water while adding 2 lbs. of detergent for every 40 gallons of water. Operate the pump to circulate the detergent solution through the sprayer system for 5 to 10 minutes and discharge a small amount of the solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
8. Flush the detergent solution out of the spray tank through the boom.
9. Repeat step 1, and follow with steps 2,3, and 4.

WEED LIST

This is a list of weeds which may be treated with DICAMBA DMA SALT 5 in accordance with this label as required under the RATES AND TIMING sections of the individual use headings. Proper usage of this product will give control of growth suppression of many annual, biennial, and perennial broadleaf weeds, and many woody brush and vine species including:

ANNUALS

| | |
|------------------------------------|--|
| Amaranth, Spiny (Spiny Pigweed) | Mustard (Yellowtops) |
| Aster, Slender | Nightshade, Black |
| Bedstraw | Pennycress, Field (Fanweed, Frenchweed, Stinkweed) |
| Beggarweed, Florida | Pepperweed, Virginia (Peppergrass) |
| Broomweed, Common | Pigweed, Prostrate |
| Buckwheat, Wild | Pigweed, Redroot (Carelessweed) |
| Buffalobur | Pigweed, Rough |
| Burclover, California | Pigweed, Smooth |
| Burcucumber | Pigweed (triazine resistant) |
| Buttercup Roughseed | Pigweed, Tumble |
| Carpetweed | Poorjoe |
| Catchfly, Nightflowering | Puncturevine |
| Chamomile, Corn | Purslane, Common |
| Chickweed, Common | Pusley, Florida |
| Clovers (Annual) | Radish, Wild |
| Cockle, Corn | Ragweed, Common |
| Cockle, Cow | Ragweed, Giant (Buffaloweed) |
| Cocklebur, Common | Ragweed, Lance-Leaf |
| Croton, Tropic | Rubberweed, Bitter (Bitterweed) |
| Croton, Woolly | Sesbania, Hemp |
| Daisy, English | Shepherdspurse |
| Evening Primrose, Cutleaf | Sicklepod |
| Fleabane, Annual | Sida, Prickly (Teaweed) |
| Goosefoot, Nettleleaf | Smartweed, Green |
| Henbit | Smartweed, Pennsylvania |
| Jimsonweed | Sneezeweed, Bitter |
| Knotweed | Sowthistle, Annual |
| Kochia | Sowthistle, Spiny |
| Ladysthumb | Spikeweed, Common |
| Lambsquarters, Common | Spurge, Prostrate |
| Lambsquarters (triazine resistant) | Spurry, Corn |
| Lettuce, Prickly | Starbur, Bristly |
| Mallow, Common | Sumpweed, Rough |
| Mallow, Venice | Sunflower, Common (Wild) |
| Mare's Tail (Horseweed) | Sunflower, Volunteer |
| Mayweed | Thistle, Russian |
| Moringglory, Ivyleaf | Velvetleaf |
| Moringglory, Tall | Waterhemp |
| Mustard, Tansy | Waterprimrose, Winged |
| Mustard, Wild | Wormwood, Annual |

BIENNIALS

| | |
|----------------------------------|-------------------|
| Burdock, Common | Plantain, Bracted |
| Carrot, Wild (Queen Anne's Lace) | Ragwort, Tansy |

Cockle, White
Evening Primrose, Common
Geranium, Carolina
Gromwell
Knapweed, Diffuse
Knapweed, Spotted
Mallow, Dwarf

Starthistle, Yellow
Sweetclover
Teasel
Thistle, Bull
Thistle, Milk
Thistle, Musk
Thistle, Plumeless

PERENNIALS

*Alfalfa
Artichoke, Jerusalem
Aster, Spiny
Aster, Whiteheath
Beadstraw, Smooth
Bindweed, Field
Bindweed, Hedge
Blueweed, Texas
*Bursage (Bur Ragweed, Lakeweed, Povertyweed)
Bursage, Woollyleaf (Lakeweed)
Buttercup, Tall
Campion, Bladder
Chickweed, Field
Chickweed (Mouseear, Canada)
Chicory
*Clover, Hop
*Dandelion, Common
*Dock, Broadleaf (Bitterdock)
*Dock, Curly
Dogbane, Hemp
*Dogfennel (Cypressweed)
Fern, Bracken
Garlic, Wild
Goldenrod, Canada
Goldenrod, Missouri
Goldenweed, Common
Hawkweed
Henbane, Black
Horsenettle, Carolina
Ironweed
Knapweed, Black
Knapweed, Russian

Milkweed, Climbing
Milkweed, Common
Milkweed, Honeyvine
Milkweed, Western Whorled
Nettle, Stinging
Nightshade, Silverleaf (White Horsenettle)
Onion, Wild
*Plantain, Broadleaf
Plantain, Buckhorn

Pokeweed
Ragweed, Western
Redvine
Sericia Lespedeza
Smartweed, Swamp
Snakeweed, Broom
*Sorrel, Red (Sheep Sorrel)
Sowthistle
Sowthistle, Perennial
Spurge, Leafy
Sundrop, Halfshrub (Evening Primrose)
Thistle, Canada
Toadflex, Dalmation
Tropical Soda Apple
Trumpet creeper (Buckvine)
Vetch
Waterhemlock
Waterprimrose, Creeping
*Woodsorrel, Creeping Common Yellow
Wormwood, Common
Wormwood, Louisiana
*Yankee weed
Yarrow, Common

*Noted perennials may be controlled using DICAMBA DMA SALT 5 at rates lower than those for other listed perennial weeds. (See APPLICATION RATES AND TIMING section in this label.)

WOODY PLANTS

| | |
|---------------|------------------------|
| Alder | Locust, Black |
| Ash | Maple |
| Aspen | Mesquite |
| Basswood | Oak |
| Beech | Oak, Poison |
| Birch | Olive, Russian |
| *Blackberry | Persimmon, Eastern |
| *Blackgum | Pine |
| *Cedar | *Plum Sand (Wild Plum) |
| Cherry | Poplar |
| Chinquapin | Rabbitbrush |
| Cottonwood | *Redcedar, Eastern |
| *Creosotebush | *Rose, McCartney |

| | |
|-------------------------|-------------------|
| Cucumbertree | *Rose, Multiflora |
| *Dewberry | Sagebrush, Fringe |
| *Dogwood | Sassafras |
| Elm | Serviceberry |
| Grape | Spicebush |
| *Hawthorn, (Thornapple) | Spruce |
| Hemlock | Sumac |
| Hickory | *Sweetgum |
| Honeylocust | Sycamore |
| Honeysuckle | Tarbush |
| Hornbeam | Willow |
| Huckleberry | Witchhazel |
| Huisache | *Yaupon |
| Ivy, Poison | *Yucca |
| Kudzu | |

*Growth Suppression

CROP SPECIFIC USE INSTRUCTIONS

FIELD, SEED, POPCORN* AND SILAGE CORN

Observe all precautions, mixing and application instructions.

Corn Use Restrictions:

- **Maximum Single Application Rate:** DO NOT exceed a total of 12.8 fluid ounces (0.5 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Annual Application Rate:** DO NOT exceed a total of 19.2 fluid ounces (0.75 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Number of Applications per Year:** 2
- **Retreatment Interval:** 2 weeks.
- **Pre-Harvest Interval:** Corn may be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity.
- DICAMBA DMA SALT 5 is not registered for use on sweet corn.
- Up to 2 applications of DICAMBA DMA SALT 5 may be made per year.
- Do not exceed a total of 19.2 fl. oz. of DICAMBA DMA SALT 5 per treated acre per crop year.
- Sequential applications must be separated by 2 weeks or more. See appropriate section for rate information. For combination options or sequential treatments, refer to appropriate section.
- DICAMBA DMA SALT 5 contains 0.625 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.
- Do not use adjuvants containing penetrants including petroleum-based oils after crop emergence or crop injury may result.
- For Preemergence in Conventional or Reduced Tillage Corn Use Only: DO NOT apply to coarse textured soils (sand, sandy loam, and loamy sand) until after crop emergence (see EARLY POSTEMERGENCE uses below).

Corn Use Precautions:

- Do not apply DICAMBA DMA SALT 5 to seed corn or popcorn without first verifying with your local seed corn company (supplier) the DICAMBA DMA SALT 5 selectivity on your inbred line or variety of popcorn. This precaution will help avoid potential injury of sensitive varieties.
- Direct contact of DICAMBA DMA SALT 5 with corn seed must be avoided. If corn seeds are less than 1 ½ inches below the surface, delay application until corn has emerged.
- Applications of DICAMBA DMA SALT 5 to corn during periods of rapid growth may result in temporary leaning. Corn will usually become erect within 3 to 7 days.
- Cultivation should be delayed until after corn is growing normally to avoid breakage.
- Avoid using crop oil concentrates after crop emergence as crop injury may result.

Agriculturally approved surfactants or sprayable fertilizers (1/2 to 1 gallon per acre of 28%, 30% or 32% urea ammonium nitrate or 2.5 pounds per acre spray grade ammonium sulfate¹) may be added to the spray mixture to improve postemergence weed control, particularly in dry growing conditions.

[¹Not for use in California.]

Corn may be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity.

Several synthetic pyrethroid insecticides are labeled for tank mix applications of DICAMBA DMA SALT 5. Refer to their label for specific instructions.

WEEDS CONTROLLED

DICAMBA DMA SALT 5 will control many annual broadleaf weeds or give growth suppression of many perennial broadleaf weeds commonly found in corn. (Refer to the WEED LIST.)

For best performance, make application when weeds have emerged and are actively growing.

Preemergence control of cocklebur, velvetleaf, and jimsonweed may be reduced if conditions including low temperature or lack of soil moisture cause delayed or deep germination of weeds.

PREPLANT/PREEMERGENCE IN NO-TILLAGE CORN

Applications of DICAMBA DMA SALT 5 may be made before, during, or after planting to emerged and actively growing broadleaf weeds. Apply DICAMBA DMA SALT 5 at 12.8 fl. oz. per treated acre on medium or fine textured soils containing 2% or greater organic matter. Use 6.4 fl. oz. per treated acre on coarse textured soils (sand, sandy loam, and loamy sand) or medium and fine textured soils with less than 2% organic matter.

When planting into a legume sod (e.g., alfalfa or clover), apply DICAMBA DMA SALT 5 after 4 to 6 inches of regrowth has occurred.

PREEMERGENCE IN CONVENTIONAL OR REDUCED TILLAGE CORN

DICAMBA DMA SALT 5 may be applied after planting and prior to corn emergence. Application at 12.8 fl. oz. per treated acre may be made to medium or fine textured soils which contain 2% or greater organic matter.

Preemergence application of DICAMBA DMA SALT 5 does not require mechanical incorporation to become active. A shallow mechanical incorporation is required if application is not followed by adequate rainfall or sprinkler irrigation. Avoid tillage equipment (e.g., drags, harrows) which concentrate treated soil over seed furrow.

EARLY POSTEMERGENCE (ALL TILLAGE SYSTEMS) (SPIKE THROUGH 8 INCH TALL CORN)

DICAMBA DMA SALT 5 at 12.8 fl. oz. per treated acre may be applied during the period from corn emergence through the five-leaf stage or 8 inches tall, whichever comes first. Reduce the rate to 6.4 fl. oz. per treated acre if corn is growing on coarse textured soils (sand, sandy loam, and loamy sand). See LATE POSTEMERGENCE APPLICATIONS given below if the 6th true leaf is emerging from whorl or corn is greater than 8 inches tall.

LATE POSTEMERGENCE (ALL TILLAGE SYSTEMS) (8 TO 36 INCH TALL CORN)

Application of DICAMBA DMA SALT 5 at 6.4 fl. oz. per treated acre may be made from 8 to 36 inch tall corn or 15 days before tassel emergence, whichever comes first. For best performance, make applications when weeds are less than 3 inches tall.

Make directed spray application when (1) corn leaves prevent proper spray coverage; (2) sensitive crops are growing nearby; (3) tank mixing with 2,4-D.

DO NOT apply DICAMBA DMA SALT 5 when soybeans are growing nearby if any of these conditions exist:

- corn is more than 24 inches tall
- soybeans are more than 10 inches tall
- soybeans have begun to bloom

OVERLAY (SEQUENTIAL) TREATMENTS

DICAMBA DMA SALT 5 may be applied to ground previously treated with one or more of the following herbicides:

| | |
|--------------|---------------|
| acetochlor | halosulfuron |
| alachlor | metolachlor |
| atrazine | paraquat |
| dimethenamid | pendimethalin |
| EPTC | propachlor |
| glyphosate | simazine |

- If applying DICAMBA DMA SALT 5 to ground previously treated with a product containing dicamba do not exceed 0.75 lbs a.e./A per year for all applications.
- Allow at least 2 weeks between applications.

Read and follow label directions for each of the above products.

TANK MIX TREATMENTS FOR CORN

DICAMBA DMA SALT 5 may be tank mixed with one or more of the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, and other restrictions.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

2,4-D
 Acetochlor
 Alachlor
 Atrazine
 Clopyralid
 Dimethenamid
 Glyphosate
 Nicosulfuron
 Paraquat
 Pendamethalin
 Primisulfuron
 Pyridate
 S-Metolachlor
 Simazine

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, APPLICATION RATES, AND OTHER RESTRICTIONS. Consult product labels for rate recommendations for tank mix partners.

SORGHUM (MILO)

Observe all restrictions and precautions, including the reference to crops growing under stress.

This product may be applied preplant, postemergence, or preharvest in sorghum.

Read and follow MIXING AND APPLICATION instructions.

Sorghum Use Restrictions:

- **Maximum Single Application Rate:** DO NOT exceed a total of 6.4 fluid ounces (0.25 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Annual Application Rate:** DO NOT exceed a total of 12.8 fluid ounces (0.5 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Number of Applications per Year:** 2
- **Retreatment Interval:** 7 days.
- **Pre-Harvest Interval:**
 - Sorghum Grain: 30 days
 - Sorghum Fodder: 30 days
 - Sorghum Forage: 20 days
- **DO NOT** graze or feed treated sorghum forage or silage prior to mature grain stage. Restrictions for sorghum that is grazed or cut for hay are indicated in Table 6 in Pasture, Hay, Rangeland, and General Farmstead section of this label.
- **DO NOT** apply this product to sorghum grown for seed production.
- DICAMBA DMA SALT 5 contains 0.625 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.

Sorghum Use Precautions:

- Applications of DICAMBA DMA SALT 5 to sorghum during periods of rapid growth may result in temporary leaning of plants or rolling of leaves. These effects are usually outgrown within 10 to 14 days.

WEEDS CONTROLLED

DICAMBA DMA SALT 5, when applied at the label rate for sorghum, will control many actively growing annual broadleaf weeds and will reduce competition from established perennial broadleaf weeds as well as control their seedlings. (Refer to WEED LIST.)

RATES AND TIMINGS

DICAMBA DMA SALT 5 may be applied to emerged and actively growing weeds at least 15 days prior to planting. Postemergence application of DICAMBA DMA SALT 5 must be made after sorghum is in the spike stage (all sorghum emerged) but before sorghum is 15 inches tall. For best performance, make applications when sorghum is in the 3 to 5 leaf stage and weeds are small (less than 3 inches tall). Use drop pipes (drop nozzles) if sorghum is taller than 8 inches. Keeping the spray off the sorghum leaves and out of the whorl will reduce the likelihood of crop injury and improve spray coverage of weed foliage.

BROADCAST RATE PER TREATED ACRE: 6.4 fl. oz. of this product (1/4 lb. a.e.)

TANK MIX TREATMENTS**DICAMBA DMA SALT 5 PLUS ATRAZINE**

For improved control of emerged, actively growing broadleaf weeds including triazine-resistant species and added suppression of perennial broadleaf weeds, tank mix 6.4 fl. oz. DICAMBA DMA SALT 5 with labeled rates of atrazine per treated acre. For control of grasses (less than 1.5 inches tall), tank mix 6.4 fl. oz. DICAMBA DMA SALT 5 with 2 lbs. a.i. atrazine per treated acre. For best performance and minimal crop injury, make application when sorghum is 3-8 inches tall and when broadleaf weeds are small (less than 6 inches tall). Application of atrazine must be made before sorghum is beyond 12 inches tall. The atrazine rate will depend upon soil texture and length of residual weed control desired. Follow all State and Federal restrictions pertaining to atrazine applications.

DICAMBA DMA SALT 5 PLUS BROX™ 2EC HERBICIDE or BUCTRIL®

For improved control of broadleaf weeds, tank mix 6.4 fl. oz. DICAMBA DMA SALT 5 with labeled rates of BROX™ 2EC Herbicide or Buctril® per treated acre. Make application at 4-leaf to 15-inch tall sorghum. Use drop nozzles to direct spray beneath sorghum leaves when sorghum is greater than 8 inches tall.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from on product, spray drift management from another).

OVERLAY (SEQUENTIAL) TREATMENTS

DICAMBA DMA SALT 5 may be applied to ground previously treated with one or more of the following herbicides registered for use in sorghum:

| HERBICIDE |
|-----------------------|
| Alachlor |
| atrazine ¹ |
| metolachlor |
| propachlor |

¹Maximum use rate for atrazine is determined by soil type, tillage practices, surface residue, and state or local restrictions. Follow the more restrictive requirements when determining the maximum use rate for atrazine.

PREHARVEST USES (FOR USE ONLY IN THE STATES OF TEXAS AND OKLAHOMA)

DICAMBA DMA SALT 5 may be applied for weed suppression any time after the sorghum has reached the soft dough stage. An agriculturally approved surfactant may be used to improve performance. For aerial application use at least 2 gallons of water-based carrier per treated acre.

Do not harvest until 30 days after treatment.

BROADCAST RATE PER TREATED ACRE: 6.4 fl. oz. of this product (0.25 lbs. a.e.)

SMALL GRAINS (WHEAT, BARLEY AND OATS) NOT UNDERSEEDED TO LEGUMES

Observe all use restrictions and precautions. Read and follow cleaning, mixing and application instructions.

Application Timing: Apply this product before, during, or after planting small grains. See specific small grain crop uses below for maximum crop stage. For best performance, apply this product when weeds are in the 2 - 3 leaf stage and rosettes are less than 2" across. Applying this product to small grains during periods of rapid growth may result in crop leaning. This condition is temporary and will not reduce crop yields. Applications to small grains may be made with aerial applications with 1 gallon of water or more per acre. Where dense foliage is present, 2 - 3 gallons of water per acre should be used.

Grazing Restrictions for small grain areas that are grazed or cut for hay are indicated in TIMING RESTRICTIONS FOR LACTATING DAIRY ANIMALS FOLLOWING TREATMENT in Pasture, Hay, Rangeland, and General Farmstead section of this label.

Do not apply within 7 days of harvest for wheat, barley or oat for grain.

If small grains are used for pasture or hay, all restrictions found in the **PASTURE, HAY, RANGELAND AND GENERAL FARMSTEAD (NON-CROPLAND)** section below apply. These restrictions also must be followed:

- Animals cannot be removed from treated area for slaughter prior to 30 days after last application.
- There is no waiting period between treatment and grazing for non-lactating dairy animals.
- Treated areas may not be grazed by lactating dairy animals before 7 days after treatment.
- Do not harvest hay from treated areas before 37 days after treatment.

DICAMBA DMA SALT 5 contains 0.625 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.

Note: Observe all precautions and restrictions on the labels of products used in tank mix treatments.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from on product, spray drift management from another).

WEEDS CONTROLLED

DICAMBA DMA SALT 5 or combinations with listed tank mix partners will provide control or suppression of the annual broadleaf weeds listed below. For improved control of listed weeds, DICAMBA DMA SALT 5 may be applied in a tank mix with other herbicides. Refer to specific crop for tank mix options.

| | |
|--|---|
| Alkanet ¹ | Mustard, Blue (Purple) ¹ |
| Bedstraw, Catchweed ¹ | Mustard, Tansy |
| Bindweed, Field ² | Mustard, Treacle ¹ |
| Buckwheat, Tartary | Mustard, Tumble (Jim Hill) ¹ |
| Buckwheat, Wild | Mustard, Wild ¹ |
| Carpetweed ¹ | Nightshade, Black |
| Chamomile, Corn | Nightshade, Cutleaf ¹ |
| Chervil, Bur ¹ | Nightshade, Silverleaf ² (White Horsenettle) |
| Chickweed, Common ¹ | Pennycress, Field (Fanweed, Frenchweed, Stinkweed) |
| Cockle, Corn | Pigweed, Redroot (Carelessweed) |
| Cockle, Cow | Pigweed, Rough |
| Cocklebur, Common | Pigweed, Tumble |
| Cornflower (Bachelorbutton) ¹ | Pineappleweed ¹ |
| Dandelion, Common ² | Plantain, Broadleaf ² |
| Dock, Curly ² | Poppy, Red Horned ¹ |
| Dragonhead, American ¹ | Puncturevine ¹ |
| Evening Primrose, Cutleaf ¹ | Purslane, Common ¹ |

| | |
|---|---|
| Falseflax, Smallseeded ¹ | Radish, Wild ¹ |
| Fiddleneck (Tarweed) ¹ | Ragweed, Common ¹ |
| Flixweed ¹ | Ragweed, Giant (Buffaloweed) ¹ |
| Fumitory ¹ | Rocket, London ¹ |
| Gromwell, Corn ¹ | Rocket, Yellow ¹ |
| Groundsel, Common ¹ | Salsify (Goatsbeard) ¹ |
| Hempnettle ¹ | Shepherdspurse ¹ |
| Henbit | Smartweed, Green |
| Jacobs Ladder ¹ | Smartweed, Pennsylvania |
| Knawel (German Moss) | Sorrel, Red (Sheep Sorrel) ¹ |
| Knotweed, Prostrate | Sowthistle, Annual |
| Kochia | Starthistle, Yellow ¹ |
| Ladysthumb | Sunflower, Common (Wild) |
| Lambsquarters, Common | Thistle, Canada ² |
| Lettuce, Miners ¹ | Thistle, Russian |
| Lettuce, Prickly | Velvetleaf |
| Mallow, Common | Vetch ¹ |
| Mayweed, Chamomile (Dogfennel) ¹ | Yarrow, Common ² |
| Pepperweed, Peppergrass ¹ | |

¹These weeds will be controlled with DICAMBA DMA SALT 5 tank mixtures. Refer to tank mix label for specific weeds controlled.

²DICAMBA DMA SALT 5 tank mixes will provide suppression of established broadleaf weeds and control of their seedlings.

RATES AND TIMINGS

Application of DICAMBA DMA SALT 5 may be made before, during or after planting of small grains. For best performance, make applications when weeds are in the 2-3 leaf stage and rosettes are less than 2 inches across. Application of DICAMBA DMA SALT 5 to small grains during periods of rapid growth may result in crop leaning. This condition is temporary and will not reduce crop yields.

Use DICAMBA DMA SALT 5 at 1.6 to 3.2 fluid ounces per treated acre in wheat, fall seeded barley, and oats, and at 1.6 to 2.4 fluid ounces per treated acre in spring seeded barley. Use the higher level of listed rate ranges when treating difficult to control weeds including kochia, Russian thistle and prickly lettuce or dense vegetative growth.

Applications to small grains may be made with aerial applications of 1 gallon or more water per acre. Where dense foliage is present, use 2-3 gallons of water per acre.

DICAMBA DMA SALT 5 used in a tank mix with other herbicides offers the best spectrum of weed control and herbicide tolerant or resistant weed management. Refer to specific crop for DICAMBA DMA SALT 5 rate and application timing.

For applications prior to the emergence of weeds or when sulfonylurea-resistant weeds are present or suspected, use a minimum of 2.4 fluid ounces per treated acre of DICAMBA DMA SALT 5 with a tank mix herbicide. Non-sulfonylurea herbicides, including 2,4-D or MCPA tank mixed with DICAMBA DMA SALT 5 will offer more consistent control of sulfonylurea resistant weeds.

When tank mixing with sulfonylurea herbicides, use an agriculturally approved surfactant of at least 80% active ingredient at the rate of 1-4 pints/100 gallons of spray or not more than 0.25-0.5% by volume. Use the highest rate of surfactant when using the lower rate ranges of the tank mix and/or when treating more mature and difficult to control weeds or dense vegetative growth.

WHEAT

FALL AND SPRING SEEDED WHEAT

DICAMBA DMA SALT 5 must be applied to fall seeded wheat prior to the jointing stage. Applications to spring seeded wheat must be made before wheat reaches the 6-leaf stage. Early developing wheat varieties including TAM 107,

Madison, or Wakefield must receive application between early tillering and the jointing stage. Care should be taken in staging these varieties to be certain that the application occurs prior to the jointing stage.

Wheat Use Restrictions:

- **Maximum Single Application Rate: DO NOT** exceed a total of 6.4 fluid ounces (0.25 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Annual Application Rate: DO NOT** exceed a total of 12.8 fluid ounces (0.5 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Number of Applications per Year: 2**
- **Retreatment Interval:** 7 days.
- **Pre-Harvest Interval:** 7 days. Restrictions for wheat that is grazed or cut for hay are indicated in the table titled TIMING RESTRICTIONS FOR LACTATING DAIRY ANIMALS FOLLOWING TREATMENT in the Pasture, Hay, Rangeland, and General Farmstead section of this label.
- **DO NOT** use pre-harvest treated wheat for seed unless a germination test is performed on the seed with an acceptable result of 95% germination or better.
- **DO NOT** make pre-harvest applications in California.

TANK MIX TREATMENTS

DICAMBA DMA SALT 5 may be tank mixed with one or more of the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled, and geographic and other restrictions.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

Apply 1.6 to 3.2 fluid ounces of DICAMBA DMA SALT 5¹ per treated acre in wheat, fall seeded barley and oats with:

2,4-D (DO NOT tank mix this product with 2,4-D in early season applications on spring-seeded barley.)

MCPA

Metsulfuron-methyl

Triasulfuron

Thifensulfuron

Tribenuron-methyl

Chlorsulfuron

Chlorsulfuron

Bromoxynil²

Clopyralid

Diuron^{2, 3}

Metribuzin^{2,3}

Fenoxaprop-ethyl⁴

¹ Early developing wheat varieties including TAM 107, MADISON, OR WAKEFIELD must receive application between early tillering and the jointing stage. Care should be taken in staging these varieties to be certain that the application occurs prior to the jointing stage.

² Herbicides with the same active ingredient and/or different formulations may be used.

³ Tank mixtures for fall seeded wheat only.

⁴ Use 1.6 fluid ounces of DICAMBA DMA SALT 5 only. Do not use if wild oats is the target weed. Do not use DICAMBA DMA SALT 5 as a tank mix treatment on Durum wheat.

Apply 2.4-3.2¹ fluid ounces of DICAMBA DMA SALT 5 per treated acre in spring seeded barley with:

2, 4-D amine or ester⁴

MCPA

metsulfuron-methyl²

triasulfuron²

thifensulfuron²
chlorsulfuron²
tribenuron-methyl²
glyphosate⁵

¹ DICAMBA DMA SALT 5 may be used at 4.8 fluid ounces on fall seeded wheat in western Oregon as a spring application only. In CO, KS, NM, OK and TX up to 8 fluid ounces of DICAMBA DMA SALT 5 may be applied on fall seeded wheat after it exceeds the 3-leaf stage for suppression of perennial weeds, including field bindweed. Applications may be made in fall following a frost but before a killing freeze. DICAMBA DMA SALT 5 may be tank mixed with 2,4-D amine after wheat begins to tiller. Periods of extended stress, including cold and wet weather, may enhance the possibility of crop injury. For fall applications only, do not use if the potential for crop injury is not acceptable.

² Do not use low rates of sulfonylurea herbicides on more mature weeds and/or on dense vegetative growth.

³ NOTE: For use on Fall Seeded Wheat Only. Do not use unless potential crop injury will be acceptable.

⁴ Use for improved control of Russian thistle, flxweed, gromwell, mayweed and fiddleneck.

⁶ DICAMBA DMA SALT 5 may be applied at 1.6 fluid ounces with any glyphosate formulation labeled for use as a preplant application to small grains with no waiting period prior to planting. Read and follow label directions of the tank mix product for adjuvant use instructions.

BARLEY

Preharvest applications: This product can be used to control weeds that may interfere with harvest of fall- and spring-seeded barley. Apply 6.4 fluid ounces of this product per acre as a broadcast or spot treatment to annual broadleaf weeds when barley is in the hard dough stage and the green color is gone from the nodes (joints) of the stem. Best results will be obtained if application can be made when weeds are actively growing, but before weeds canopy.

Barley Use Restrictions:

- **Maximum Single Application Rate: DO NOT** exceed a total of 6.4 fluid ounces (0.25 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Annual Application Rate: DO NOT** exceed a total of 8.8 fluid ounces (0.34 lbs a.e.) of this product per treated acre, per crop year to spring seeded barley.
- **Maximum Annual Application Rate: DO NOT** exceed a total of 9.6 fluid ounces (0.38 lbs a.e.) of this product per treated acre, per crop year to fall seeded barley.
- **Maximum Number of Applications per Year: 2**
- **Retreatment Interval:** 7 days.
- **Pre-Harvest Interval:** 7 days. Restrictions for small grain areas that are grazed or cut for hay are indicated in the table titled TIMING RESTRICTIONS FOR LACTATING DAIRY ANIMALS FOLLOWING TREATMENT in the Pasture, Hay, Rangeland, and General Farmstead section of this label.

FALL SEEDED BARLEY

DICAMBA DMA SALT 5 must be applied to fall seeded barley prior to the jointing stage.

NOTE: For spring seeded barley varieties that are seeded during the winter months or later, follow the rates and timings given for spring seeded barley.

TANK MIX TREATMENTS

DICAMBA DMA SALT 5 may be tank mixed with one or more of the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled, geographic and other restrictions.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from on product, spray drift management from another).

BROADCAST RATE PER TREATED ACRE

Apply 1.6-3.2 fluid ounces DICAMBA DMA SALT 5 with:

2,4-D
MCPA
metsulfuron-methyl¹
triasulfuron¹
thifensulfuron¹
tribenuron-methyl¹
chlorsulfuron¹
metribuzin²
bromoxynil

¹ Do not use low rates of sulfonylureas (Ally[®], Amber[®], Express[®], Glean[®], and Harmony[®] Extra) on more mature weeds and/or on dense vegetative growth.

² Herbicides with the same active ingredient and/or different formulations may be used.

SPRING SEEDED BARLEY

DICAMBA DMA SALT 5 must be applied before spring seeded barley exceeds the 4-leaf stage.

TANK MIX TREATMENTS

DICAMBA DMA SALT 5 may be tank mixed with one or more of the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled, geographic and other restrictions.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

BROADCAST RATE PER TREATED ACRE

Apply 1.6-3.2 fluid ounces of DICAMBA DMA SALT 5 with:

MCPA
metsulfuron-methyl¹
triasulfuron¹
thifensulfuron¹
tribenuron-methyl¹
chlorsulfuron¹
metribuzin²
bromoxynil

¹ Do not use low rates of sulfonylureas on more mature weeds and/or on dense vegetative growth.

² Herbicides with the same active ingredient and/or different formulations may be used.

OATS

FALL AND SPRING SEEDED OATS

DICAMBA DMA SALT 5 must be applied before spring seeded oats exceed the 5-leaf stage. Applications to fall seeded oats must be made prior to the jointing stage. **DO NOT** tank mix this product with 2,4-D in oats.

Oat Use Restrictions:

- **Maximum Single Application Rate:** **DO NOT** exceed a total of 3.2 fluid ounces (0.125 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Annual Application Rate:** **DO NOT** exceed a total of 3.2 fluid ounces (0.125 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Number of Applications per Year:** 1
- **Retreatment Interval:** N/A
- **Pre-Harvest Interval:** 7 days. Restrictions for small grain areas that are grazed or cut for hay are indicated in the table titled TIMING RESTRICTIONS FOR LACTATING DAIRY ANIMALS FOLLOWING TREATMENT in the Pasture, Hay, Rangeland, and General Farmstead section of this label.
- **DO NOT** tank mix this product with 2,4-D in oat.

TANK MIX TREATMENTS

DICAMBA DMA SALT 5 may be tank mixed with one or more of the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled, geographic and other restrictions.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from on product, spray drift management from another).

BROADCAST RATE PER TREATED ACRE

Apply 1.6-3.2 fluid ounces of DICAMBA DMA SALT 5 with: label rate of MCPA

PREHARVEST APPLICATIONS FOR BARLEY AND WHEAT:

DICAMBA DMA SALT 5 can be used to control weeds that may interfere with harvest of fall- and spring-seeded barley and wheat.

Apply 6.4 fluid ounces of DICAMBA DMA SALT 5 per acre as a broadcast or spot treatment to annual broadleaf weeds when barley or wheat is in the hard dough stage and the green color is gone from the nodes (joints) of the stem. Best results will be obtained if application can be made when weeds are actively growing, but before weeds canopy.

A waiting interval of 7 days is required before harvest.

DO NOT use pre-harvest treated barley or wheat for seed unless a germination test is performed on the seed with an acceptable result of 95% germination or better.

For control of additional broadleaf weeds or grasses, DICAMBA DMA SALT 5 may be tank mixed with other herbicides, including 2,4-D, that are labeled for pre-harvest uses in barley or wheat.

DO NOT make pre-harvest applications in California.

SUGARCANE

Observe all use restrictions and precautions. Read and follow MIXING AND APPLICATION instructions.

Sugarcane Use Restrictions:

- **Maximum Single Application Rate: DO NOT** exceed a total of 25.6 fluid ounces (1.0 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Annual Application Rate: DO NOT** exceed a total of 51.2 fluid ounces (2.0 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Number of Applications per Year: 2**
- **Retreatment Interval: 7 days.**
- **Pre-Harvest Interval: 87 days.** Restrictions for sugarcane that is grazed or cut for hay are indicated in the table titled TIMING RESTRICTIONS FOR LACTATING DAIRY ANIMALS FOLLOWING TREATMENT in the Pasture, Hay, Rangeland, and General Farmstead section of this label.

Consult your local or state authorities for possible application restrictions, especially concerning aerial applications and advice concerning special local use situations.

WEEDS CONTROLLED

DICAMBA DMA SALT 5, when applied at the label rates, will control many annual, biennial and perennial broadleaf weeds commonly found in sugarcane. (Refer to WEED LIST.)

RATES AND TIMINGS

Application of DICAMBA DMA SALT 5 may be made any time after weeds have emerged and are actively growing but before the close-in stage of sugarcane. Application rates and timings of DICAMBA DMA SALT 5 are given below. Use the higher level of listed rate ranges when treating dense vegetative growth.

| Weed Stage & Type | Product Amount | Broadcast Rate Per Treated Acre (lbs. a.i.) |
|-------------------|----------------|---|
|-------------------|----------------|---|

| | | |
|-------------------------|-------------------|---------|
| Annual | | |
| Small, actively growing | 6.4-12.8 fl. oz. | 1/4-1/2 |
| Established growth | 12.8-19.2 fl. oz. | 1/2-3/4 |
| Biennial | 12.8-25.6 fl. oz. | 1/2-1 |
| Perennial | 25.6-51.2 fl. oz. | 1-2* |

*Application made over the top of actively growing sugarcane may result in crop injury.

When possible, direct the spray beneath the sugarcane canopy in order to minimize the likelihood of crop injury. The use of directed sprays will also aid in maximizing spray coverage of weed foliage.

TANK MIX TREATMENTS

DICAMBA DMA SALT 5 may be tank mixed with one or more of the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, weeds controlled, geographic and other restrictions.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

| HERBICIDE |
|-----------|
| ametryn |
| asulam |
| atrazine |
| 2,4-D* |

*Application of DICAMBA DMA SALT 5 plus 2,4-D tank mix at the higher listed ranges may result in crop injury.

PASTURE, HAY, RANGELAND AND GENERAL FARMSTEAD (NON-CROPLAND)

DICAMBA DMA SALT 5 is labeled for use on pasture, hay, rangeland, general farmstead (non-cropland including fence rows and non-irrigation ditchbanks) for broadleaf weed and brush control. DICAMBA DMA SALT 5 may also be applied to non-cropland areas for the control of broadleaf weeds in noxious weed control programs, districts or areas including broadcast or spot treatment of roadsides and highways, utilities, railroad and pipeline rights-of-way. Noxious weeds must be recognized at the state level but programs may be administered at state, county or other level.

DICAMBA DMA SALT 5 uses described in this section also pertain to small grains (including barley, forage sorghum, oats, rye, sudangrass or wheat) grown for pasture use only.

Observe all use restrictions and precautions. Read and follow mixing and application instructions.

Pasture, Hay, Rangeland, and General Farmstead (noncropland) Restrictions:

- **Maximum Single Application Rate:** DO NOT exceed a total of 25.6 fluid ounces (1.0 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Annual Application Rate:** DO NOT exceed a total of 25.6 fluid ounces (1.0 lbs a.e.) of this product per treated acre, per crop year.
- **DO NOT** apply more than 12.8 fluid ounces (0.5 lbs a.e.) of this product per acre to small grains grown for pasture.
- **Maximum Number of Applications per Year:** 1
- **Retreatment Interval:** N/A
- **Pre-Harvest Interval:** 7 days. Restrictions for sugarcane that is grazed or cut for hay are indicated in the table titled TIMING RESTRICTIONS FOR LACTATING DAIRY ANIMALS FOLLOWING TREATMENT in the Pasture, Hay, Rangeland, and General Farmstead section of this label

Newly seeded areas, including small grains grown for pasture may be severely injured if rates of DICAMBA DMA SALT 5 greater than 12.8 fl. oz./A are applied.

Established grass crops growing under stress can exhibit various injury symptoms that may be more pronounced if herbicides are applied. Furthermore, rates of DICAMBA DMA SALT 5 in excess of 51.2 fl. oz. (2 lbs. a.e.) per treated acre may cause temporary injury to many grass species.

Bentgrass, carpetgrass, buffalograss and St. Augustine grass may be injured at rates exceeding 12.8 fl. oz. DICAMBA DMA SALT 5 (0.5 lb. a.e.) per treated acre. Usually colonial bentgrasses are more tolerant than creeping types. Velvetgrasses are most easily injured. Treatments will kill or injure alfalfa, clovers, lespedeza, wild winter peas, vetch and other legumes.

Animals cannot be removed from treated area for slaughter prior to 30 days after last application. There is no waiting period between treatment and grazing for non-lactating animals.

TIMING RESTRICTIONS FOR LACTATING DAIRY ANIMALS FOLLOWING TREATMENT

| DICAMBA DMA SALT 5 Rate per Treated Acre | Days Before Grazing | Days Before Hay Harvest |
|---|---------------------|-------------------------|
| Up to 12.8 fl. oz. (0.5 lb. a.e.) | 7 days | 37 days |
| Up to 25.6 fl. oz. (1.0 lb. a.e.) | 21 days | 51 days |
| Up to 51.2 fl.oz. (2.0 lbs. a.e.) | 40 days | 70 days |

NOTE: Observe all precautions and restrictions on labels of products used in tank mixtures.

MIXING AND APPLICATION

Tank Mixture Instructions: DICAMBA DMA SALT 5 contains 0.625 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.

DICAMBA DMA SALT 5 can be applied using water, oil in water emulsions including invert systems, or sprayable fluid fertilizer as a carrier. A compatibility test (see COMPATIBILITY TEST section) should be made prior to tank mixing.

To prepare oil in water emulsions, half-fill spray tank with water, then add the appropriate amount of emulsifier with continuous agitation. Slowly add the herbicide and then the oil (including diesel oil or fuel oil) or a premix of oil plus additional emulsifier to spray tank. Complete filling of spray tank with water. Maintain vigorous agitation during spray operation to prevent oil and water from forming separate layers.

Read and follow the label of each tank mix product used for precautionary statements, directions for use, application rates and other restrictions.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from on product, spray drift management from another).

DICAMBA DMA SALT 5 may be tank mixed with one or more of the following herbicides for control of grasses, additional broadleaf weeds, and woody brush and vines.

| HERBICIDE |
|---|
| Pasture, hay, rangeland and general farmstead (non-cropland) use: |
| glyphosate |
| metsulfuron methyl |
| paraquat |
| picloram |
| tricolopyr |
| 2,4-D |

Due to variations that may occur in formulated products and specific use ingredients (e.g. water supplies), a compatibility test (see COMPATIBILITY TEST section) should be performed prior to actual tank mixing if applicator has no prior experience with that tank mix.

Application Instructions: DICAMBA DMA SALT 5 may be applied broadcast using either ground or aerial application equipment. When using ground equipment, apply 3 to 600 gallons of diluted spray per treated acre. Volume of spray applied will depend on the height, density, and type of weeds or brush being treated and on the type of equipment being used. When using aerial equipment, apply 2 to 40 gallons of diluted spray per treated acre in a water-based carrier.

DICAMBA DMA SALT 5 may be applied to individual clumps or small areas (spot treatment) of undesirable vegetation using handgun or similar types of application equipment. Apply diluted sprays to allow complete wetting (up to run-off) of foliage and stems.

Herbicide adjuvants or other spray additives (emulsifiers, surfactants, wetting agents, drift control agents, or penetrants) may be used for wetting, penetration, or drift control. Spray additives must be agriculturally approved when used in pasture applications. If spray additives are used, read and follow all use instructions and precautions on product label.

WEEDS CONTROLLED

DICAMBA DMA SALT 5, when applied at label rates, will give control of many annual, biennial, and perennial broadleaf weeds, and many woody brush and vine species commonly found in pasture, hay, rangeland and general farmstead (non-cropland) areas. (Refer to WEED LIST.) Perennial weeds noted with an asterisk (*) may be controlled with lower rates of either DICAMBA DMA SALT 5 or DICAMBA DMA SALT 5 plus 2,4-D. See RATES AND TIMINGS below.

RATES AND TIMINGS

Application rates and timing of DICAMBA DMA SALT 5 are given below. Use the higher level of listed rate ranges when treating dense or tall vegetative growth.

| WEED STAGE & TYPE | PRODUCT AMOUNT | BROADCAST RATE PER TREATED ACRE (LBS. A.I.) |
|--|-------------------|---|
| Annual Small, actively growing | 6.4-12.8 fl. oz. | 1/4-1/2 |
| Established weed growth | 12.8-19.2 fl. oz. | 1/2-3/4 |
| Biennial ¹ Rosette diameter Less than 3 inches | 6.4-12.8 fl. oz. | 1/4-1/2 |
| 3 inches or more | 12.8-25.6 fl. oz. | 1/2-1 |
| Bolting | 25.6-38.4 fl. oz. | 1-1 1/2 |
| Perennial Suppression or top growth control Noted (*) Perennials | 12.8-25.6 fl. oz. | 1/2-1 |
| | 25.6 fl. oz. | 1-2* |
| Other Perennials | 51.2 fl. oz. | 2* |
| Woody Brush & Vines Top growth suppression | 12.8-25.6 fl. oz. | 1/2-1 |
| Top growth control ² | 25.6-51.2 fl. oz. | 1-2* |
| Stems and stem suppression | 51.2 fl. oz. | 2* |

¹For best performance, make application when biennial weeds are in the rosette stage.

²Species noted in WEED LIST section will require tank mixtures for adequate control.

*Rates of 2.0 lb. a.i./A are spot treatments only. Do not broadcast apply more than 1.0 lb. a.e./A.

Retreatments may be made as needed; however, do not exceed a total of 51.2 fl. oz (2.0 lbs. a.e.) of DICAMBA DMA SALT 5 per treated acre per year.

OTHER APPLICATION INSTRUCTIONS:

Cut surface tree treatments: DICAMBA DMA SALT 5 may be applied as a cut surface treatment for control of unwanted trees and prevention of sprouts of cut trees. A mix ratio of 3.2 fluid ounces of DICAMBA DMA SALT 5 with 4 to 12 fluid ounces of water must be used in application. Use the lower dilution when treating difficult-to-control species.

FRILL OR GIRDLE TREATMENTS: Make a continuous cut or a series of overlapping cuts using an axe to girdle tree trunk. Spray or paint cut surface with the DICAMBA DMA SALT 5/water mix.

STUMP TREATMENTS: Spray or paint freshly cut surface with the water mix. The area adjacent to the bark should be thoroughly wet.

NOTE: For more rapid foliar effects, 2,4-D may be added to the DICAMBA DMA SALT 5/water mix.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank

mixing. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

Dormant applications for control of multiflora rose: DICAMBA DMA SALT 5 can be applied when plants are dormant as an undiluted SPOT-CONCENTRATE directly to the soil or as a LO-OIL BASAL BARK treatment using an oil-water emulsion solution.

SPOT-CONCENTRATE applications of DICAMBA DMA SALT 5S: Apply directly to the soil as close as possible to the root crown but within 6-8 inches of the crown. On sloping terrain, make the application to the uphill side of the crown. Do not make application when snow or water prevents applying DICAMBA DMA SALT 5 directly to the soil. The use rate of DICAMBA DMA SALT 5 is dependent on the canopy diameter of the multiflora rose. Examples: Use DICAMBA DMA SALT 5 at 0.2, 0.8 or 1.8 fluid ounces of product respectively, for 5, 10, or 15 feet canopy diameters. Do not exceed a total of 51.2 fl. oz. of DICAMBA DMA SALT 5 per acre per year.

LO-OIL BASAL BARK application of DICAMBA DMA SALT 5: Apply to the basal stem regions from the ground line up to a height of 12 to 18 inches. Spray until runoff, with special emphasis on covering the root crown. For best results, make application when plants are dormant. Do not make application after bud break or when plants are showing signs of active growth. Do not make application when snow or water prevents applying DICAMBA DMA SALT 5 to the ground line. Refer to MIXING AND APPLICATIONS above in this section for method of preparing oil-in-water emulsion. Example for making approximately 2 gallons of a LO-OIL spray mixture: combine 1 ½ gallons water plus 1 ounce emulsifier plus 12.8 fl. oz. DICAMBA DMA SALT 5 plus 2 ½ pints of No. 2 diesel fuel. Adjust amounts of materials used proportionately to the amount of final spray solution desired. Do not exceed 8 gallons of spray solution mix applied per acre per year.

Conservation Reserve Program (CRP) acres: DICAMBA DMA SALT 5 can be used on both newly seeded and established grasses grown in Conservation Reserve or Federal Set-Aside Programs.

DICAMBA DMA SALT 5 treatment will injure or may kill alfalfa, clovers, lespedeza, wild winter peas, vetch, and other legumes.

Agriculturally approved surfactants may be added to the spray mixture to improve postemergence weed control, particularly in dry growing conditions.

Do not use adjuvants containing penetrants including petroleum based oils after grass emergence on newly seeded grasses.

DICAMBA DMA SALT 5 contains 0.625 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.

Retreatments may be made as needed however, do not exceed a total of 51.2 fluid ounces of this product (2.0 lbs a.e.) per acre.

Newly seeded areas: DICAMBA DMA SALT 5 may be applied either preplant or postemergence to newly seeded grasses or small grains including barley, oats, rye, sudangrass, or wheat grown as a cover crop. Postemergence applications may be made after seedling grasses exceed the 3-leaf stage. Rates of DICAMBA DMA SALT 5 greater than 12.8 fl. oz. per treated acre may severely injure newly seeded grasses.

Preplant applications: Injury to new seedlings may occur if intervals between application and grass planting are less than 45 days per pint of DICAMBA DMA SALT 5 per treated acre west of the Mississippi River or 20 days per 12.8 fl. oz. east of the Mississippi River.

Established grass stands: Established grass stands are perennial grasses planted one or more seasons prior to treatment. Certain species, bentgrass, carpetgrass, smooth brome, buffalograss or St. Augustine grass, may be injured when treated with DICAMBA DMA SALT 5 at rates exceeding 12.8 fl. oz. per treated acre.

DICAMBA DMA SALT 5, when applied at label rates, will control many annual and biennial weeds and provide control or suppression of many perennial weeds. (Refer to WEED LIST.)

Application rates and timings of DICAMBA DMA SALT 5 treatments are given below. Use the higher rate of the rate range when vegetation is either dense or tall, or when weeds are growing under stressed conditions including drought or cool temperature.

| Weed Type* & Stage | Broadcast Rate Per Treated Acre | |
|--|---|-----------------------------|
| | Amount of formulated DICAMBA DMA SALT 5 (fl. oz.) | Equivalent lbs. a.e. |
| Annuals Small, actively growing Established weed growth | 3.2-12.8 12.8 | 1/8-1/2 1/2 |
| Biennials** Rosette diameter Less than 3 inches 3 inches or greater Bolting biennial | 6.4-12.8 12.8-25.6 25.6-38.4 | 1/4-1/2 1/2-1 1-1 1/2 |
| Perennials** Suppression/Control | 25.6-51.2 | 1-2 |

*For best results, treat biennial weeds with DICAMBA DMA SALT 5 when they are in the rosette stage of growth. Retreatments may be made as needed; however, do not exceed a total of 51.2 fl. oz. (2.0 lbs. a.e.) of DICAMBA DMA SALT 5 per treated acre per year.

**biennial and perennial weeds will require follow-up (sequential) treatments for seedling control and escapes.

TANK MIX TREATMENTS

To control grasses and additional broadleaf weeds, DICAMBA DMA SALT 5 may be tank mixed with other herbicides registered for use in Conservation Reserve Programs including 2,4-D, glyphosate (Gly Star™ Original or Roundup®), paraquat (Gramoxone®), metsulfuron (Ally®) and others.

Read and follow the label of each tank mix product used for precautionary statements, directions for use, application rates, and other restrictions.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from on product, spray drift management from another).

ASPARAGUS

FOR USE ONLY IN THE STATES OF CALIFORNIA, OREGON AND WASHINGTON

Observe all use restrictions and precautions. Read and follow mixing and application instructions.

If spray contacts emerged spears, crooking (twisting) of some spears may result. If such crooking occurs, discard affected spears.

Asparagus Use Restrictions:

- **Maximum Single Application Rate:** DO NOT exceed a total of 12.8 fluid ounces (0.5 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Annual Application Rate:** DO NOT exceed a total of 12.8 fluid ounces (0.5 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Number of Applications per Year:** 2
- **Retreatment Interval:** 14 days
- **Pre-Harvest Interval:** DO NOT harvest prior to 24 hours after treatment.
- **DO NOT** use in the Coachella Valley of California.

DICAMBA DMA SALT 5 contains 0.625 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.

RATES AND TIMINGS

Apply DICAMBA DMA SALT 5 to emerged and actively growing weeds in 40 to 60 gallons of diluted spray per treated acre immediately after cutting the field, but at least 24 hours before the next cutting.

| WEEDS | RATE PER TREATED ACRE |
|---|---------------------------------------|
| Mustard, Black Pigweed, Redroot (Carelessweed) Sowthistle, Annual | 6.4-12.8 fl. oz (1/4-1/2 lb. a.e.) |

| | |
|---|--------------------------------|
| *Thistle, Canada Thistle, Russian | |
| *Bindweed, Field Chickweed, Common Goosefoot, Nettleleaf Radish, Wild Thistle, Milk | 12.8 fl. oz. (1/2 lb. a.e.) |

DICAMBA DMA SALT 5 may be applied in a tank mixture with either 2,4-D, Gly Star™ Original or Roundup® for improved control of noted (*) weeds. Read and follow 2,4-D, Gly Star™ Original or Roundup® product labeling for precautionary statements, directions for use, application rates and timings, and other restrictions.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from on product, spray drift management from another).

TURF AND LAWNS

Including Golf Course (Fairways, Aprons, Tees, and Rough), Parks, Recreational Areas, Lawn Care application, and Sod farms.

Observe all use restrictions and precautions. Read and follow mixing and application instructions.

Turf and Lawns Use Restrictions:

- **Maximum Single Application Rate:** DO NOT exceed a total of 25.6 fluid ounces (1.0 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Annual Application Rate:** DO NOT exceed a total of 25.6 fluid ounces (1.0 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Number of Applications per Year:** 2
- **Retreatment Interval:** 7 days.
- **Pre-Harvest Interval:** N/A

To avoid injury to newly seeded grasses, application of DICAMBA DMA SALT 5 should be delayed until after the second mowing. Furthermore, application rates in excess of 12.8 fl. oz. (0.5 lb. a.e.) per treated acre may cause noticeable stunting or discoloration of sensitive grass species including bentgrass, carpetgrass, buffalograss, and St. Augustine grass.

In areas where roots of sensitive plants extend, do not apply in excess of 3.2 fl. oz. (0.125 lb. a.e.) of DICAMBA DMA SALT 5 per treated acre on coarse textured (sandy-type) soils, or in excess of 6.4 fl. oz. (0.25 lb. a.e.) per treated acre on fine textured (clay-type) soils. Do not make repeat applications in these areas for 30 days and until previous applications of DICAMBA DMA SALT 5 have been activated in the soil by rain or irrigation.

DICAMBA DMA SALT 5 contains 0.625 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.

WEEDS CONTROLLED

DICAMBA DMA SALT 5, when applied at label rates, will give control of many annual, biennial, and noted (*) perennial broadleaf weeds commonly found in turf. DICAMBA DMA SALT 5 will also give growth suppression of many other listed perennial broadleaf weeds and woody brush and vine-species. (Refer to WEED LIST.)

MIXING AND APPLICATION

Apply 30 to 200 gallons of diluted spray per treated acre (3 qts. to 4 ¼ gals. per 1,000 sq. ft.), depending on density or height of weeds treated and on the type of equipment used.

RATES AND TIMINGS

Use the higher level of listed rate ranges when treating dense vegetative growth.

| Weed Stage & Type | DICAMBA DMA SALT 5 | |
|-------------------|--------------------------|------------------------------|
| | Fl. Oz. per treated acre | Pounds a.e. per treated acre |
| | | |

| | | |
|--|-----------------------|----------------------|
| Annuals Small, actively growing Established weed growth | 6.4-12.8 12.8-19.2 | 0.25-0.5 0.5-0.75 |
| Biennials, Rosette diameter Less than 3 inches 3 inches or more | 6.4-12.8 12.8-25.6 | 0.25-0.5 0.5-1.0 |
| Perennials, Woody Brush and Vines | 12.8-25.6 | 0.5-1.0 |

For best performance, apply when weeds are emerged and actively growing.

Retreatments may be made as needed; do not exceed a total of 25.6 fl. oz. (1.0 lb. a.e.) DICAMBA DMA SALT 5 per treated acre per year.

TANK MIX TREATMENTS

Read and follow the label of each tank mix product used for precautionary statements, directions for use, application rates and timings and other restrictions.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another). Read and follow the applicable restrictions and limitation and directions for use on all product labels involved in tank mixing.

Tank mix treatments of DICAMBA DMA SALT 5 may be made with 2,4-D, MCPA, MCPP, or bromoxynil for control of additional weeds listed on the tank mix product label.

Apply 2.6 to 6.4 fl. oz. (0.1 to 0.25 lb. a.e.) of DICAMBA DMA SALT 5 per treated acre with labeled rates of 2,4-D, MCPA, MCPP, or bromoxynil. Use the higher level of the listed rate ranges when treating established weeds. Repeat treatments may be made as needed; however, do not exceed 25.6 fl. oz. (1.0 lb. a.e.) of DICAMBA DMA SALT 5 per treated acre per year.

GRASS SEED CROPS

GRASSES GROWN FOR SEED INCLUDING BERMUDAGRASS, BLUEGRASS, FESCUE AND RYEGRASS

Observe all use restrictions and precautions. Read and follow mixing and application instructions.

Grass Grown For Seed Use Restrictions:

- **Maximum Single Application Rate:** **DO NOT** exceed a total of 25.6 fluid ounces (1.0 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Annual Application Rate:** **DO NOT** exceed a total of 51.2 fluid ounces (2.0 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Number of Applications per Year:** 2
- **Retreatment Interval:** 7 days.
- **Pre-Harvest Interval:** N/A. Refer to the Pasture, Hay, Rangeland and General Farmstead section for grazing and feeding restrictions.
- **DO NOT** apply this product after the grass seed crop begins to joint.

Refer to the PASTURE, HAY, RANGELAND, AND GENERAL FARMSTEAD (NON-CROPLAND) section for possible grazing and feeding restrictions.

Do not use on bentgrass unless possible crop injury can be tolerated.

DICAMBA DMA SALT 5 contains 0.625 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.

WEEDS CONTROLLED

DICAMBA DMA SALT 5 will provide control or suppression of annual broadleaf weeds listed below. For improved control of listed weeds plus additional weeds, DICAMBA DMA SALT 5 may be applied in a tank mix with other herbicides.

Alfalfa¹
Bedstraw, Catchweed
Bindweed, Field
Buttercup, Corn
Buttercup, Creeping
Buttercup, Western Field
Catchfly, Nightflowering
Chamomile, Corn
Chickweed, Common
Chickweed, Mouseear
Clover
Cockle, White
Dock, Broadleaf
Dock, Curly
¹Top growth only.

Hemlock, Poison
Knapweed, Russian¹
Knawel
Knotweed, Prostrate
Kochia
Ladysthumb
Lambsquarters, Common
Lettuce, Prickly
Mayweed (Dogfennel)
Ragwort, Tansy
Sorrel, Red (Sheep Sorrel)
Sowthistle, Annual
Starwort, Little
Thistle, Canada¹

RATES AND TIMINGS

Apply 6.4 to 12.8 fl. oz. of DICAMBA DMA SALT 5 per treated acre on seedling grass after the crop reaches the 3 to 5 leaf stage. Apply up to 25.6 fl. oz. of DICAMBA DMA SALT 5 on well-established perennial grass. Do not apply after the grass seed crop begins to joint. For best performance, make applications when weeds are in the 2 to 4 leaf stage and rosettes are less than 2 inches across. Use the higher level of listed rate ranges when treating more mature weeds or dense vegetative growth.

TANK MIX TREATMENTS

For control of grasses or additional broadleaf weeds, DICAMBA DMA SALT 5 may be tank mixed with all broadleaf herbicides registered for use in Grass Seed Production. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled and geographic and other restrictions.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from on product, spray drift management from another).

BROADCAST RATE PER TREATED ACRE

Apply 6.4-25.6 fl. oz. DICAMBA DMA SALT 5 with:

2,4-D
MCPA
bromoxynil
clopymidifen
diuron
clopymidifen

ANNUAL GRASS CONTROL

For suppression of annual grass weeds including:

Brome, Downy (Cheatgrass)
Brome, Ripgut
Fescue, Rattail
Windgrass

Apply up to 51.2 fl. oz. of DICAMBA DMA SALT 5 per treated acre in the fall or late summer after harvest and burning of established grass seed crops. Applications should be made immediately following the first irrigation when the soil is moist and before weeds have more than 2 leaves.

PREPLANT DIRECTIONS (POST HARVEST/FALLOW/CROP STUBBLE/SET-A-SIDE) FOR BROADLEAF WEED CONTROL BEFORE WHEAT, CORN, SORGHUM, SOYBEANS

Observe all precautions. Read and follow mixing and application instructions.

WEEDS CONTROLLED

DICAMBA DMA SALT 5 may be applied alone or in tank mix combinations with other herbicides registered for this use.

DICAMBA DMA SALT 5 can be applied either post harvest in the fall, spring or summer, during the fallow period or to crop stubble/set-a-side acres. DICAMBA DMA SALT 5, when applied at the label rates, will control many annual broadleaf weeds. See the WEEDS CONTROLLED section under small grains. In addition, DICAMBA DMA SALT 5 will control or suppress the following biennial and perennial broadleaf weeds:

| | |
|---|------------------------------------|
| Alfalfa ¹ | Knapweed, Diffuse |
| Artichoke, Jerusalem | Nightshade, Silver |
| Bindweed, Field | Redvine |
| Bindweed, Hedge | Smartweed, Swamp |
| Blueweed, Texas | Sowthistle, Perennial ¹ |
| Bursage (Bur Ragweed, Povertyweed, Lakeweed) ¹ | Spurge, Leafy |
| Dandelion, Common ¹ | Thistle, Bull |
| Dock, Curly ¹ | Thistle, Canada ² |
| Dogbane, Hemp | Thistle, Milk |
| Garlic, Wild ² | Thistle, Musk |
| Horsenettle, Carolina | Thistle, Plumeless |
| Knapweed, Spotted | Thistle, Scotch |
| | Trumpet creeper (Buckvine) |

¹Perennials may be controlled using DICAMBA DMA SALT 5 at rates lower than those for other listed perennial weeds. (See RATES AND TIMINGS under this heading).

²See the SPECIAL TANK MIX TREATMENTS section under this heading for specific control programs for these weeds.

RATES AND TIMINGS

Apply DICAMBA DMA SALT 5 as a broadcast or spot treatment to emerged and actively growing weeds after crop harvest (post harvest) and before a killing frost or in the fallow cropland or crop stubble the following spring or summer.

Agriculturally approved spray additives, including surfactants or oils, may be used to enhance spray coverage and the herbicide's penetration of weed foliage. See CROPPING RESTRICTIONS for required interval between application and planting to prevent crop injury.

For best performance, make application when annual weeds are less than 6 inches tall, when biennial weeds are in the rosette stage, and to perennial weed regrowth in late summer or fall following a mowing or tillage treatment. Most effective control of upright perennial broadleaf weeds, including Canada thistle and Jerusalem artichoke, occurs if application is made when the majority of weeds, including field bindweed and hedge bindweed, are in or beyond the full bloom stage.

Avoid disturbing treated areas following application. Treatments may not kill weeds which develop from seed or underground plant parts, including rhizomes or bulblets, after the effective period for DICAMBA DMA SALT 5. For seedling control, a follow-up program or other cultural practices could be instituted. For small grain in-crop uses of DICAMBA DMA SALT 5, see the RATES AND TIMINGS section under the SMALL GRAINS heading for details.

DICAMBA DMA SALT 5 RATES PER TREATED ACRE

| WEED TYPE | AMOUNT OF PRODUCT PER ACRE |
|-----------------------|-------------------------------|
| Annual | ½-1 pt. (6.4 to 12.8 fl. oz.) |
| Biennial | (12.8 to 25.6 fl. oz.) |
| Perennial | (12.8 to 51.2 fl. oz.) |
| Perennial suppression | (12.8 to 25.6 fl. oz.) |
| Noted (1) perennials | (25.6 to 51.2 fl. oz.) |
| Other perennials | (51.2 fl. oz.) |

Retreatments may be made as needed; however, do not exceed a total of 51.2 fl. oz. of DICAMBA DMA SALT 5 per treated acre during any given period.

TANK MIX TREATMENTS

DICAMBA DMA SALT 5 may be tank mixed with one or more of the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, weeds controlled, geographic and other restrictions.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another). Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing.

DICAMBA DMA SALT 5 BROADCAST RATE PER TREATED ACRE FOR ANNUAL WEED CONTROL

Apply 3.2 to 12.8 fl. oz. DICAMBA DMA SALT 5 with:

2,4-D
Atrazine¹
metsulfuron-methyl²
triasulfuron²
paraquat
glyphosate
chlorsulfuron²
paraquat
pronamide
metribuzin¹

¹ Tank mixes of DICAMBA DMA SALT 5 with these products may be subject to special restrictions. See the product label of the tank mix partner for intended use rates, restrictions and other precautions.

² When tank mixing with sulfonylurea herbicides, refer to the product label for rates and restrictions. Use a surfactant of at least 80% active ingredient at the rate of 1-2 quarts/100 gallons of spray or not more than 0.25-0.5% by volume. Use the highest rate of surfactant when using the lower rate ranges of the tank mix and/or when treating more mature weeds or dense vegetative growth. Sulfonylurea-resistant weeds may not be controlled by tank mixes of DICAMBA DMA SALT 5 and a sulfonylurea. Refer to the DICAMBA DMA SALT 5 tank mix section for alternative tank mixes.

DICAMBA DMA SALT 5 BROADCAST RATE PER TREATED ACRE FOR BIENNIAL AND PERENNIAL WEED CONTROL

Apply 12.8-51.2 fl. oz. of DICAMBA DMA SALT 5 with:

clopyralid
2,4-D
glyphosate
picloram

SPECIAL TANK MIX TREATMENTS

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow the applicable restriction and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

For suppression of perennial weeds, apply 6.4-12.8 fl. oz. of DICAMBA DMA SALT 5 per acre with labeled rates of glyphosate.

For wild garlic control, apply 12.8 fl. oz. DICAMBA DMA SALT 5 per acre with labeled rates of 2,4-D LV ester. Apply when wild garlic is 4 to 8 inches tall.

For Canada thistle control, use DICAMBA DMA SALT 5 with labeled rates of clopyralid or DICAMBA DMA SALT 5 plus glyphosate tank mix treatments.

Application may be made during fallow periods for control of volunteer barley, bulbous bluegrass, downy brome, jointed goatgrass, common rye and volunteer wheat when they are actively growing. Use 12.8 fl. oz. DICAMBA DMA SALT 5 per acre with labeled rates of Kerb[®] 50-W (EPA Reg. No. 62719-397). Fall seeded wheat may be planted 9 months or more after application. For best performance, make application between mid-October and mid-December, prior to soil freeze up.

During fallow periods, apply DICAMBA DMA SALT 5 plus glyphosate herbicide to give improved control of kochia, wild buckwheat, prickly lettuce, field bindweed and Canada thistle.

CROPPING RESTRICTIONS

The following instructions are based on DICAMBA DMA SALT 5 use rates up to 51.2 fl. oz. (2.0 lbs. a.e. per treated acre.)

Corn, sorghum, and soybeans may be planted in the spring following applications made during the previous year. If less than 1 inch of rainfall occurs between application and first killing frost, treated areas should be cultivated to allow herbicide to come in contact with moist soil. Cultivation may take place before or immediately after ground thaw.

Soybean injury may occur if the interval between application and planting is less than specified. In areas with greater than 30 inches of rainfall, delay planting for 30 days per pint of DICAMBA DMA SALT 5 per treated acre. In areas with less than 30 inches of rainfall, delay planting for 45 days per pint of DICAMBA DMA SALT 5 per treated acre. Exclude days when ground is frozen.

Wheat may be planted in the fall or spring following applications. Also, spot application may be made any time prior to crop emergence if crop injury can be tolerated in treated areas. Wheat injury may occur if the interval between application and planting is less than specified.

East of the Mississippi River, the interval is 20 days per 12.8 fl. oz. of DICAMBA DMA SALT 5 per treated acre or 1.25 days per 0.8 ounce. Moisture is essential for DICAMBA DMA SALT 5 degradation. Exclude days when ground is frozen.

West of the Mississippi River, the interval is 45 days per pint of DICAMBA DMA SALT 5 per treated acre or 3 days per 0.8 fl oz. Moisture is essential for DICAMBA DMA SALT 5 degradation. Exclude days when ground is frozen.

Following a normal harvest of barley, oats, or wheat, any rotational crop may be planted. If the interval before harvest is shortened, including when cover crops will be plowed under, do not follow up with the planting of a sensitive crop.

COTTON

PREPLANT APPLICATION

Observe all use restrictions and precautions. Read and follow mixing and application instructions. Refer to the WEED LIST section of this label for a list of weeds controlled or suppressed.

Cotton Use Restrictions:

- **Maximum Single Application Rate:** **DO NOT** exceed a total of 6.4 fluid ounces (0.25 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Annual Application Rate:** **DO NOT** exceed a total of 12.8 fluid ounces (0.50 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Number of Applications per Year:** 2
- **Retreatment Interval:** 7 days
- **Pre-Harvest Interval:** N/A
- **DO NOT** apply preplant to cotton west of the Rockies.
- **DO NOT** make this product preplant applications to cotton in geographic areas with average annual rainfall less than 25".
- If applying a spring preplant treatment following application of a fall preplant (postharvest) treatment, then the combination of both treatments may not exceed 2.0 lbs of Dicamba acid equivalent per acre.
- DICAMBA DMA SALT 5 contains 0.625 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.

RATES AND TIMINGS

Apply DICAMBA DMA SALT 5 as a broadcast or spot treatment to emerged and actively growing weeds at a rate of up to 6.4 fl. oz./acre prior to planting cotton. Most effective control of weeds occurs if application is made when weeds are in the 2-4 leaf stage and rosettes are less than 2" across.

CROPPING RESTRICTIONS

Do not plant cotton for at least 21 days after application and after allowing for a minimum accumulation of 1" of rainfall or overhead irrigation.

Do not apply to cotton west of the Rockies or make preplant applications to cotton in geographic areas with average annual rainfall less than 25".

TANK MIX TREATMENTS

For control of grasses or additional broadleaf weeds, DICAMBA DMA SALT 5 may be tank mixed with Prometryn, Paraquat, and Glyphosate herbicides.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

CONTROL OF PERENNIAL BROADLEAF WEEDS IN FALLOW LAND

(SPOT APPLICATION ONLY)

FOR USE ONLY IN THE STATES OF IDAHO, MONTANA, NEVADA, OREGON, UTAH, AND WASHINGTON

Observe all use restrictions and precautions. Read and follow mixing and application instructions.

Do not treat sub-irrigated non-cropland or areas where the soil remains saturated with water throughout the year.

Make only one application of DICAMBA DMA SALT 5 per year.

DICAMBA DMA SALT 5 contains 0.625 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.

WEEDS CONTROLLED

DICAMBA DMA SALT 5, when applied at label rates, will control many broadleaf weeds including:

| | |
|------------------------------|-------------------|
| Bindweed, Field | Knapweed, Russian |
| Dock, Broadleaf (Bitterdock) | Ragwort, Tansy |
| Dock, Curly | Spurge, Leafy |
| Knapweed, Black | Thistle, Canada |

RATES AND TIMINGS

DICAMBA DMA SALT 5 may be applied at any time following a crop harvest to stubble, or fallow land. Application should be made when weeds are actively growing and prior to a killing frost.

Apply 51.2 fl. oz. (2.0 lbs. a.e.) of DICAMBA DMA SALT 5 per treated acre. Application may be made up to one month prior to the planting of wheat.

NOTE: Do not use unless injury to wheat or rotated barley will be acceptable.

Barley, oats, corn, sorghum (milo), annual or perennial grass crops may be planted into treated areas one year after application. Crops grown for seed (other than perennial grass seed) should not be planted into treated areas until three years after application. Do not plant broadleaf crops including alfalfa, beans, peas, potatoes, or sugar beets into treated areas until two years after application.

In most cases, treatments will not kill perennial weed seedlings which germinate from seed one or two years after treatment. Once the effect of the chemical has been lost, a follow-up program for seedling control or other cultural practices should be instituted.

WIPER APPLICATION USES

Important. Observe all precautions. DICAMBA DMA SALT 5 may be applied through wiper application equipment to control or suppress actively growing broadleaf weeds, brush and vines. Use a solution containing 1 part DICAMBA DMA SALT 5 to 1 part water. Do not contact desirable vegetation with herbicide solution.

Only make wiper application to crops (including PASTURES) and NON-CROPLAND AREAS described in this label with the exception of Cotton, Sorghum, or Soybean.

RIGHTS-OF-WAY, UTILITY AND INDUSTRIAL AREAS, AND FENCEROWS

DICAMBA DMA SALT 5 can be used on non-crop land areas including rights-of-way (roadways, rest areas, utility, railroad, highway, pipeline, and rights-of-way that run through pasture and rangeland); utility facilities (substations, pipelines, tankfarms, pumping stations, parking and storage areas, fencerows, and non-irrigated ditchbanks).

Observe all Use Restrictions and Precautions on this label. Read and follow the Mixing and Application section.

RIGHTS-OF-WAY

DICAMBA DMA SALT 5 can be used to control many broadleaf weeds on rights-of-way. This use includes applications to roadside, roadway and highways; to areas along utilities including cable and powerlines; railroad track and embankment; highways, highway medians, bridge abutments, pipelines, and rights-of-way that run through pasture and rangeland. Use controlled application techniques that minimize the risk of off-target movement.

UTILITY AND INDUSTRIAL AREAS

DICAMBA DMA SALT 5 can be used to control many broadleaf weeds and brush in noncrop areas on or surrounding substations, pipelines, tankfarms, pump stations, production facilities, and bareground situations. It may also be used on parking and storage areas (refer to Best Stewardship Practices to avoid direct runoff from impervious surfaces).

FENCEROWS

DICAMBA DMA SALT 5 can be used to control many broadleaf weeds and brush in fencerows.

Rights-of-way, Utility and Industrial areas, and Fencerows use restrictions:

- **Maximum Single Application Rate:** **DO NOT** exceed a total of 25.6 fluid ounces (1.0 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Annual Application Rate:** **DO NOT** exceed a total of 51.2 fluid ounces (2.0 lbs a.e.) of this product per treated acre, per crop year.
- **Maximum Number of Applications per Year:** 2
- **Retreatment Interval:** 7 days.
- **Pre-Harvest Interval:** N/A

Mixing and Application

Read and observe Management of Off-Site Movement recommendations in this label.

DICAMBA DMA SALT 5 can be applied using water, oil in water emulsions including invert systems, or sprayable fluid fertilizer as a carrier. A compatibility test (see Compatibility Test section) should be made prior to tank mixing.

To prepare oil in water emulsions, half-fill spray tank with water, then add the appropriate amount of emulsifier. With continuous agitation, slowly add the herbicide and then the herbicidal oil or a pre-mix of oil plus additional emulsifier to spray tank. Complete filling of spray tank with water. Maintain vigorous agitation during spray operation to prevent oil and water from forming separate layers.

DICAMBA DMA SALT 5 may be applied broadcast using either ground or aerial application equipment. When using ground equipment, apply low or high volume sprays of between 3 - 600 gals, of diluted spray per treated acre. Volume of spray applied will depend on the height, density, and type of weeds or brush being treated and on the type of equipment being used. When using aerial equipment, apply 5 - 40 gals, of diluted spray per treated acre.

DICAMBA DMA SALT 5 may be applied to individual clumps or small areas (spot treatment) of undesirable vegetation using handgun or similar types of application equipment. Apply diluted sprays to allow complete wetting (up to runoff) of foliage and stems.

Herbicide adjuvants or other spray additives (emulsifiers, spreader stickers, surfactants, wetting agents, drift control agents, or penetrants) may be used for wetting, penetration, or drift control. Spray additives must be agriculturally approved when used in pasture applications. If spray additives are used, read and follow all use recommendations and precautions on product label.

WEEDS AND BRUSH CONTROLLED

DICAMBA DMA SALT 5, when applied at specified rates, will give control of many annual, biennial, and perennial broadleaf weeds, and many woody brush and vine species commonly found in non-crop land areas. (Refer to Weed List.) Noted (*) perennial weeds may be controlled with lower rates of either DICAMBA DMA SALT 5 or DICAMBA DMA SALT 5 plus tank mix combinations. See Rates and Timings below.

RATES AND TIMINGS

Application rates and timings of DICAMBA DMA SALT 5 are given below. Use the higher level of listed rate ranges when treating dense or tall vegetative growth.

| Weed Stage and Type | Amount of Product Per Acre Fl. oz. | Gals, of Spray Mixture Per Acre** | Spray Concentration for Low Volume Application**** (%vol/vol) |
|--|--------------------------------------|-----------------------------------|---|
| Annual Small, Actively Growing Established weed growth | 6.4 – 12.8 12.8-19.2 | 25-50 50-75 | 3 3 |
| Biennial* - Rosette diameter Less than 3" 3" or more Bolting | 6.4 – 12.8 12.8-25.6 25.6-38.4 | 25 - 50 50 - 100 100 - 150 | 3 - 4 3 - 4 3 - 4 |
| Perennial Suppression or top growth control Noted (*) Perennials Other Perennials | 6.4 – 12.8 25.6-51.2 51.2 | 50 - 100 100 - 200 200 | 4 4 5 |
| Woody Brush and Vines*** Top Growth Stems and Roots | 6.4 – 51.2 51.2 | 50 - 200 200 | 5 5 |

* For best performance, make application when biennial weeds are in the rosette stage.

** Assuming typical application rate of 25.6 fl. oz. of DICAMBA DMA SALT 5/100 gals.

*** Tank mixes may be required for optimal control. Refer to Weed List.

**** Low volume rates must not exceed 51.2 fl. oz. of DICAMBA DMA SALT 5 maximum per acre per year (5% volume/volume = 10 gals, maximum solution per acre per year).

Retreatments may be made as needed; however, do not exceed a total of 51.2 fl. oz (2.0 lbs. a.e.) of DICAMBA DMA SALT 5 per treated acre per year.

Tank Mix Options for Rights-Of-Way, Utility and Industrial Areas, and Fencerows

DICAMBA DMA SALT 5 may be tank mixed with other herbicides for additional weed control. The following table lists example options, but does not limit tank mix options.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from on product, spray drift management from another).

| Herbicide | Rates Per Treated Acre (lbs. a.i.) |
|--------------------|---|
| norflurazon | Consult product labels for rate restrictions. |
| prodiamine | |
| glufosinate | |
| glyphosate | |
| metsulfuron methyl | |
| pendimethalin | |
| triclopyr | |
| clopyralid | |
| bromacil | |
| chlorsulfuron | |

| Herbicide | Rates Per Treated Acre (lbs. a.i.) |
|---------------------|------------------------------------|
| diquat | |
| simazine | |
| diuron | |
| fosamine ammonium | |
| hexazinone | |
| imazapyr | |
| imazemethapyr | |
| sulfometuron methyl | |
| sulfosate | |
| tebuthiuron | |
| 2,4-D | |

Due to the differences that may occur between specific formulated products and specific use ingredients (e.g., water supplies), a compatibility test (see Compatibility Test section) is recommended prior to actual tank mixing.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. This product may not be mixed, loaded, or used within 50 feet of all wells including abandoned wells, drainage wells, and sinkholes.

PESTICIDE STORAGE: Store in original container in a well-ventilated area separately from fertilizer, feed and foodstuffs. Avoid cross-contamination with other pesticides. Spillage or leakage should be contained and absorbed with clay granules, sawdust, or equivalent material for disposal.

PESTICIDE DISPOSAL: Triple rinse pesticide from containers and use rinsate in the pesticide application. Wastes which cannot be used according to label instructions must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

[non-refillable ≤5 gallons]

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10

seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

[non-refillable >5 gallons]

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows (all sizes): Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use for disposal. Insert pressure rinsing nozzle inside of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[Refillable container]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of ALBAUGH, LLC, its Supplemental Distributors, or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

ALBAUGH, LLC, its Supplemental Distributors and the Seller warrant that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW NEITHER ALBAUGH, LLC NOR ITS SUPPLEMENTAL DISTRIBUTORS MAKE ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR ANY PARTICULAR PURPOSE OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND THE EXCLUSIVE LIABILITY OF ALBAUGH, LLC, ITS SUPPLEMENTAL DISTRIBUTORS AND THE SELLER FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. To the extent consistent with applicable law, when Buyer suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), Buyer must promptly notify Seller in writing of any claims to be eligible to receive either remedy stated above. TO THE EXTENT CONSISTENT WITH APPLICASBLE LAW, IN NO CASE SHALL ALBAUGH, LLC, ITS SUPPLEMENTAL DISTRIBUTORS, OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALBAUGH, LLC, its Supplemental Distributors, and the Seller offer this product, and the Buyer accepts it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of ALBAUGH, LLC

No employee or agent of ALBAUGH, LLC, its Supplemental Distributor, or the Seller is authorized to vary or exceed the terms of this Warranty in any other manner.

REGISTERED TRADEMARKS

All trademarks are the property of their respective owners.

041922

LABEL HISTORY
(Not included in final printed label)

| File name | Version mark | Comment |
|-----------------------------|---------------------|----------------------------|
| 045002-000##.20210610.DRAFT | 061021 | Section 3 Application |
| 045002-000GI.20220317.DRAFT | 031722 | Revised as per EPA request |
| 045002-000GI-20220419.DRAFT | 041922 | Revised as per EPA request |
| | | |
| | | |
| | | |
| | | |
| | | |