

US ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (TS-767)
WASHINGTON, DC 20460

EPA REGISTRATION NO.

45017-39

DATE OF ISSUANCE

APR 27 1988

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION
(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

TERM OF ISSUANCE

NAME OF PESTICIDE PRODUCT

Slime-Trol DPP-140

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

WELLS LABORATORIES, INC.
50000 Road
Lewistown, PA 17047

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(B) provided that you:

1. Submit/site all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Add the phrase "EPA Registration No. 45017-39" to your label before you release the product for shipment.
3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Walter C. Francis
Acting Product Manager (32)
Antimicrobial Program Branch
Registration Division (H7505.1)

Enclosures

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

BETZ Slime-tro

SLIME CONTROL AGENT

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

DANGER

CORROSIVE. CAUSES EYE AND SKIN DAMAGE. HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING. AVOID BREATHING VAPORS OR SPRAY MIST. WEAR GOGGLES OR FACE SHIELD AND RUBBER GLOVES WHEN HANDLING. IMMEDIATELY REMOVE AND WASH CONTAMINATED CLOTHING BEFORE REUSE. WASH THOROUGHLY AFTER HANDLING.

ENVIRONMENTAL HAZARDS

THIS PESTICIDE IS TOXIC TO FISH AND WILDLIFE. DO NOT DISCHARGE EFFLUENT CONTAINING THIS ACTIVE INGREDIENT INTO LAKES, STREAMS, PONDS, ESTUARIES, OCEANS, OR PUBLIC WATERS UNLESS THIS PRODUCT IS SPECIFICALLY IDENTIFIED AND ADDRESSED IN AN NPDES PERMIT. DO NOT DISCHARGE EFFLUENT CONTAINING THIS PRODUCT TO SEWER SYSTEMS WITHOUT PREVIOUSLY NOTIFYING THE SEWAGE TREATMENT AUTHORITY. FOR GUIDANCE CONTACT YOUR STATE WATER BOARD OR REGIONAL OFFICE OF THE EPA.

RESTRICTION STATEMENT

THIS PRODUCT IS NOT CLEARED FOR USE IN THE MANUFACTURE OF PAPER AND PAPERBOARD PRODUCTS, COATINGS AND ADHESIVES THAT MAY COME IN CONTACT WITH FOOD.

Active Ingredients:

Bis (trichloromethyl) sulfone.....10%
N-Alkyl (C12-40%, C14-50%, C16-10%)
dimethyl benzyl ammonium chloride.... 8%
Inert Ingredients N:.....82%
Total.....100%
* Inert ingredients include solubilizing and dispersing agents.

CONTENTS: LIQUID
POUNDS PER GALLON: 8.8
EPA REG. NO.: 45017-C0

EPA EST. NO.

DANGER

KEEP OUT OF REACH OF CHILDREN.
In case of contact with skin, wash thoroughly with soap and water. In case of contact with eyes, flush with water for 15 minutes. Immediately contact a physician. In case of ingestion, immediately contact a physician. A Material Safety Data Sheet is available upon request.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Keep container tightly closed. Protect from freezing. Store in a dry place. Do not store in areas where food is stored.
PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide, spray mixture, or rinseate is a violation of Federal law. If these wastes cannot be disposed of by other means, contact the nearest Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest Federal Government facility.
CONTAINER DISPOSAL: METAL AND PLASTIC CONTAINERS: Triple rinse (or equivalent). Then offer for recycling, if available, or dispose of in a sanitary landfill, or incinerate or burn, if allowed by state and local authorities. If burned, state and local authorities must be notified.
FIBER DRUMS WITH LINERS: Completely empty liner by shaking and tapping sides and bottom. Dispose of liner and drum in a sanitary landfill or incinerate if allowed by state and local authorities.

PULP AND PAPER

This product aids in the control of objectionable bacteria and fungi in pulp, paper mill and the additive system, and for the preservation of pulp, pigment slurries, alums, emulsions, adhesives, defoamers, polymers, and paper products. Additions can be made on a continuous or intermittent basis, depending upon the severity of the contamination.

BADLY FOULED SYSTEMS must be cleaned before treatment is begun.

FOR SLIME CONTROL

This product should be added directly to the pulp and paper mill systems. Apply at a point in the system where the product will be uniformly mixed.
INTERMITTENT OR SLUG METHOD- INITIAL DOSE: When the system is noticeably fouled, add this product at a rate of 0.8 to 2.0 lbs per ton of pulp or paper produced. Addition of this product to the additive system should be made directly at the rate of 1.0 to 2.0 pounds (120 to 240ppm) per 1000 gallons. Repeat until control is achieved.
SUBSEQUENT DOSE: When microbial control is evident, add this product at the rate of 0.8 to 1.5 pounds per ton of pulp or paper produced. Treat the system as needed to maintain control. Additions of this product to the additive system may be reduced to 1.0 to 1.5 pounds (120 to 180 ppm) per 1000 gallons.

NET WT. : LBS.
LOT NO.

FOR INDUSTRIAL USE. Technical advice regarding specific site problems is available from BETZ. 1908 8904 PNEEL 1 OF 2

JACKSONVILLE, FL 32256 BUSINESS: 904-733-7110 EMERGENCY (

BETZ Slime-trol DPB-1

SLIME CONTROL AGENT

USE DIRECTIONS CONTINUED FROM PANEL ONE

RECIRCULATING COOLING WATER SYSTEMS

This product aids in the control of bacterial, fungal and algal slimes in evaporative condensers, heat exchange water systems, commercial and industrial cooling towers, influent systems such as flow through filters and lagoons, and industrial water scrubbing systems and brewery pasteurizers.

This product may be added to the system either continuously or intermittently or as needed. The frequency of feeding and duration of the treatment will depend upon the severity of contamination.

POORLY FOULED SYSTEMS must be cleaned before treatment is begun.

FOR CONTROL OF BACTERIA, FUNGI AND ALGAE

INTERMITTENT OR SLUG METHOD

INITIAL DOSE: When the system is noticeably fouled, add this product at the rate of 0.5 to 1.0 pound (60 to 120 ppm) per 1000 gallons of water in the system. Repeat until control is achieved.

SUBSEQUENT DOSE: When control is evident, add this product at the rate of 0.5 to 0.75 pound (60 to 90 ppm) per 1000 gallons of water in the system every 3 days or as needed to maintain control.

CONTINUOUS FEED METHOD

INITIAL DOSE: When the system is noticeably fouled, add this product at the rate of 0.5 to 1.0 pound (60 to 120 ppm) per 1000 gallons of water in the system.

SUBSEQUENT DOSE: Continuously feed this product to maintain a dosage of 0.5 to 0.75 pound (60 to 90 ppm) per 1000 gallons of blowdown (or water loss) from the system.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

APR 27 1966

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Docket No.

45017-39

AUXILIARY WATER AND WASTE WATER SYSTEMS

This product is effective for the control of odor-forming and slime forms in auxiliary water, waste water and waste material disposal, holding or storage tanks, storage piles, associated piping, settling ponds or lagoons, and disposal wells.

This product may be added to the system by slug or intermittent feed or by spray. The frequency of feed or spray and the duration of treatment will depend upon the degree of microbial contamination. Additions to water systems should be made during periods of low flow or close to the pump as possible to insure adequate mixing.

INTERMITTENT OR SLUG METHOD

When treatment is required, add this product at the rate of 0.5 to 1.0 pound (60 to 120 ppm) per 1000 gallons of water already in the system, or being added to the system, for per week or as needed to achieve the desired level of control.

When control is obtained, add this product at the rate of 0.5 to 0.75 pound (60 to 90 ppm) per 1000 gallons of water in the system.

USE IN OIL FIELD SYSTEMS

For the control of slime-forming bacteria, fungi, algae and sulfide-producing water, polymer or micellar floods, water disposal systems or other oil field systems. This product may be added to the systems either continuously or intermittently. The frequency of feeding and duration of the treatment will depend upon the severity of contamination.

POORLY FOULED SYSTEMS should be cleaned before treatment is begun.

INTERMITTENT OR SLUG METHOD

INITIAL DOSE: When the system is noticeably fouled, add this product at the rate of 0.5 to 1.0 pound (60 to 120 ppm) per 1000 barrels of water intermittently for 4 to 8 hours per week or as needed. Repeat until control is achieved.

SUBSEQUENT DOSE: When microbial control is evident, add this product at the rate of 0.5 to 0.75 pound (60 to 90 ppm) per 1000 barrels of water every 3 days or as needed to maintain control.

CONTINUOUS FEED METHOD

INITIAL DOSE: When the system is noticeably fouled, add this product at the rate of 0.5 to 1.0 pound (60 to 120 ppm) per 1000 barrels of water in the system. Continue until control is achieved.

SUBSEQUENT DOSE: Continuously feed this product to maintain a dosage of 0.5 to 0.75 pound (60 to 90 ppm) per 1000 barrels of water in the system.