

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
(AND DOMESTIC ANIMALS)
CAUTION**

Harmful if swallowed. If swallowed, do not induce vomiting. Call a Physician immediately. Avoid contact with eyes or skin. In case of eye contact, immediately flush eyes with fresh water for 15 minutes. If irritation persists, call a Physician. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling. Avoid inhaling of spray mist or vapor.

Wash hands, arms, and face thoroughly with soap and water after handling and before eating or smoking. Routinely shower or bathe after work and wash all clothing with soap and hot water before reusing.

ENVIRONMENTAL HAZARDS


This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from treated areas. Do not apply where run-off is likely to occur. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

1. **PROHIBITIONS:** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.
2. **PESTICIDE DISPOSAL:** Pesticide, spray mixture or residue that cannot be used according to label instructions must be disposed of according to Federal, State, or local procedures under the Resource Conservation and Recovery Act.
3. **CONTAINER DISPOSAL:** Triple rinse (or equivalent) and dispose of in a sanitary landfill or by other approved State and local procedures.
4. **GENERAL:** Consult Federal, State, or Local Disposal Authorities for approved alternative procedures such as limited open burning.
5. **STORAGE:** Store in a cool, dry place.



SUPERIOR OIL

Dormant Spray and Summer Spray

ACTIVE INGREDIENT:	BY WT.
Petroleum Oil*	97.0%
INERT INGREDIENTS:	3.0%
	100.0%

*Petroleum Oil, minimum 92% Unsulfonated residue

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting or give anything by mouth to an unconscious person. Call a Physician immediately.


IF INHALED: Remove victim to fresh air and apply respiration if indicated.

IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly.

IF IN EYES: Rinse eyes for at least 15 minutes with water and call a Physician immediately.

Manufactured by Agricultural Products Company
Ft. Worth, Texas 76112

EPA Est. 43789-TX-01
EPA Reg. No. 43789-43



Net Content

DORMANT SPRAY ONLY

ALTERNATE SIMPLIFIED APPLICATION FOR PLANTS

PLANTS	INSECTS
Peaches, Plums, Apples	San Jose scale
Pears, Hackberry, Dogwood	Spider mite eggs
Pecans	Obscure scale
Oak	(Scale hemispherical)
Birch	Terrapin scale
Chinaberry	White peach scale
Pine	Pine needle scale

DILUTIONS

Mix 10 tablespoons in 1 gal. water or 1 qt. in 6 gal. water. Repeat applications at 2-3 week intervals until control. **NOTE:** Do not spray trees when temperature is below 45 deg. F. Agitate well.

BEST AVAILABLE COPY

DIRECTIONS

Dilute with water, as per directions below. Spray plants thoroughly, making sure underside of leaves are completely covered. For oil type applications, spray tank must be free of any sulphur residue. Do not combine with sulphur sprays and do not precede or follow a sulphur product application with this product until sufficient time has elapsed for safety (at least 14 days; in some cases longer). If in doubt, spray a portion of one tree previously treated with sulphur using 1 quart to 25 gallons of water. If sufficient time has not elapsed, burn will show in several days. Do not use this product when temperatures are above 90 degrees F. When temperatures drop below 32 degrees F (nighttime low), do not spray until daytime temperatures have warmed to a point above 35 degrees F, and plants or trees are free from frosted frost or heavy dew. Do not use on foliage closer than 60 days after or 90 days before using Captain or Phaltan products. Do not use in combination with Sevin on crops other than citrus. Application to trees weakened by disease, drought, driving winds, or high nitrogen applications may result in oil injury. When using this product in combination with other pesticides, observe all labeling precautions and limitations for the products involved. Do not use in combination with other treemakers unless previous experience assures compatibility.

MIXING

With agitator in motion add water, agitator shaft is just covered. When combining wettable powders (insecticides or fungicides) with this product as directed on this label add these first. Then, with agitator still running, add this product and finish filling with water. Continue agitation to insure good emulsification.

Do not use if this spray oil does not emulsify. Correct emulsification is indicated if, when this product is added to water in the tank, a uniform milky white spray mixture results during agitation. If the spray mixture, after adding this product under agitation, is not milky-white do not use the resulting spray or the remainder of this product. To test for correct emulsification, add a small amount of this product to water in a mixer before wettable powders or other insecticides or fungicides as directed on this label are added.

AMOUNTS GIVEN FOR 100 GALLONS OF DILUTED SPRAY

ALFALFA (SEED CROP ONLY). Use 1 quart per acre in 20 to 30 gallons water by ground applicator or 10 to 15 gallons water by air when used in combination and at the recommended rates with either CYTHION® or Toxaphene for control of pests listed on the product labels. For Spider Mites, use 1 quart per acre at the dilution noted above with Kelthane® at the recommended label rates.

ALMONDS, APRICOTS, CHERRIES, NECTARINES, PEACHES, PLUMS, PRUNES. Dormant or delayed dormant. Use 1½ to 2 gals. for brown apricot scale, black scale, San Jose scale, olive scale (parlatona scale), oleander scale, frosted scale, Italian pear scale, European fruit lecanium, putnam scale, parlatona, soft brown scale, eggs of aphids, red spider mites. For Summer Use, use 3 to 4 quarts for control of red spider mites, 1 gallon for control of summer stages of brown apricot scale.

APPLES: Use 1½ to 2 gallons delayed dormant for control of rust mite, spider mite eggs.

APPLES, PEARS. Dormant or delayed dormant. Use 1½ to 2 gallons for San Jose scale, Italian pear scale, eggs of aphids, red spider mites. For summer use, apply 3 to 4 qts. for red spider mites, and 1 to 1½ gal. for summer forms of San Jose and Italian pear scale.

FIGS. Dormant or Delayed Dormant. Use 3 gals. for control of Fig Scale.

GRAPES. Dormant spray. Use 1 gal. plus recommended amount of Ethyl Parathion. Use 250 to 400 gallons spray per acre, depending upon size of vines, for control of mealy bugs.

LEMONS, GRAPEFRUIT. Use 1 to 1½ gals. plus recommended rate of Cythion®. For best results spray during winter and spring months for control of red scale, yellow scale, purple scale, black scale, and red spider mites. Do not use within 8 weeks of harvest on lemons.

OLIVES. Use 1½ gals. or combine 1 gal. with the recommended rate of Ethyl Parathion per 100 gals. water. Apply in pre-bloom or post-bloom up to buckshot, for control of parlatona scale, spider mites, oleander scale, black scale.

NAVEL ORANGES. Use 3 qts. plus recommended rate of Cythion® for control of red scale, black scale, yellow scale, purple scale, red spider mites. Do not apply later than September 15.

VALENCIA ORANGES. Use 3 qts. plus recommended rate of Cythion® for control of red scale, black scale, yellow scale, purple scale, red spider mites. Do not apply later than September 15.

PEARS. Delayed Dormant. Use 2 gals. for pear psylla. For Summer Use, apply 1 to 1½ gals. for blister mite.

ORNAMENTALS AND SHADE TREES. Conifers (except those conifers and ferns not tolerant to oil sprays), Holly, Pyracantha, Sycamore, Hawthorne, Flowering Peach, Flowering Cherry, Apple, Magnolia, Maple, Dogwood, Oak, Privet, Birch. Use 1 gal. for soft brown scale, oyster shell scale, brown apricot scale, black scale, San Jose scale, red spider mites and eggs. For Dormant Spray on Delicous Varieties use 2 gals. Dormant spray on other varieties use 1 gal. NOTE: May cause discoloration of blue spruce.

WALNUTS. Late spring to mid-summer: only. Use 2 to 4 quarts for control of summer stages of frosted scale, European fruit scale, San Jose scale, and other softshelled scales, red spider mites. Do not apply after husk split.

*Cythion is a Reg. TM of American Cyanamid Co.

**Kelthane is a Reg. TM of Rohm and Haas Co.

NOTICE - READ CAREFULLY

CONDITIONS OF SALE: Seller(s) offer(s) this product for sale subject to and buyer and all users are deemed to have accepted the following conditions of sale and warranty which may only be varied by written agreement of a duly authorized representative of Seller(s).

WARRANTY LIMITATION: Seller(s) warrant(s) that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use on the label subject to the inherent risks referred to below. Seller(s) make(s) no other express warranties.

THERE IS NO IMPLIED WARRANTY OF MERCHANTABILITY and there are no warranties which extend beyond the description on the label hereof.

INHERENT RISKS: The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with its use. Buyer assumes all risks associated with use or application of the product contrary to label instructions or resulting from: (a) extraordinary weather conditions.

LIMITATION OF LIABILITY: In no case shall Seller(s) be liable for special, indirect, or consequential damages resulting from the use or handling of this product and no claim of any kind shall be greater in amount than the purchase price of the product in respect of which such damages are claimed.

HOME AND GARDEN USE

SPRING & SUMMER SPRAY

PLANTS	INSECTS
Camellia	Tea scale
Azalea	Azalea scale, mealy bugs, red spider
Euonymus	Euonymus & red scale, mealy bugs
Rose	Rose scale, red spider
Juniper	Juniper scale, red spider
Privet	White, black scale, red spider
Magnolia	Magnolia scale, red spider
Citrus	Scale, mealy bugs, white flies, mites
Boxwood	Scale, mealy bugs, red spider
Holly	Tea scale
Sage Plum	Scale

DILUTIONS

Mix 5 tablespoons in 1 gal. water or 1 qt. in 12½ gals. water. Repeat at 2 or 3 week intervals until control.

NOTE: Do not spray plants when temperature is above 90 deg. F.

NOTE: Do not combine with any other spray when spraying citrus. See detailed instructions on this label for use of this product before or after sulphur products, Captain, or Phaltan.

BEST AVAILABLE COPY