# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

# WARNING

Causes eye irritation. May cause skin irritation. Harmful it swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist. Do not get in eyes, on skin, or on clothing. Wash with soap and water after handling and before eating or smoking. Wash contaminated clothing before reuse. Do ript use or store near food or feed.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically Atropine is antidotal

#### **FNVIRONMENTAL HAZARDS**

This product is toxic to fish, birds, and other wildlife. Do not apply directly to takes streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this tabel.

This product is highly toxic to bees exposed to direct treatment or risidues on plants. Protective information may be obtained from your Agricultural Extension Service.

#### PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

### **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

#### STORAGE AND DISPOSAL

- PROHIBITIONS: Do not contaminate water, food, or feed by storage or oisposal. Open dumping is prohibiter
- 2. CONTAILIER DISPOSAL: Do not reuse empty container. Wild ap container and put in trash collection
- 3. STORACE: Store in a cool, dry place



## LAWN AND ORNAMENTAL GARDEN SPRAY

Controls pests on Turf: Chinch Bugs, Sod Webworms, Ants. Brown Dog Ticks, Crickets, Cutworms, Earwigs, Grasshoppers, Hyperodes Weevils. Controls pests on Ornamental Plants. Aphids, Bagworms. Cutworms, Grasshoppers, Leafhoppers, Mealybugs, Mites, Spittlebugs. Exposed Thrips, Whiteflies

#### **ACTIVE INGREDIENTS:**

Chlorpyrifos [O, O-diethyl O-(3,5,6-trichloro-2-pyridyl)	
phosphorothicate]*	2.0%
Xylene	89.7%
INERT INGREDIENTS:	8.3%
	100 00/

\*Dursban Reg. Trademark Dow Chemical Company Contains 2 5 ounces Chlorpyrifos per gallon

# KEEP OUT OF REACH OF CHILDREN WARNING

#### STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. Do not induce vomiting. Vomiting may cause appration pneumonia.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth

IF ON SKIN: Wash the affected area with soap and warm water IF IN EYES: Flush eyes with plenty of water. Get immediate medical attention

See side panel for additional precautionary statements

# A. C. C. E. P. T. E. D. 789-TX-01 EPA Reg. No. 43789-63

2 1981

Under the Fether 1.1 in the Fungicide, and 1 ferrior Act, as amended for the pessicide registered under EPA Reg. No.

OCT.

Net Contents

AGRICULTURAL PRODUCTS COMPANY Ft. Worth, TX 76112

#### DIRECTIONS

For control of chinchbugs, sod webworms (lawn moths), and, brown dog ticks, crickets earwigs, and grasshoppers infesting lawns and turfs. Thoroughly water immediately after treatment to wash the insecticide into the turf. However, for sod webworm control delay watering or mowing the treated area for 12 to 24 hours after treatment. Dilute spray at the rage of 2 tablespoons per gallon of water to freat 50 square feet or 1 gallon per 125 to 150 gallons of water and apply over 6,600 square foot area. When using a lawn hose-on sprayer (15-gallon to 20-gallon sprayer), fill quart jar to 20-ounce mark with spray and finish filting jar to top with water. Then apply this mixture over 1,000 square feet of lawn grass. Ten steps in one direction and elever in another direction is approximately 1,000 square feet. Repeat application if chinch bug damage reoccurs or when pests reappear. Keep children and pets off of treated area until spray has dried. Keep out of lawn and garden fishpools and other bodies of water.

**AROUND BUILDINGS:** 

To help prevent infestation of buildings by ants, crickets, earwigs, and sowbugs (pillbugs) treat a band of soil six to ten feet wide around and adjacent to the building and also the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. Difute spray at the same rate as above. Apply as a coarse (large droplet) spray to thoroughly and uniformly wet the band area.

To control hyperodes (turfgrass) weevils use 2 gallons spray per 125 to 150 gallons of water and apply over 6.600 square foot area or 4 tablespoons per gallon of water to treat 50 square feet. Spray suspected problem creas in mid-April and again in mid-May. Consult your local Agricultural Extension Service Specialist for best time to treat in your area. NOTE:

Do not spray vegetable gardens or other plants used for food purposes PEACH TREE BORER:

Dilute this product in water it the rate of 25 oz of this product per gallon of water. A ply this spray mixture to flowering trees and shrubs of the Genus Prunus as a trunk spray before newly hatched larvae enter the trees. Apply as a coarse, low-pressure spray. Thoroughly wet all bark areas from ground level to scalfold limbs. Consult your Agriculturat Experiment Station or Extension Service specialist for proper time to treat in your area.

ASH BORER, LILAC BORER, DOGWOOD BORER:

Dilute this product in water at the rate of 7° oz of this product per gallon of water. Apply this spray mixture to trunks and lower limbs of trees when the adults begin to emerge. Consult your State Agricultural Experiment Station or Extension Service specialist for proper time to treat in your area. Also controls scale crawlers. Oak lecanium. Pine needle Toa crawlers, and White peach crawlers.

IMPORTED FIRE ANT:

For the treatment of potted containerized balled or birriapped nursery stock, use 8.9 oz of this product diluted in 10 gallons of water. For control of lire ants in soci attached to the roots of these plants, completely submerge the ball of soil in a tank containing the mixture. Do not remove burlap wrap or plastic containers with drain holes prior to submerging. Keep soil submerged until complete safuration has occurred, normally about 30 seconds. Wear rubber gloves, boots, and apron during this operation.

ORNAMENTALS:

To control aphids bagworms, grasshoppers, miles spittlebugs, whiteflies Dilute 2 ounces (4 tabli-spoons) in 1 gallon of water Avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals if needed. May be used on Shrubs. Shade and Flowering Trees, and liowers except as noted below. NOTE: Do not use, on azaleas, camellias, poinsettias, roses, or variegated by because of possible injury to these plants.

#### NOTICE — READ CAREFULLY

CONDITIONS OF SALE. SePerial offer(s) this product for sale subject to land buyer and all users are deemed to have accepted the following conditions of sale and warranty which may only be varied by written agreement of a duty authorized representative of Seller(s).

WARRANTY LIMITATION. Sefer(s) warrants) mat this product comprise to the chemical description on the label and is reasonably in for the purposes referred to in the exerctions for use on the label subject to the inherent naks referred to below. Sefer(s) make(s) no other express warranties. THERE IS NO IMPLIED WARRANTY OF MERCHANTABILITY and there are no warrantes which extend beyond the description on the label hereof IMHERENT RISKS. The directions to luse of this product are believed to be reliable and should be followed carefulty. However, it is impossible to eliminate all risks associated with its use. Buyer assumes all risks as sociated with use or application of this product contrary to label instructions or resulting from extraordinary weal their conditions.

LIMITATION OF LIABILITY: In no case shall Select(s) be hable for special indirect, indirect, in constituential damages resulting from the use or handling of this product and no claim of any kind shall be creater in amount than the purchase price of the product in respect of which such damages are claimed.

· Jan 1980 198

BEST AVAILABLE COP