

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING: Corrosive. Causes eye damage. Harmful or fatal if swallowed. Do not get in eyes or clothing. Wear goggles or face shield when handling. Avoid contact with skin. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Get Medical Attention.

IF IN EYES: Flush with plenty of water. Get Medical Attention.

IF ON SKIN: Wash with plenty of soap and water. Get Medical Attention if irritation persists.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

GENERAL INFORMATION:

Do not allow spray to contact adjacent crops or other plants that may be defoliated. Clean and wash spray equipment before and after use and do not allow wash water to contact desirable plant. To avoid fire or explosion do not mix with insecticides. Do not contaminate water by cleaning of equipment or disposal of wastes.

FARMWORKER SAFETY

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure.) When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information. **WARNING**—Area treated with Sodium Chlorate on (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure refer to the Statement of Practical Treatment on the Knock'um Off label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOTE: Do not allow spray to contact adjacent crops or other plants that may be defoliated. Clean and wash spray equipment before and after use and do not allow wash water to contact desirable plants. To avoid fire or explosion do not mix with insecticides. Do not contaminate water by cleaning of equipment or disposal of wastes.

COTTON DEFOLIATION: Plants must be mature to get results with a defoliant. The usual signs of maturity include a slight yellowing of the foliage and a reddish tinge on the leaves with a clean break by a slight downward pull. Plants should not be drought-stricken because the tough wilted leaves do not permit uptake of the chemical, or growth of an abscission layer. When defoliants are applied to drought-stricken plants the leaves will die but usually not drop, thereby causing extra trash to be picked up in harvesting.

Weather conditions that favor good defoliation are high temperatures, high humidity, low wind velocity and high to adequate soil moisture. Plant conditions which favor good defoliation are ample fertility and moisture, complete insect control, heavy boll load and bolls mature, vegetable growth ceased, and little or no second growth.

Apply 1/2 to 3/4 gallons of KNOCK'UM OFF per acre in sufficient water to give thorough coverage. When using ground equipment apply 15 to 20 gallons of spray solution per acre, and by air, 5 gallons of spray solution per acre. Thorough coverage is essential.

Apply KNOCK'UM OFF to mature cotton plants after the youngest bolls expected to make cotton are at least 30 days old. Do not apply later than 7 days before harvest.

Do not graze treated areas or feed gin waste to livestock,

GRAIN SORGHUM - KNOCK'UM OFF may be used to reduce the moisture content in grain sorghum prior to harvest.

KNOCK'UM OFF may be applied by aircraft or ground equipment. Thorough coverage of the grain sorghum is essential.

To reduce the moisture content of grain while the crop is standing apply 3/4 to 1 gallon KNOCK'UM OFF per acre in sufficient water to give thorough coverage. Apply a total of 5 to 10 gallons of spray per acre by airplanes and 10 to 20 gallons per acre by ground equipment. Use the lower rate when grain moisture is low and the weather is clear and dry. Use the higher rate when conditions for desiccation are poor.

SUNFLOWER - To facilitate harvest and reduce moisture in the fully mature seed heads apply 1 gallon of KNOCK'UM OFF in sufficient water to give good coverage, 7 or more days before normal harvest when sunflower heads are fully mature. Apply in 5 to 10 gallons of water per acre by air or in 20 to 30 gallons of water per acre by ground with a high clearance sprayer. Do not apply later than 7 days before harvest. Do not graze treated fields or feed treated forage.

SAFFLOWERS - To facilitate harvest and reduce moisture in the fully mature seed heads apply 1 gallon of KNOCK'UM OFF in sufficient water to give good coverage, 7 or more days before normal harvest when safflower heads are fully mature. Apply in 5 to 10 gallons of water per acre by air or in 20 to 30 gallons of water per acre by ground. Do not apply later than 7 days before harvest. Do not graze treated fields or feed treated forage.

RICE: To facilitate harvest and reducing moisture in the fully mature seed heads and to desiccate immature red rice in mature white rice, apply by aircraft 3/4 to 1 gallon of KNOCK'UM OFF per acre in 5 or more gallons of water per acre. Use the lower rate when crop conditions and weather are right to enhance crop conditions and weather are right to enhance rapid drying. Apply 7 to 10 days before anticipated harvest. Thorough coverage of the crop is essential to achieve good results. Do not graze treated fields or feed treated straw or crop residues within 14 days of application.

SOYBEANS - As a soybean harvest aid to dry weeds in early maturing soybeans and to facilitate the harvest, apply 1 gallon of KNOCK'UM OFF per acre in enough water to give thorough coverage; application may be made by ground equipment or aircraft. Make application 7 to 10 days before anticipated harvesting date when beans are mature and ready for harvest. For air application use a minimum of 5 gallons ready-to-apply spray solution per acre; for ground equipment apply a minimum of 20 gallons of ready-to-apply spray solution per acre. Do not graze treated fields or feed treated soybean foliage.

CHILIPEPPERS - Apply 1/2 to 1 1/2 gallons per acre in 5 to 15 gallons of water by aircraft and 20 to 40 gallons of water by ground rig. To be used on peppers for processing only. Apply 10 days before harvest. When grown for processing, consult a processor before applying.

GUAR BEANS - To facilitate harvest and reduce bean seed moisture, apply 1 gallon of KNOCK'UM OFF in enough water to provide thorough coverage either by aircraft or ground equipment. Make application 7 - 10 days before anticipated harvest date. Do not apply under conditions of extreme heat during the middle of the day. For air application, use 5 - 10 gallons of total solution per acre; for ground application, use a minimum of 20 gallons of total solution per acre. Do not graze treated fields or feed treated fodder or forage to livestock.

FLAX - For Use As A Harvest Aid: To desiccate trashy weeds and desiccate the mature crop, apply 1 gallon Knock'um Off per acre in enough water to give thorough coverage. Make application on bright, sunny days (preferably when temperature is above 60°F) 7 to 10 days before anticipated harvest date. For air application, use minimum of 5 gallons ready-to-apply material per acre. Activity of the spray will be enhanced by the addition of a good non-ionic surfactant at the rate of 1 quart per 100 gallons of spray.

DO NOT graze treated fields or feed treated flax foliage and fodder for 14 days after application.

CORN - Knock'um Off may be used as a harvest aid for desiccation of trashy weeds in early maturing corn. Use 1 gallon of Knock'um Off in 5 to 7 gallons of water per acre by air. Apply at least 14 days before anticipated harvest date on clear days when temperatures are expected to go above 70°F. Use the 5 gallon rate when weeds are small and the crop canopy is fairly open.

DO NOT graze treated fields or feed fodder, forage or residual seeds within 14 days of application.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE INSTRUCTIONS: Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool dry area away from any heat or ignition source. Do not stack plastic containers (4 x 1 gal. or 2 x 2.5 gal.) over 2 pallets high. Do not stack metal containers over 4 pallets high. Move containers by handles or cases. Do not move containers from one area to another unless they are securely sealed. Keep container tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed, and fertilizer to avoid contamination. Store in original containers only. Unloading storage and work areas must be as clean as possible to prevent contamination if spilled. If concentrate or spray solution is allowed to dry on flammable objects, the chlorate component can act as an oxidizer. If the contents are leaking or material is spilled follow these steps:

1. Contain spill, absorb with a material such as saw dust, clay granules, or dirt.
2. Collect and place in suitable containers for disposal.
3. Wash area with water and soap to remove remaining pesticide.
4. Follow washing with clean water rinse.
5. Clothing contaminated with chlorate solution must be removed at once before solution dries. Dried contaminated clothes can be ignited easily. Wash contaminated clothing immediately.
6. Do not allow run off to enter sewer or contaminate water supplies.
7. Dispose of waste as indicated below:

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Metal - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY - CONDITION OF SALE:

Directions for Use of this product are based on field use and tests believed reliable and should be followed carefully. If it is, however, impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of Ida, Inc. or the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result.

The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such of the results to be obtained if not used in accordance with directions or established safe practice. The buyer assumes all responsibility, including injury or damage resulting from its misuse as such or in combination with other materials.