



8

A Flowable 8 Pounds Elemental Sulfur

ACTIVE INGREDIENT:	
Elemental Sulfur	65.00%
INERT INGREDIENTS: <u>35.00%</u>	
TOTAL 100.00%	

Product contains 8.0 lbs. of sulfur per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Statement of Practical Treatment
 IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
 IF SPLASHED IN EYES: Immediately flush eyes with plenty of water. Get medical attention if irritation develops.
 IF SPILLED ON SKIN: Immediately remove contaminated clothing, INCLUDING SHOES, and wash skin with soap and plenty of water. If irritation develops, send for a physician.
 See back Panel for Additional Precautionary Statements.

EPA Reg. No. 45115-
EPA Est. No.

Ida, Inc. EPA Reg. No. 45115-
EPA Est. No.

Net Contents: Gallons Liters

PRECAUTIONARY STATEMENTS

Hazardous to humans and domestic animals.

CAUTION: Harmful if swallowed, avoid breathing spray mist. May cause irritation of nose, throat, and skin. Avoid contact with eyes, skin and clothing. Causes moderate eye and skin irritation. Do not contaminate feed and food stuffs. Wash thoroughly with soap and water after handling.
Environmental Hazards

Do not apply directly to water. Do not contaminate water by disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proven that sulfur is safe in that locality.

During periods of high temperatures sulfur may burn foliage and fruit. Do not make sulfur application at excessively high temperatures. Do not use Sulfur 8 with oil or within four (4) weeks of an oil application.

When Sulfur 8 is used with arsenicals, lime should be added to prevent plant injury.

Use of this material in a manner other than in accordance with this label may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

Consult your State Agricultural Experiment Station or State Extension Service Specialist for complete details on the spraying program best suited to your local conditions.

Unless otherwise specified for specific tree fruits, dosage rates are given as pints of Sulfur 8 per 100 gallons of water for use in a full coverage spray.

Shake well before using.

Pour recommended amount into partially filled spray tank. Keep agitator running during filling and spraying operation. Do not allow mixture to stand. Failure to maintain agitation will cause Sulfur 8 to settle and may necessitate manual stirring to redisperse.

APPLE, PEAR

Scab: Prebloom through calyx sprays $2\frac{1}{2}$ to $5\frac{1}{2}$ cups. Cover sprays $\frac{3}{4}$ to $1\frac{1}{2}$ pints. Powdery mildew: Prebloom through calyx sprays $2\frac{1}{2}$ to $5\frac{1}{2}$ cups. Cover sprays $1\frac{1}{2}$ pints.

PEACH

Brown rot, powdery mildew, Scab: Pink and bloom sprays $2\frac{1}{2}$ to $5\frac{1}{2}$ cups. Petal fall, shuck and cover sprays $\frac{3}{4}$ to $1\frac{1}{3}$ pints.

CHERRY, PLUM, PRUNE

Brown rot, leaf spot: Pink and bloom sprays $2\frac{1}{2}$ to $5\frac{1}{2}$ cups. Petal, shuck and cover sprays $\frac{1}{2}$ to $1\frac{1}{3}$ pints.

CITRUS

(Grapefruit, lemons, limes, oranges, tangelos and tangerines).
Rust Mite: $\frac{3}{4}$ to 3 pints. Do not apply within 4 weeks of an oil application.

PEANUTS

Rust: Apply $1\frac{1}{2}$ to 3 pints per acre in sufficient water for thorough coverage. Begin application approximately 6 weeks after planting. Repeat at 7 to 10 day intervals as long as rust control is required. Do not apply within 4 weeks of an oil application.

COTTON

Atlantic Mite: $1\frac{1}{2}$ pints for early season control and up to 6 pints for mid to late season control.

GRAPES

(California only) - Powdery mildew: $\frac{3}{4}$ to $1\frac{1}{2}$ pints per 100 gallons water, maximum of 6 pints Sulfur 8 per acre in spring and summer. Make first application when shoots are 6 to 8 inches long and third at or about bloom stage.

The timing rate and number of applications will vary with local conditions. Consult State Agricultural Service representative or State Agricultural Experiment Station in your region for specific information. Follow use PRECAUTIONS given on this label.

POTATOES

Powdery mildew: Use $2\frac{1}{2}$ to 3 pints in 8 gallons water per acre by air. Apply at first sign of infection and repeat at 3 week intervals.

STRAWBERRIES

Powdery mildew: Use 3/4 pint. Apply at first sign of infection and repeat at 3 week intervals. For strawberries that will be canned, discontinue application of sulfur well before harvest in accordance with local canner's recommendation.

SUGAR BEETS

Powdery mildew: Use 3/4 to 2 1/4 pints in 5 to 8 gallons water per acre by air. Apply at first sign on infection and repeat at 3 week intervals.

TOMATOES

Tomato Russet Mite: Use 3 pints to 6 pints per acre depending upon size of plants. Use full coverage spray.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment.

PESTICIDE DISPOSAL:

Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

CONTAINER DISPOSAL:

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or disposal of in a sanitary landfill, or by incineration if allowed by state and local authorities.

WARRANTY - CONDITION OF SALE:

DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result.

ALL SUCH RISKS ARE ASSUMED BY THE BUYER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use. Seller makes no other warranties, express or implied, including FITNESS OR MERCHANTABILITY. In no case shall the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.