

17-5  
4/21/90

Les Shockey  
Labeling Specialist  
IDA, Inc.  
2215 West Street, 2nd Floor  
Germantown, TN 38138

JUN - 4 1990

Dear Ms. Shockey:


Subject: IDA, Inc. Atrazine 5F  
EPA Reg. No. 45115-69  
Re: Atrazine Voluntary Exposure Reductions Measures  
Your Amendment Resubmission Dated May 24, 1990

The labeling referred to above, submitted in connection with the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable with the following comment:

1. On page 4, it appears that the heading, "GENERAL INFORMATION" and the 1st paragraph under that heading are missing. Please restore and be sure that there are no references to pineapples or other unsupported crop uses.

A stamped approved copy of your label is attached. Please submit five (5) copies your final printed label incorporating this amendment and the above mentioned comment before you release the product for shipment. All technical and end-use products released for shipment must have amended labels after September 1, 1990.

Sincerely yours,



Robert J. Taylor  
Product Manager (25)  
Fungicide-Herbicide Branch  
Registration Division (H7505C)

CONCURRENCES

SYMBOL	H7505C						
SURNAME	ma						
DATE	4/4/90						

# RESTRICTED USE PESTICIDE

(GROUND WATER CONCERNS)

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

THIS PRODUCT IS A RESTRICTED USE HERBICIDE DUE TO GROUND WATER CONCERNS. USERS MUST READ AND FOLLOW ALL PRECAUTIONARY STATEMENTS AND INSTRUCTIONS FOR USE IN ORDER TO MINIMIZE POTENTIAL FOR ATRAZINE TO REACH GROUND WATER.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

IDA, INC.

45115-69

## ATRAZINE 5 FLOWABLE

For season-long weed control in corn and sorghum, for weed control in certain other crops in non-crop and industrial sites.

ATRAZINE 5F CONTAINS 5 LBS. ACTIVE INGREDIENTS PER GALLON.

### ACTIVE INGREDIENTS:

Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine.....	50.81%
Related Compounds.....	2.67%
INERT INGREDIENTS:.....	46.52%
TOTAL:.....	100.00%

Keep Out Of Reach Of Children

### CAUTION

SEE SIDE PANEL FOR STATEMENT OF PRACTICAL TREATMENT.

SHAKE WELL BEFORE USING

EPA REGISTRATION NUMBER: 45115-69

EPA ESTABLISHMENT NUMBER:

NET CONTENTS: 2.5 Gallons / 9.45 Liters

MANUFACTURED FOR: IDA, INC.

2215 West Street, Suite 2  
Germantown, TN 38138

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful or fatal if swallowed. Causes moderate eye irritation. Do not get in eyes, on skin, or on clothing. Wear goggles. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating. Remove contaminated clothing and wash before reuse. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Users are required to wear long sleeve shirts and long pants or equivalent, chemical resistant gloves, and boots (waterproofed). In addition, persons involved in mixing/loading operations are required to use chemical resistant rubber or neoprene gloves and face shield or goggles.

#### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Call a Physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF IN EYES:** Immediately flush eyes with plenty of water. Get medical attention if irritation develops.

**IF ON SKIN:** Immediately flush skin with plenty of soap and water while removing contaminated clothing and shoes. If irritation develops, seek medical attention.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

Atrazine can travel (seep and leach) through soil and enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

#### FARMWORKER SAFETY STATEMENT

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must inform workers of areas or fields that may not be entered without specific protective clothing until sprays have dried, and appropriate actions to take in case of accidental exposure, as described under Precautionary Statements on this label. When oral warnings are given, warnings shall be given in a language customarily understood by the workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: "CAUTION: Area treated with atrazine on (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure, refer to Statement of Practical Treatment on the ATRAZINE 5F label."

P27404

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Postemergence application to corn and sorghum must be made before corn and sorghum reaches 12 inches in height.

The maximum application rate for corn and sorghum is 3 lbs. a.i. per acre per calendar year. Applications for quackgrass suppression in corn and sorghum are restricted to a spring application only. No fall applications are permitted.

Application rates to noncrop land for industrial weed control cannot exceed a maximum of 10 lbs. active ingredient per acre per calendar year.

## NOTE :

Ground water contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material.

This product may not be mixed/loaded, or used within 50 feet of all wells including abandoned wells, drainage wells, and sink holes.

Do not apply this product through any type of irrigation system.

sequence with other registered herbicides which are not triazines. If any resistant biotypes are expected to be present, use a registered non-triazine herbicide. Consult with your state Agricultural Extension Service for specific recommendations.

Since ATRAZINE SF acts mainly through root absorption, its effectiveness depends on rainfall or irrigation to move it into the root zone. Broad weeds develop a shallow cultivation or rotary hoeing will generally result in better weed control.

ATRAZINE SF is nonflammable.

Care should be taken to avoid using ATRAZINE SF near adjacent desirable plants or in greenhouses or injury may occur.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.

**NOTE:** The Sener does not recommend the use of ATRAZINE SF in combination with other herbicides or oils except as specifically described on the label or in literature published by the Sener. FAILURE TO FOLLOW ALL PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

#### APPLICATION PROCEDURES

**Ground Application:** Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. Be certain that nozzles are uniformly spaced and are the same size. Calibrate sprayer before use and recalibrate at the start of each season and when changing carriers. Unless otherwise specified, use a minimum of 10 gal. of spray mixture/A for all preplant incorporated, preplant surface, preemergence, and postemergence applications (with or without oil or surfactant) with ground equipment.

Use a pump with capacity to (1) maintain 35-40 psi at nozzles, (2) provide sufficient agitation in tank to keep mixture in suspension, and (3) to provide a minimum of 20% bypass at all times. Use centrifugal pumps which provide propeller shear action for dispersing and mixing the product. The pump should provide a minimum of 10 gal./min./100 gal. tank size circulated through a correctly positioned sparger tube or jets. Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump should be a 15-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Check netze manufacturer's recommendations.

For band application, calculate the amount to be applied per acre as follows:

$$\frac{\text{Band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of band}$$

#### Aerial Application

Use aerial application only where broadcast applications are specified. Apply a minimum of 1 qt. of water for each quart of ATRAZINE SF applied per acre. For postemergence treatments on corn and sorghum, apply recommended rate in a minimum of 2 gal. of water/A. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin, and should wash thoroughly before eating and at the end of each day's operation.

#### ATRAZINE SF IN WATER APPLICATIONS

ATRAZINE SF liquid formulation, should be mixed with water and applied as a spray. Pour ATRAZINE SF into the tank during or after filling. Hydraulic jet or mechanical agitation is recommended during mixing and application to keep the materials in suspension. All return lines to the tank must discharge below liquid level and agitation should not be so violent as to cause air bubbles to form in the liquid. Wash sprayer thoroughly after use.

#### ATRAZINE SF IN LIQUID FERTILIZER APPLICATIONS

Nitrogen solutions or complete liquid fertilizers may replace all or part of the water as a carrier for preemergence and preplant applications of ATRAZINE SF on corn and sorghum. Mixing should be accomplished as described under ATRAZINE SF IN WATER APPLICATIONS. Check the compatibility of this product with liquid fertilizer and/or nitrogen solution as shown below before use. Do not apply in nitrogen solution or complete liquid fertilizer after corn or sorghum emerges, except as noted under Crop Injury by treatment for corn or crop injury may occur.

**Compatibility Test:** Since liquid fertilizers can vary, even within the same analysis, always check compatibility with herbicides at each time before use. Be especially careful when using complete fertilizer or fluid fertilizers as serious compatibility problems are more likely to occur. Commercial application of fertilizers may improve compatibility in some instances. The following test assumes a spray volume of 25 gal./A. For other spray volumes, make appropriate changes in the ingredients. Check compatibility using this procedure:

1. Add 1 pt. of fertilizer to each of 2 one-quart jars with tight lids.
2. To one of the jars add 1/2 tsp. or 1.2 ml. of a compatibility agent approved for this use such as Compac® or Unifac®. 1/2 tsp. is equivalent to 2 pt. per 100 gal. spray. Shake or stir gently to mix.
3. To both jars add the appropriate amount of herbicide(s). If more than one herbicide is used, add them separately with dry herbicides first, flowables next, and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix. The appropriate amount of herbicides for this test follows:  
**DRY HERBICIDES:** For each pound to be applied per acre, add 1/2 level tsp. to each jar.

**LIQUID HERBICIDES:** For each pint to be applied per acre, add 0.5 tsp. or 2.5 ml. to each jar.

4. After adding all ingredients, put lids on and tighten, and invert each jar 10 times in row. Let the mixtures stand 15 min. and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the two jars. If either mixture separates but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (A) stir the dry herbicide(s) in water before addition, or (B) add 1/2 of the compatibility agent to the fertilizer and the other 1/2 of the emulsifiable concentrate or flowable herbicide before addition to the mixture. If incompatibility is still observed, do not use the mixture.

**Note:** In the event of a compatibility problem when mixing oil with this product and water, a compatibility agent such as Compac® or Unifac® should be used. Any of the above oils contaminated with water or other materials can cause compatibility problems and/or crop injury.

#### ATRAZINE SF PLUS EMULSIFIABLE OIL OR OIL CONCENTRATE

Adding emulsifiable oil (petroleum-derived, petroleum-derived oil concentrate, or single or mixed crop-derived concentrate) to postemergence water-based sprays in corn and sorghum may improve weed control. However, under certain conditions, the use of either type of oil may seriously injure the crop. To minimize this possibility, observe the following directions.

Use one of the following properly emulsified:

1. A suitable oil concentrate containing at least 1% but not more than 20% suitable emulsifier or surfactant blend.
2. Petroleum-derived oil containing at least 1% suitable emulsifier.

#### MIXING PROCEDURES

All uses: (1) Be sure sprayer is clean and not contaminated with any other materials or crop residue. Sprayer clogging may result. (2) Fill tank with clean water, nitrogen solution, or complete fertilizer. (3) Start agitation. (4) Be certain that the agitation system is working properly. (5) Add herbicide(s) or other materials to the liquid surface. (6) Pour product directly from container into the tank. (7) Increase agitation if necessary to maintain surface action. (8) Add emulsifiable concentrate to tank mix herbicide(s) after this product is thoroughly suspended. (9) Flush tank. (10) Empty tank as completely as possible to prevent buildup of oil or emulsifiable concentrate residue. Maintain agitation to avoid separation of materials. (11) If an oil or emulsifiable concentrate tank starts to build up in tank, drain and clean with strong detergent solution or solvent. (12) Clean sprayer thoroughly immediately after use by flushing system with water containing a detergent.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

#### GENERAL INFORMATION

ATRAZINE SF will control most annual broadleaf and grass weeds in corn, sorghum, sugarcane, and in certain other crops as specified on this label such as barnyardgrass, witchgrass (Panicum capillare), yellow foxtail, green foxtail, wild oats, large (hairy) crabgrass, giant foxtail, velvetleaf, morning glory, lambsquarters, pigweed, ragweed, nightshade, purslane and mustard. It is also effective in non-crop areas and industrial sites for control of most annual and many perennial broadleaf and grass weeds. ATRAZINE SF may be applied before or after weeds emerge.

In each case where a range of rates is given, the lower rate should be used on coarse textured soil and sows low in organic matter, and the higher rate should be used on fine textured soil and sows high in organic matter.

Following many years of continuous use of this product and chemically related products, biotypes of some of the weeds listed on this label have been reported which cannot be effectively controlled by this and related herbicides. Where this is known or suspected and weeds controlled by this product are expected to be present along with resistant biotypes, we recommend the use of this product in combinations or in

#### INSTRUCTIONS FOR SAFE HANDLING

Avoid application directly to animals or humans. Flagmen or loaders should avoid inhalation of spray mist or contact with skin and should wash thoroughly before eating and at the end of each day's operation. See Precautionary Statements for further information.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

#### Instructions For Use

**Corn:** Atrazine 5F may be applied either before planting, at planting or after planting as indicated below.

#### Preplant (Broadleaf and Grass Control):

Broadcast in the spring after plowing at the rate indicated in Table I. Application may be made before, during or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation of Atrazine 5F. Best results have been obtained when Atrazine 5F is applied within two weeks prior to planting.

#### Pre-emergence (Broadleaf and Grass Control):

Apply during or shortly after planting prior to weed emergence at the rate indicated in Table I.

#### Post-emergence (Broadleaf and Grass Control):

Apply before weeds exceed 1 1/2 inches in height, at the rate indicated in Table I.

TABLE I

#### For Control of Broadleaf and Grasses\* Atrazine 5F

Soil Texture	Broadcast Rate/A
Light Soils: sands, loamy sands and sandy loams	3.25 pints
Medium Soils: silt and clay loams that are low in organic matter	3.80 pints
Heavy soils: silt and clay loams with medium to high organic matter and clay (including the dark prairie soils of the Corn Belt)	4.80 pints
Peats, muck and high organic clays (apply post-emergence only)	4.80 pints

\*For Preplant or Pre-emergence Applications in Western Kansas, Western Nebraska, Eastern Colorado, Eastern Wyoming, New Mexico, West Texas and the Pan Handle of Oklahoma. On sands, loamy sands, sandy loams, mild to strongly alkaline soils and all recently leveled soils, apply 1.9 pints per acre for broadleaf weed control. Broadleaf weeds such as pigweed, lambsquarters, nightshade, purslane and kochia will be controlled. On other soil types in the areas above, make applications at the rate shown in Table I for broadleaf and grass control.

**Layby Treatment (Broadleaf and Grass Control):** Broadcast 1.6-3.2 pints per acre at a minimum of 5 gals. of water or nitrogen solution before weeds are 1 1/2 inches high and corn is 12 inches high. When nitrogen

solutions are used, direct the spray to the lower 1/4 inches of cornstalk to avoid corn foliage injury. Agitation in the spray tank during application is essential. **Postemergence Broadleaf and Grass Weed Control With Atrazine 5F Plus Emulsifiable Oil In Water:** Broadcast 3 1/4 pints per acre after weed emergence, but before weeds reach 1 1/2 inches in height. Add emulsifiable oil at the rate of 1 gallon per acre for ground applications and 1/2 gallon per acre for aerial application.

**Postemergence Broadleaf Weed Control With Atrazine 5F Plus Emulsifiable Oil In Water:** Broadcast 1.9 pts. per acre for control of broadleaf weeds such as annual morning glory, cocklebur, lambsquarters, mustard, pigweed, ragweed, smartweed, and wild buckwheat. Application should be made before pigweed and lambsquarters reach six inches in height and before all other weeds reach four inches in height. Add emulsifiable oil at the rate of 1 gallon per acre for ground application and 1/2 gallon per acre for aerial applications. A cultivation may be necessary if all weeds are not controlled or if regrowth of weeds occurs.

#### Precautions for Applications of Atrazine 5F Plus Emulsifiable Oil In Water to Corn.

1. Do not use oil in Atrazine 5F sprays when corn is under stress from prolonged cold, wet weather, poor fertility, or other factors, or when corn is wet and succulent from recent rainfall as crop injury may occur.
2. Do not use oils in sprays when treating inbred lines or any breeding stock as injury may occur.
3. Adding other insecticides, herbicides, liquid fertilizers or other materials is not recommended with Atrazine 5F and emulsifiable oil in water because they cause compatibility problems or crop injury.
4. Store and handle emulsifiable oil carefully. Oil contaminated with even a small amount of water may not emulsify properly when added to the tank.
5. Do not make more than one application of Atrazine 5F and emulsifiable oil in water per season (except as recommended for control of yellow nutsedge and Canada thistle on this label).

#### PROBLEM WEEDS

##### For control of yellow nutsedge and Canada Thistle

Atrazine 5F will control yellow nutsedge (*Cyperus esculentus*) and Canada thistle (*Cirsium arvense*) when applied as indicated below. For best results, Atrazine 5F should be used each year until yellow nutsedge or Canada thistle is eliminated or reaches a level of infestation where neither weed species is a problem. If regrowth of yellow nutsedge or Canada thistle occurs following the last application of Atrazine during the season, cultivate once. When Atrazine 5F is applied postemergence to the weeds, 1 gallon of emulsifiable crop oil should be added per acre.

For control of yellow nutsedge and Canada thistle with Atrazine 5F several alternative methods of use are available. These methods are listed in order of preference below. If other weed species, such as annual grasses, are also expected, use an alternative method that includes a preplant or pre-emergence oil or apply all the Atrazine 5F preplant or pre-emergence.

1. Broadcast 2 1/4 pts. of Atrazine 5F plus 1 gallon of emulsifiable oil per acre after the crop has emerged and after yellow nutsedge or Canada thistle plants emerge, but before yellow nutsedge plants reach a height of 3 inches or Canada thistle plants reach a height of 6 inches. Follow with a second application of 2 1/4 pts. of Atrazine 5F plus 1 gal. of oil per acre 10 to 20 days but prior to lay-by (12 inches) after the initial application has been made.
2. Broadcast 2 1/4 pts. of Atrazine 5F per acre preplant. Follow with an application of 2 1/4 pts. of Atrazine 5F plus 1 gallon of oil per acre after the corn and weeds emerge, but before nutsedge plants reach a height of 3 inches (for nutsedge control only).
3. Broadcast 2 1/4 pts. of Atrazine 5F per acre during or shortly after planting, but prior to crop or weed emergence. Follow with an application of 3.2 pts. of Atrazine 5F plus 1 gallon of oil per acre after the corn and weeds emerge, but before yellow nutsedge plants reach a height of 3 inches or Canada thistle plants reach a height of 6 inches.
4. Broadcast 4 1/2 pts. of Atrazine 5F plus 1 gallon of oil per acre after the crop has emerged, but prior to lay-by (12 inches) and after yellow nutsedge and Canada thistle plants emerge, but before nutsedge reaches a height of 3 inches or Canada thistle is more than 6 inches tall.
5. Broadcast 4 1/2 pts. of Atrazine 5F per acre preplant (for yellow nutsedge control only).

6. Broadcast 4.8 pts. of Atrazine 5F per acre during or shortly after planting, but prior to crop or weed emergence. (For yellow nutsedge control only).

**NOTE:** Oil should not be used in Atrazine sprays when corn is wet or under stress, especially when using 4.8 pts. of Atrazine 5F per acre. See precaution section under Post-emergence Application of Atrazine 5F plus Emulsifiable Oil in Water on Corn for additional directions.

#### For Quackgrass Control On Land Going Into Corn Production:

**Split Application:** Broadcast 2.4 pts. per acre in the fall or spring and plow 1-3 weeks later.

Broadcast a second application at the rate of 2.4 pts. per acre in the spring before, during or after planting, but before weeds are 1.5 inches high. This split application will control both quackgrass and most annual broadleaf grass weeds.

**Single Application:** Broadcast 4.8 pts. per acre in the spring. Plow 1-3 weeks after application.

#### Tank Mixtures with Atrazine 5F Atrazine 5F plus Paraquat CL

For control of existing vegetation and residual control where corn will be planted directly into a cover crop, established sod or in previous crop residues—Broadcast 3.25-4.8 pts. Atrazine 5F and 1-2 pts. Paraquat CL per acre in 20-60 gallons of water per acre. Add 8 oz. of nonionic surfactant per 100 gallons of diluted spray. Add Atrazine 5F to spray tank first and thoroughly mix with water. Add the Paraquat and surfactant last.

Refer to the Paraquat CL label for further directions, limitations and cautions.

#### Atrazine 5F plus Alachlor 4EC

Use a tank mixture of Atrazine 5F plus Alachlor 4EC for the control of most annual broadleaf and grass weeds in corn (field and silage corn only) such as barnyardgrass, black nightshade, bracharia, carpetweed, crabgrass, fall panicum, Florida pusley, giant foxtail, green foxtail, yellow foxtail, goosegrass, lambsquarter, pigweed, purslane, mustard, common ragweed, smartweed, and witchgrass. This combination also reduces competition from buttonweed (velvetleaf), cocklebur and annual morningglory.

Make applications at the rate indicated in Table II. For preplant applications, apply within 7 days of planting and incorporate to a depth of 2 to 3 inches. For pre-emergence applications, apply to the soil surface after planting, but before the crop or weeds emerge. For post-emergence treatments, make applications until weeds reach the two-leaf stage, and the corn is no more than three inches tall.

**TABLE II**  
Tank Mixtures of Atrazine 5F Plus Alachlor 4EC On  
Corn (Field and Silage Only)

Broadcast Rate Per Acre

Soil Texture	Less than 3% organic matter		More than 3% organic matter	
	Atrazine 5F (Pts/A)	Alachlor 4EC (Qts/A)	Atrazine 5F (Pts/A)	Alachlor 4EC (Qts/A)
Light sandy soils	1.75	1.5	1.75	1.5
Medium loam soils	1.75-1.9	1.75	1.75-1.9	2.0
Heavy clay soils	1.9-2.5	2.5	1.9-2.5	2.5

**NOTE:** Apply in a minimum of 20 gallons of water per acre. Non-pressure fluid fertilizer may replace all or part of the water used as a carrier for applications applied to the soil surface before crop emerges. Add the Atrazine 5F to the spray tank first and thoroughly mix with water. Add the alachlor last. At least 12 weeks must elapse following treatment with Atrazine 5F plus Alachlor 4EC before immature corn forage can be harvested or fed to cattle. Refer to the Alachlor 4EC label for further directions, limitations and cautions.

#### Atrazine 5F plus Propachlor 65W

Use a tank mixture of Atrazine 5F plus Propachlor 65W for control of most annual broadleaf and grass weeds in corn (field, hybrid seed, silage and sweet corn only), such as annual ryegrass, barnyardgrass (watergrass), carpetweed,

crabgrass, fall panicum, Florida pusley, giant foxtail, green foxtail, yellow foxtail, goosegrass, groundsel, jimsonweed, lambsquarters, mustard, nightshade, pigweed, purslane, ragweed, smartweed and sunflower. This combination also reduces competition from buttonweed (velvetleaf), cocklebur and annual morningglory.

Broadcast 1.75-2.5 pts. of Atrazine 5F plus 3.8 to 6.0 lbs. of Propachlor 65W per acre on the surface any time from immediately after planting until broadleaf and grass weeds reach the two-leaf stage. Use the lower rates of Atrazine 5F and Propachlor 65W on the light-textured soils low in organic matter. Use the higher rates on heavy-textured soils high in organic matter.

A minimum of 2.25 pts. per acre of Atrazine 5F in the tank mixture will give better control of annual morningglory, buttonweed (velvetleaf), cocklebur and sunflower.

Apply in a minimum of 20 gallons of water per acre. Non-pressure fluid fertilizer may replace all or part of the water used as a carrier for applications applied to the soil surface before crop and weeds emerge. Add the Atrazine 5F to the spray tank first and thoroughly mix with water. Cut the Propachlor 65W bag and fill with water to the full line. Grasp the neck of the bag firmly and shake vigorously and add to the spray tank.

Refer to Propachlor 65W label for further directions, limitations and cautions.

#### Precautions For All Applications to Corn

1. Post-emergence application to corn must be made before corn reaches 12 inches in height.
2. The maximum application rate for corn is 5 lbs. A.I. per acre per calendar year. Applications for quackgrass suppression in corn are restricted to spring applications only. No fall applications are permitted.
3. To avoid crop injury and illegal residues do not apply more than 4.8 pts. per acre of ATRAZINE 5F to corn in one year.
4. Following harvest of a treated crop, plow (moldboard disk-plow) and thoroughly till the soil in the fall or spring to minimize possible injury to rotational spring seeded crops, regardless of the rate used.
5. Do not graze treated area or feed treated forage to livestock for 21 days following application or illegal residues may result.

For post-emergence applications plus emulsifiable oil, see additional precautions under that section of this label.

#### Suggestions for Rotational Crops

1. Land treated with Atrazine 5F should not be planted to any crop except corn or sorghum until the following year or injury may occur.
2. If Atrazine 5F is applied after June 10, do not rotate with crops other than corn or sorghum the next year or injury may occur.
3. If Atrazine 5F is used at a broadcast rate higher than 4.8 pts. per acre (or comparable rates in a band application), a crop of untreated corn or sorghum should precede the next rotational crop.
4. In the High Plains and Intermountain area of the West where rainfall is sparse and erratic or where irrigation is required, use Atrazine 5F for control of weeds in corn only when corn is to follow corn or a crop of untreated corn, or sorghum is to precede other rotational crops.
5. In Western Mountain and eastern parts of the Dakotas, Nebraska and Kansas, corn treated with Atrazine 5F should not be followed with soybeans if the broadcast rate applied was more than 2.2 pts. per acre (or comparable rate in a band) or injury may occur.
6. Injury may occur to soybeans planted in North Central Iowa and South Central Minnesota the year following an Atrazine 5F application on Harps, Canisteo, Stroden or other soils having a calcareous surface layer.
7. Do not plant sugar beets, tobacco, vegetables (including dry beans), spring-seeded small grains or small-seeded legumes and grasses the year following Atrazine 5F application or injury may occur.

#### Sorghum and Sorghum-sudan Hybrids (Grain and Forage Types):

Atrazine 5F may be applied either before planting, at planting or after planting as indicated below.

#### Preplant (Broadleaf and Grass Control):

Broadcast in the spring after plowing at the rate indicated in Table III. Application may be made before, during or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation of Atrazine 5F. Best results have been obtained when Atrazine 5F is applied within two weeks prior to planting.

Pre-emergence Broadleaf and Grass Weed Control: Apply during or shortly after planting, but prior to weed emergence at the rate indicated in Table III.

and the desert regions of California (Arizona, apply when sorghum is 6-10 inches in height, but before it reaches a boot stage. In all other areas, apply after sorghum reaches the three-leaf stage. Add emulsifiable oil at the rate of 1 gallon per acre for ground applications and .5 gallon per acre for aerial applications. A cultivation may be necessary if all weeds are not controlled or if regrowth of weeds occur.

**Precautions for Application of Atrazine 5F plus Emulsifiable Oil in Water to Sorghum:**

1. Do not use oil in Atrazine 5F sprays when sorghum is under stress from prolonged cold, wet weather, poor fertility or other factors or when sorghum is wet and succulent from recent rainfall as crop injury may occur.
2. Do not use oil in Atrazine 5F sprays when treating inbred lines or any breeding stock as injury may occur.
3. Adding other insecticides, herbicides, liquid fertilizers or other materials is not recommended because they may cause compatibility problems or crop injury.
4. Store and handle emulsifiable oil carefully. Oil contaminated with even a small amount of water may not emulsify properly when added to the tank.
5. Do not make more than one application of Atrazine 5F emulsifiable oil in water per season.

**Post-emergence Broadleaf Weed Control with Atrazine 5F plus Surfactant in Oklahoma, New Mexico, Texas, Western Kansas, Colorado and Desert regions of Arizona and California only:**

Broadcast 1.9 pts. of Atrazine 5F plus .75-1.5 pts. of surfactant per acre after sorghum reaches 6 inches in height, but before weeds reach 1.5 inches in height. Apply only on sandy loam and finer textured soils.

**Precautions for all Applications to Sorghum:**

1. Post-emergence application to sorghum must be made before sorghum reaches 12 inches in height.
2. The maximum application rate for sorghum is 5 lbs. active ingredient per acre per calendar year. Application for quackgrass suppression in sorghum are restricted to a spring application only. No fall applications are permitted.
3. Heavy rains immediately following application tend to result in excessive concentrations of herbicide in seed furrow, resulting in possible crop injury. Applications to furrow-planted sorghum should not be made until furrows are leveled (plow-in). Deep planter marks or seed furrow should also be leveled before application.
4. Application made to sorghum growing under stress caused by minor element deficiency or to sorghum growing on highly calcareous soils may result in crop injury.
5. Do not graze or feed forage from treated areas for 21 days following application or illegal residues may result.
6. Following harvest of a treated crop, plow (moldboard or disk-plow) and thoroughly till the soil in the fall or spring to minimize possible injury to rotational spring-seeded crops, regardless of rate used.

For applications to furrow-irrigated bedded sorghum in Arizona and California and for post-emergence applications plus emulsifiable oil, see additional precautions under these sections of this label.

**Suggestions for Rotational Crops:**

1. Land treated with Atrazine 5F should not be planted to any crop except corn or sorghum until the following year or injury may occur.
2. If Atrazine 5F is applied after June 10, do not rotate with crops other than corn or sorghum the next year or injury may occur.
3. If Atrazine 5F is used at a broadcast rate higher than 4.8 pts. per acre (or comparable rates in a band application), a crop of untreated corn or sorghum should precede the next rotational crop.
4. In the High Plains and intermountain areas of the West where rainfall is sparse and erratic or where irrigation is required, use Atrazine 5F to control weeds in sorghum only when sorghum is to follow sorghum or a crop of untreated sorghum, or corn is to precede other rotational crops.
5. In Western Minnesota and Eastern parts of the Dakotas, Nebraska, and Kansas, sorghum treated with Atrazine 5F should not be followed with soybeans if the broadcast rate applied was more than 3.2 pints per acre (or comparable rate in a band) or injury may occur.
6. Injury may occur to soybeans planted in North Central Iowa and South Central Minnesota the year following an Atrazine 5F application on Haros, Cansted, Stroden or other soils having calcareous surface layer.
7. Do not plant sugar beets, tobacco, vegetables, (including dry beans), spring-seeded small grains or small-seeded legumes and grasses the year

**TABLE III  
Preplant and Pre-emergence Applications\*  
For Broadleaf and Grass Weed Control in Sorghum**

Soil Texture	Organic Matter	Broadcast Rate of Atrazine 5F Per Acre
loamy sands, sandy loams and sandy clay loams	any level	(except for pre-emergence use on bedded sorghum in Arizona and California as indicated below).
Medium and Heavy Soils:	Less than 1%	
silt loams, clay loams	1% to 1.5%	2.6-3.2 pts.
and clays	More than 1.5%	3.2-3.8 pts.

\*Atrazine 5F should not be applied preplant to sorghum grown in N. Mex., Okla., Texas, Ark., La., Tenn., Miss., Ala., Ga., Fla., S. Car. and N. Car. or pre-emergence to sorghum grown in N. Mex., Texas and Okla., except in N.E. Okla., and the Texas Gulf Coast.

In case of planting failures, sorghum can be replanted into soil previously treated with Atrazine 5F. Do not make a second broadcast application or injury may occur. If Atrazine 80W is applied in a band and sorghum is replanted in the untreated row middles, Atrazine 5F can be applied in a band to the second planting.

**Pre-emergence Broadleaf Weed Control in Furrow Irrigated Bedded Sorghum (Arizona and California only):**

For pre-emergence control of broadleaf weeds such as ground cherry, lambsquarters, morningglory, mustard, pigweed and purslane, broadcast 1.3 to 1.9 pts. per acre. Use the lower rate on fine-textured soils and soils high in organic matter. Make application after bed preparation, during or after planting, but before sorghum and weeds have emerged and before the first furrow irrigation. Several regular irrigations should follow the application, making sure that all soil is thoroughly wet.

**Precautions for Pre-emergence Applications of Atrazine 5F to Furrow Irrigated Bedded Sorghum Grown in Arizona and California:** To avoid possible sorghum injury, do not use on sand or loamy sand soils or on sorghum planted in the furrow. Additionally, applications made to sorghum growing on alkali soils or where cuts, fills or erosions have exposed calcareous or alkali subsoils, may result in crop injury. In case of crop failure, do not replant sorghum for eight months following application. Corn may be planted immediately.

**Post-emergence Broadleaf and Grass Weed Control:** Apply before weeds exceed 1.5 inches in height at the rate indicated in Table IV. Application may be made up to sorghum reaching 12 inches in height.

**TABLE IV  
Post-emergence Broadleaf and Grass  
Weed Control in Sorghum**

Soil Texture	Minimum Height of Sorghum at Treatment	Broadcast Rate/Atrazine 5F
Sandy or loamy sand		DO NOT USE
Sandy loam		See directions for broadleaf weed control below
Silt loam to sandy clay loam	Completely emerged	3.2-3.8 pts.
Oilon and Pullman clay soils	At least 6 inches high	3.2-3.8 pts.
Silty clay loams and heavier soils	Completely emerged	4.8 pts.

**Post-emergence Broadleaf Weed Control with Atrazine 5F plus Emulsifiable Oil in Water:**

Broadcast 1.9 pts. per acre for control of broadleaf weeds such as annual morningglory, cocklebur, lambsquarters, mustard, pigweed, ragweed, smartweed and wild buckwheat. Application should be made before pigweed and lambsquarters reach six inches in height and before all other weeds reach four



**Sugar Cane:** For control of nonrhizomatous weeds, such as crabgrass, jungle rice, wire grass, foxtail, amaranths, Florida's paint brush, fireweed and similar plants. Broadcast 1.75-3.25 qts. per acre in 20-50 gallons of water for adequate coverage of the soil surface at time of planting or ratooning, but before the cane emerges. One additional application may be made over the cane as it emerges, and two additional applications may be made interline after emergence, as a directed spray.

For control of emerged Perennial Weeds (artillery weed) (Florida only): Apply 4-5 pints per acre in 40 gallons of water per acre as a directed spray. Add 2 quarts of surfactant for each 50 gallons of spray and be sure weed foliage is thoroughly covered.

#### Precautions

1. Do not apply Atrazine 5F after "close-in."
  2. Do not apply more than 7.6 qts. of Atrazine 5F to any one crop of cane.
- Forest and Christmas Tree Plantations:** Douglas Fir, Grand Fir, Noble Fir, White Fir, Lodgepole Pine, Ponderosa Pine and Scotch Pine (Pacific Northwest—west of Cascade only). Annual broadleaf and grass weed control. Broadcast 1.75-3.25 qts. in 20-40 gallons of water per acre to assure thorough coverage. For band applications, reduce rate of Atrazine 5F and volume of water in proportion to the acre treated. For example, when treating a 4 foot band over trees planted in rows 8 feet apart, apply .8 to 1.75 qts. per acre. Apply between fall and early spring while trees are dormant or soon after transplanting and before weeds are 1.5 inches high.

**Quackgrass Control:** Broadcast 6.4 qts. per acre in 5 gallons of water for adequate coverage. Apply in fall or early spring while trees are dormant and before weed seedlings are more than 1.5 inches high. This application will also control most annual broadleaf and grass weeds.

#### Precautions:

1. Do not graze treated areas.
2. Do not apply to seedbeds.
3. Do not make more than one application per year.

**Nonselective Weed Control on Noncrop Land:** For use on fence rows, around utility poles, and industrial sites, highway medians and shoulders, railroad right-of-way, lumberyards, petroleum tank farms, and in noncrop areas on farms, such as around buildings, equipment, and fuel storage areas, along fences and lanes. Apply Atrazine 5F at rates recommended below before or soon after weeds begin growth. Post-emergence applications should be made when weeds are young and actively growing. Use higher rates on heavy clay and muck soils. Do not contaminate domestic or irrigation water supplies, or lakes, streams, or ponds. Use sufficient water to assure thorough coverage. Use at least 1 gallon of water for each quart of Atrazine 5F, more if practical. Mechanical by-pass or jet agitation is necessary to keep Atrazine 5F in suspension during application.

Most annual broadleaf and grass weed, broadcast 3.8 qts. to 8 qts. per acre. Hard to kill annual and many perennial broadleaf and grass weeds, broadcast 8 quarts per acre.

Hard to kill biennial and perennial weeds such as bull thistle, broadcast 8 quarts per acre.

For longer residual control in regions of high rainfall and long growing season, broadcast 8 quarts per acre.

For small areas, 3.6 oz. per 1,000 sq. ft. is equivalent to 8 quarts per acre.

#### STORAGE AND DISPOSAL

**STORAGE INSTRUCTIONS:** Storage should be under lock and key and secure from access by unauthorized persons and children. Storage should be in a cool dry area away from any heat or ignition source. Avoid storage at high temperatures. Do not stack over 2 pallets high. Move containers by handles or cases. Do not move containers from one area to another unless they are securely sealed. Keep container tightly sealed when not in use. Keep away from any puncture source. Avoid storage near water supplies, food, feed and fertilizer to avoid contamination. Avoid contamination with acids or alkalis. Store in original containers only. If the contents are leaking or material is spilled follow these steps:

1. Contain spill, absorb with a material such as saw dust, clay granules or dirt.
2. Collect and place in suitable containers for disposal.
3. Wash area with water and soap to remove remaining pesticide.
4. Follow washing with clean water rinse.
5. Place a leaking container in a plastic tub and transfer contents as soon as possible to an empty original container.
6. Do not allow run off to enter sewer or contaminate water supplies.
7. Dispose of waste as indicated below.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

#### WARRANTY—CONDITION OF SALE:

**DA** warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.