

DIRECTIONS:

Apply by pouring directly into pool, or feed through mechanical chlorinator. Apply in the evening or early morning as sunlight dissipates chlorine rapidly.

AT TIME OF POOL OPENING

1. Superchlorinate to satisfy initial chlorine demand of raw water. Apply 48 ounces per 10,000 gallons of water. (Triple the regular dosage).
2. Test the pH of the water with an appropriate test kit. If required, adjust pH to range between 7.2 and 7.6. **ALWAYS MAINTAIN THE pH BETWEEN 7.2 and 7.6 IN ORDER TO PROVIDE EFFECTIVE PERFORMANCE OF CHLORINE.** Test pH frequently.

REGULAR MAINTENANCE:

3. The day following superchlorination, test chlorine level. Apply 16 oz. per 10,000 gal. of water daily, or as needed to maintain a chlorine residual of 0.6 ppm to 1.0 ppm. Test for chlorine frequently using a suitable chlorine test kit. On hot days, when the bathing load is heavy, or when pool water is heated, it may be necessary to repeat dosage more frequently than once a day.

Residuals resulting from recommended dosage are subject to variable conditions such as the temperature of the water, intensity of sunlight, and numbers of swimmers in the pool. Therefore, the recommended dosages should not be assumed precise for all circumstances. Adjustments in dosage quantities should be made on the basis of daily testing for chlorine residuals.

DANGER: To avoid possible irritation of mucous membranes, always wait until chlorine level drops below 2.0 ppm before allowing swimmers to re-enter the pool.

Note: Sodium hypochlorite degrades with age and exposure to heat and light. Keep in a cool dark place. Increase dosage as necessary to obtain required level of available chlorine.

DANGER

Sodium hypochlorite contains 11% available chlorine, which is irritating to eyes, skin and lungs. Sodium hypochlorite is a strong oxidant. It may produce chemical burns. Do not get on skin or in eyes.

FIRST AID:

EXTERNAL: Flush at once with water. Wash with soap and rinse with clear water. If irritation (skin burn) results, see a physician.

EYES: Flush immediately with cold water. Repeat several times. Wash with 1% solution of boric acid. **SEE A PHYSICIAN AT ONCE.** Eyes should be irrigated by professional medical personnel immediately after emergency treatment.

INTERNAL: Drink raw egg white, milk, rice gruel, or mucilage. Follow immediately with an emetic such as a tablespoon of mustard in warm water. **SEE A PHYSICIAN AT ONCE.**

DANGER: Never mix this chlorine solution in the same container with any other chemical. Noxious fumes may result. Always apply unmixed with anything other than water. When container is empty, rinse thoroughly with water and discard.

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public water. Apply this product only as specified on this label.

Dist by.

ALSIP NURSERY
Frankfort, IL

EPA ESTAB. NO. 362881L01
EPA REG. NO. 36288-1

FIVE U.S. GALLONS

20