



CT<sup>®</sup> chem-tox<sup>®</sup>

#### DIRECTION FOR USE

For a description of the Federal Law and  
protection of employees under the Act, see page 1.

#### STORAGE AND DISPOSAL

##### Pesticide Disposal:

Do not contaminate water, food or feed by storage or disposal.


##### Container Disposal:

Do not contaminate water, food or feed by storage or disposal.

Directions for use contained on page 1.

# FOOD PLANT SPRAY

*Designed for Use in Edible Product Areas of Official Establishments Operating Under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs and in Other Food Processing Plants.*



CHEM TOX, Inc.  
BARRINGTON, IL 60010

Order Code  
0-301

#### ACTIVE INGREDIENTS

Pyrethrin  
Methyl Ethyl Ketone  
Petroleum Ether  
  
Equivalent to 4.0% butyl carbamate 6.0% by  
weight of active ingredients in liquid form.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

EPA Reg. No. 45385-8  
EPA Est. N. 45385 IL-01

Net 1 Gallon / 3.785 Liters





**CT** **chem - tox**

## DIRECTION FOR USE

### NOTIONARY STATEMENTS

#### HAZARDS TO HUMANS

#### DOMESTIC ANIMALS

#### AUTION

Inhaled or absorbed through the skin. Avoid mist, and provide adequate ventilation of contact with the eyes. Skin or clothing should contact, immediately flush eyes or skin with medical attention if irritation persists. Do not let pets to contact treated surfaces until processing surfaces, dishes and utensils during treatment, or thoroughly washed before entry to food handling facilities cover or remove any food equipment during application. Do not apply spraying is underway. After spraying in food wash all equipment benches, shelving etc. will be handled with portable water.

#### ENVIRONMENTAL HAZARDS

No fish and wildlife. Keep out of lakes, ponds, contaminate water by cleaning of equipment or

#### PHYSICAL HAZARDS

Or store near heat or open flame

The use of this product in food processing establishments should be confined to time periods when plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Stored food products which are infested should be fumigated or treated by other effective methods using an approved product intended for that purpose.

**SPACE SPRAY:** For control of accessible, exposed stages of flies, including fruit flies, mosquitoes, small flying moths, gnats, wasps, hornets, cheese skippers, and these common stored product pests: granary weevils, rice weevils, confused flour beetle, saw-toothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, Tobacco moths, meal worms, and grain mites. Close windows and doors and shut off ventilating systems. Use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. For best control of Wasps and Hornets, apply directly to nests in the evening when the insects have returned to the nests. Keep the area closed for at least 15

minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

**SPACE AND CONTACT SPRAY:** For control of accessible, exposed stages of silverfish, crickets, roaches, spiders, scorpions, clover mites, cheese mites, earwigs, and ants, first using equipment which can deliver a fine spray, direct spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects are known to seek shelter. Contact as many insects as possible, spray directly all ant trails. Do not allow spray to contact stored food. Then, shutting all doors, windows and shutting off all ventilating systems proceed with space spraying directions as given above. Use dosage rate of 1 ounce per 1000 cubic feet of space. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

**IN MUSHROOM PRODUCTION AND PROCESSING:** To control mushroom flies or fungus gnats in growing houses, apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results, refrigeration should be turned off. In packing and canning plants, use at a rate of 1 ounce to 2000 cubic feet of space.

#### Terms of Sale or Use, or

**Warranty:** CHEM-TOX warrants that its product, as of the time of sale, is CHEM-TOX conforms to the ingredient statement on the label, and is reasonably fit for the principal set forth in the Directions for Use. Except as so warranted, the product is sold as is. CHEM-TOX makes no other warranty express or implied.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price and shall not include incidental or consequential damages. Notice of any claim must be reported to CHEM-TOX within 10 days of discovery by buyer or user failing which buyer or user waives any claim for such damage.

**Excuse from non-controllable conditions:** Because of certain critical conditions created or insured by buyer or user or over which CHEM-TOX has no control, buyer and user, as a condition of purchase or use, assuming responsibility for and release CHEM-TOX from liability arising out of handling or use of the product attributable to such causes. No person is authorized to vary or waive any Statement contained herein.

Buyer assumes all risks of use. Storage or handling of this material must be in strict accordance with Directions Given herewith.