

BEST AVAILABLE COPY

CHEM-TOX

AQUEOUS FOOD PLANT SPRAY

DESIGNED FOR USE IN EDIBLE PRODUCT AREAS OF OFFICIAL ESTABLISHMENTS OPERATING UNDER THE MEAT, POULTRY, SHELL EGG GRADING AND EGG PRODUCTS INSPECTION PROGRAMS AND IN OTHER FOOD PROCESSING PLANTS.

Active Ingredients.

Pyrethrins	0.5%
*Piperonyl Butoxide, Technical.....	5.00%
† Inert Ingredients	92.50%

*Equivalent to 4.0% (butylcarbityl)(6-propylpiperonyl) ether and 1.0% related compounds.

† Contains 2.0% Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for additional precautionary statement

EPA Reg.No: 45385-14

EPA Est.No: 45385-IL-01

Manufactured by
CHEM-TOX, INC.
21n 988 Pepper Rd.
Barrington, IL 60010

Net Content:

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12. Active to Inert Change in Status.

DIRECTIONS FOR USE CONTINUED

Apply this product with conventional mechanical or compressed air equipment designed to deliver a TRUE AEROSOL MIST.

Close windows and doors and shut off ventilation systems. Use a non-thermal fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cu.ft. of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

The above described application will kill accessible, exposed stages of FLIES, including FRUIT FLIES, MOSQUITOES, SILVERFISH, SMALL FLYING MOTHS, GNATS, CRICKETS, ROACHES, SPIDERS, SCORPIONS, WASPS, HORNETS, CLOVER MITES, CHEESE MITES, CHEESE SKIPPERS, and these common stored product pests: GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLE, SPIDER BEETLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, ANGOUMOIS GRAIN MOTHS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, TOBACCO MOTHS, MEAL MOTHS, and GRAIN MITES.

In MUSHROOM PRODUCTION and processing: To control adult PHORIDS, SCIARDS, and CECIDOMYLIDS in growing houses, apply at a rate of 1 ounce per 1000 cu.ft. of space. Fogger should be operated 8 to 10 feet away from beds and direct slightly upward over them. Circulating fans should be left in operation, but for best results refrigeration should be turned off. In packing and canning plants use at a rate of 1 ounce to 2000 cu. ft. of space.

Order Code: 0-306

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in meat packing plants, bakeries and other food processing plants, wash all equipment, benches, shelving, etc. where exposed food will be handled, with an effective cleaning compound followed by a potable water rinse. In the home, all food processing surfaces and materials should be covered during treatment, or thoroughly washed before use.

Remove pets, birds and cover fish aquariums before spraying. **CAUTION- PRE-TEST INSECTICIDE FOR POSSIBLE DISCOLORATION OR SPOTTING OF FABRIC, MATERIAL OR CARPET TEST FOR POSSIBLE ALLERGIC REACTION**

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

DIRECTIONS FOR USE

IT IS A VIOLATION OF THE FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in a locked storage area.

PESTICIDE DISPOSAL: For Containers Over 1 Gallon: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: For Containers 1 Gallon and Under: Do not reuse empty container. Wrap container and put in trash.

For Containers Over 1 Gallon: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Directions for use continued on back panel

