

PYRONOX NO. 3

INSECTICIDE

Industrial, Mill, Livestock and Mosquito Spray

For Rapid Control of Flying and Crawling Insects Named On This Label

Use in Food Processing Plants, Canneries, Packing Houses, Mills, Elevators, Granaries, Warehouses, Box Cars, Ships' Holds, Breweries, Bottling Plants, Bakeries, Stores, Markets, Theaters, Hospitals, Schools, Hotels, Motels, Restaurants, Dairies, Barns and Milk Rooms.

Kills and Repels Flies, Mosquitoes and Gnats on Cattle, Horses and Hogs.

Active Ingredients:
Pyrethrins 0.13%
*Piperonyl Butoxide, Technical..... 1.27%
+ Inert Ingredients..... 98.60%

*Equivalent to 1.02% (butylcarbityl)(6-propylpiperonyl) ether and 0.25% related compounds
+ Contains 98.60% Petroleum Distillate

STATEMENT OF PRACTICAL TREATMENT

If swallowed - Call a Physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. **DO NOT INDUCE VOMITING.** Vomiting may cause aspiration if indicated.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in meat packaging plants, bakeries and other food processing plants, wash all equipment, benches, shelving, etc., where exposed food will be handled, with potable water. In the home, all food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use.

Remove pets, birds and cover fish aquariums before spraying.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Harmful if swallowed.

See side panel for additional precautionary statements.

EPA Est. No. 45385-IL-01
EPA Reg. No. 45385-28

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.


PHYSICAL HAZARDS

Do not use or store near heat or open flame

NET CONTENTS: 5 GALLON

CHEM-TOX, Inc.
BARRINGTON, IL 60010

Order Code:
O-307



Not Reviewed. Registrant claims to be in accordance with reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

DIRECTION FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

Pesticide Disposal:

Pesticide, spray mixture or rinse that cannot be used according to label instruction must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

Container Disposal:

Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other approved State and local procedures

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control exposed or accessible stages of roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, meal worms, grain mites, and cadelles: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets, and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings, and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, mouldings, window and door frames, and all areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

SPACE SPRAYING: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For

rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths: direct spray at an upward angle, distributing it uniformly throughout the entire area, at a rate of one gallon to 200,000 cubic feet of space.

To kill exposed or accessible stages of the above named flying insects, use a rate of one gallon to 80,000 cubic feet of space. Contact insects directly whenever possible.

BARNs, MILK ROOMs, DAIRY PLANTs: To control flies, mosquitoes, and gnats, apply as a fog or fine mist, directing spray toward the ceiling and upper corners, until the area is filled with mist, using about 2 ounces per 1000 cubic feet of space. For best results, close windows and doors before spraying and keep closed for ten to fifteen minutes. Vacate the area and ventilate before reoccupying. Repeat application as necessary.

ON ANIMALs: To protect horses, cattle and hogs from horn flies, house flies, mosquitoes, and gnats, apply a light mist sufficient to wet the tips of the hair. To control stable flies, horse flies, deer flies, apply 2 ounces per adult animal, sufficient to wet the hair but not to soak the hide. Repeat treatment once or twice daily or at intervals necessary to give continued protection.

OUTDOOR MOSQUITO CONTROL: To control adult mosquitoes, apply as a fog or fine mist using good mechanical, thermal or compressed air equipment. For best results, treat when wind velocity does not exceed 5 miles per hour. Application in the evening or early morning is recommended. Treat area thoroughly with particular attention to low and marshy places, shore lines, stagnant pools or ponds and shrubbery or vegetation where mosquitoes may rest. Apply at a rate of 1.2 gallons per acre (or 3.6 fluid ounces per 1000 square feet).

PA 117
45385-28
PA 1491

5/30/90