

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Avoid contact with eyes, skin and clothing. Avoid breathing vapors or spray mist. Keep away from food, feedstuffs and domestic water supplies. Wash hands thoroughly after using and before eating or smoking.

ENVIRONMENTAL HAZARDS

Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of waste.

PHYSICAL HAZARDS

Do not store near heat or open flame.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a Physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. DO NOT INDUCE VOMITING; vomiting may cause aspiration pneumonia. Remove victim to fresh air. Apply respiration if indicated.
If inhaled: Remove contaminated clothing and wash affected areas with soap and water.
If on skin: Flush eyes with plenty of water. Call a physician if irritation persists.

DIRECTION FOR USE

It is a violation of the Federal Law to use this product in a manner inconsistent with its labeling.

Pesticide Disposal:

Pesticide spray mixture or rinseate that cannot be used according to label instruction must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

Container Disposal:

Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other approved State and local procedures.
Directions for Use continued on side panel.

CT chem-tox

GARD-X
Oil Spray

- Dormant and growing season insect spray
- Controls Scale Insects, Mealybugs, Whiteflies, Red Spider Mites and Certain Other Garden Pests
- Acts as excellent leaf polish

EPA REG. NO. 45385-44
EPA EST. NO. 45385-1L-01

KEEP OUT OF REACH OF CHILDREN

CAUTION

Harmful if swallowed.

See side panel for additional precautionary statements

ACTIVE INGREDIENT PETROLEUM OIL 98%
INERT INGREDIENTS 2%

Net 28 - Min. U.S. Gals. - Class Light-Med.

CHEM-TOX, INC.
BARRINGTON, IL 60010

NET CONTENTS
1 QUART

Shake well before using. Use a forceful spray, wetting parts and all parts of plants thoroughly. Do not store diluted spray. Make new dilution for each use.

FOR DORMANT USE - To control Scale Insects, Red Spider Mites, and Mealybugs on Dormant Shrubs, Evergreens (note caution), Woody Plants, Fruit Trees (Apples, Plums, Pears, Apricots, Peaches, Cherries) and Shade Trees: 1/2 pt. 15% oil per gal. water.

GROWING SEASON - To Control Scale Insects, Red Spider Mites, Mealybugs and Whitefly Larvae on Citrus (Oranges, Lemons, Grapefruit, Tangerines, Limes), Fruit Trees (Apples, Plums, Pears, Apricots, Peaches, Cherries), Shade Trees, Evergreens, Camellias, Azaleas, Hardy Shrubs: 3 Tablespoons (2 1/2 oz) per gal. water. This application is also effective in the removal of sooty mold fungus through the control of whitefly larvae.

COMBINATION WINTER SPRAY - FRUIT TREES (Apples, Plums, Pears, Apricots, Peaches, Cherries), ROSES AND ORNAMENTALS (Dormant, deciduous only) - for Control of San Jose Scale, Rose Scale, Quince Shell Scale, Brown Apricot Scale, Black Scale and Overwintering Insect Eggs.

COMBINATION WITH Melathion 50 Insect Spray on Citrus, Avocado, Mangoes, Gardenias and Ilex for Control of Scale Insects, Red Spider Mites, Mealybugs, Whiteflies, Aphids, Thrips and for the removal of Sooty Mold Fungus through control of Whitefly Larvae - Use 1 Tablespoonful Oil Spray and 1 Tablespoonful Melathion 50 Insect Spray for each gal. of diluted spray. May be applied on Avocadoes up to within 7 days of harvest, to Grapefruit, Lemons, Limes and Oranges up to within 7 days of harvest but not during full bloom, and to Mangoes up to within 2 days of harvest.

Helps to destroy insect eggs such as Red Spider Mite Eggs, Oak Moth Eggs and Codling Moth Eggs.

AS A LEAF POLISH - To brighten leaves of hardy house plants. Use 3 Tablespoons (1 1/2 oz) per gal. water.

GREENHOUSES. To control Scale Insects, Red Spider Mites, Whitefly, Mealybugs on Hardy Shrubs, Gardenias, Camellias and Azaleas - 2 Tablespoons (1 oz) per gal. water. Allow at least 60 days between applications and limiting to 3 applications per year.

PALM TREES: Scale Insects, Mealybugs, Red Spider Mites - 2 Tablespoons (1 1/2 oz) per gal. water. Note Caution. **AZALEAS, CAMELLIAS**: Scale Insects, Red Spider Mites, Mealybugs, Whitefly Larvae, Lace Bug Larvae - 3 Tablespoons (2 1/2 oz) per gal. water. Apply in spring after blooming and/or fall prior to blooming.

N SPRAYETTE Garden Hose Sprayer to use - Follow directions on the bottle.

Do not use Spray on foliage within 30 days before or after using Sulfur. Avoid spraying when temperature is excessively high or when plants are suffering from drought. Oil sprays should not be applied to broadleaved evergreens or palms when freezing or near-freezing temperatures may be expected. Do not use on Maidenhair Ferns, Cocos Palms, Maples, etc. or non-woody plants unless specified. Oil Spray may cause discoloration to Blue Spruce. Do not use in combination with or closely following treatment with Garden Fungicide.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

ACCEPTED

FEB 15 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act, amended, for the pesticide registered under EPA Reg. No. 45385-44