

### DIRECTION FOR USE

It is a violation of the Federal Law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in original container in a locked storage area.

**PESTICIDE DISPOSAL:** For Containers Over 1 Gallon: Ways resulting from the use of this product may be: incineration on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** For Containers 1 Gallon and Under: Do not reuse empty container. Wrap container and put in trash.

For Containers Over 1 Gallon: Triple rinse (or equivalent). Then offer for recycling or reuse, if available, or puncture and dispose of in a certified landfill. Only those pesticides approved by the EPA are eligible for such treatment.

Directions for use continue on back panel.

# ACCEPTED

## FEB 26 1991

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 45385-47

ORDER CODE  
W-105

CHEM TOX, Inc  
BARRINGTON, IL 60010

# DURSBAN\*plus INSECT KILLER

## A COMBINATION OF INSECTICIDES OFFERING GOOD FLUSHING ACTIVITY, QUICK KNOCKDOWN AND RESIDUAL KILL OF COCKROACHES FOR UP TO 4 WEEKS.

### FOR RESIDUAL AND CONTACT CONTROL OF COCKROACHES, CRICKETS, SILVERFISH, CLOVER MITES, CENTIPEDES, MILLIPEDES, FIREBRATS, EARWIGS, SOWBUGS, SPIDERS, BROWN DOG TICKS, LONE STAR TICKS, ANTS, CARPET BEETLES, FLEAS. CAN BE USED IN NON-FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS

<b>ACTIVE INGREDIENTS:</b>	
1% Pyrethrin	0.10 % A.A.
• Chlorpyrifos (0.0 diethyl 0.3 5.6 trichloro 2 pyridyl) phosphorothioate	0.50
<b>INERT INGREDIENTS</b>	99.40
* Cis:trans isomers ratio: max 30% (1.0:0.9) and min 70% (0.9:1.0)	
U.S. Patent Nos. 3,465,007 & 3,542,928	

• U.S. Patent No. 3,244,596. Dursban is a trademark of Dow Chemical Co.  
\*\* U.S. Patent No. 3,683,078

### KEEP OUT OF REACH OF CHILDREN CAUTION

Harmful if swallowed  
See back panel for additional precautionary statements.

EPA Reg. No. 45385-47  
EPA Est. No. 45385 IL-01

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid contact with eyes, skin or clothing. Wash hands after using. Do not contaminate food or feedstuffs with this spray. Do not allow spray to contact food, feedstuffs or animal supplies. Thoroughly wash food contact surfaces with soap and water if they become contaminated by application of this product. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and livestock from areas where spraying. Do not use this product in or on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Do not induce vomiting. Call a physician or Poison Control Center immediately. **IF INHALED:** Remove to fresh air. **IF ON SKIN:** Remove contaminated clothing and wash affected areas with soap and water. Get medical attention if irritation persists. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists. **NOTE TO PHYSICIAN:** Contains piperidinium distillate. Vomiting may cause aspiration pneumonia. Chlorpyrifos is a cholinesterase inhibitor. If symptoms of cholinesterase inhibitor appear, atropine only by injection is an antidote.

### ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish, birds, and other wildlife. Do not apply directly to water. Drift and spray off from treated areas may be hazardous to aquatic organisms in adjacent streams, rivers, or impounded spills. Do not contaminate water by cleaning of equipment or disposal of waste.

For use in non-food areas indoors and outside premises of homes, hotels, preschools, apartment buildings, commercial offices, educational and storage buildings, recreational buildings, boats, ship cabins, and hospitals. **IN HOSPITALS,** remove patients prior to treatment. Ventilate room for two hours before returning patients. This product can be used in non-food areas only, such as corridors, offices, elevators, lavatories, garage, utility rooms, storage rooms and basements, of packaging and processing plants, meat handling areas of food plants, in a storage room for egg processing plants, supermarkets, restaurants, buffing plants, canneries, flour and feed mills and dairies.

Avoid for use in and around campers, mobile homes, buses and other motor vehicles. This product can be used in garages, planes, boats, cars, and ships, but only if they are well ventilated. Do not use on food or food products or on animals.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

**ATTENTION: DO NOT INDOORS!** Do not use in food areas indoors and outside premises of homes, hotels, preschools, apartment buildings, commercial offices, educational and storage buildings, recreational buildings, boats, ship cabins, and hospitals. **IN HOSPITALS,** remove patients prior to treatment. Ventilate room for two hours before returning patients.

This product can be used in non-food areas only, such as corridors, offices, elevators, lavatories, garage, utility rooms, storage rooms and basements, of packaging and processing plants, meat handling areas of food plants, in a storage room for egg processing plants, supermarkets, restaurants, buffing plants, canneries, flour and feed mills and dairies.

Avoid for use in and around campers, mobile homes, buses and other motor vehicles. This product can be used in garages, planes, boats, cars, and ships, but only if they are well ventilated. Do not use on food or food products or on animals.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.

Do not use on plants, fruits, or vegetables, processing plants, restaurants, or other areas where the food is commercially processed or prepared. Do not use on sheep, goats, cattle or birds exposed.