45385-69 3/16/2006



# UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

e 173

MAR 16 2006

H.R. MCLANE, AGENT FOR CTX-CENOL, INC. 7210 Red Road, Suite 206A Miami, Florida 33143-5321

Subject: Amendment - Addition of Clothing Treatment Perma-Tox Insecticide E.C. EPA Reg. No. 45385-69 Your application dated September 1, 2005

Dear Mr. McLane:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable subject to comments listed below. Two (2) copies of the finished labeling must be submitted prior to releasing the product for shipment. A stamped copy of the label is enclosed for your records.

- 1. Delete reference to military clothing, uniforms, gear, battle dress uniform, the MSN and military specification numbers. The product you cited with these uses is for military use only. There was no information submitted authorizing use of this product for military purposes nor information stating that this product formulation meets military specifications. Thus limit the directions and use sites to "Outer Clothing and Bed Nets only.
- 2. Revise the mixing directions to reflect a 0.5% application rate, i.e. 1% oz in 1 quart of water. This is the maximum application rate for consumer treated clothing.
- 3. Delete the misuse statement under directions for outer clothing and bed nets. It already appears under Directions for Use on the left panel of the label.
- 4. Under Directions for use delete the statements beginning, "Keep away from food, eating utensils... and ending....contaminated with this product". These precautions pertain to miltary uses.
- 5. Following the last sentenced under directions for outer clothing and bet nets add, "For further protection use in conjunction with an insect repellent registered for direct application to exposed skin".
- 6. Delete the first sentence under application procedure. This precaution pertains to EPA Reg. No. 63021-1.

- 7. Revise the first statement under For Clothing to read, "Dilute 11/2 oz into 1 quart of water in a 1 quart hand trigger sprayer". Replace the statement "Do not treat clothing a second time; one treatment is effective for life on the garment", with," "Follow standard laundering procedures. Clothing should be retreated after four weeks or after the sixth laundering to maintain adequate protection. Do not retreat clothing more than once every two weeks. Launder clothing at least once before retreating".
- 8. Note that this product is in child resistant packaging. Therefore you cannot have a separate 2 ounce bottle. The product must be mixed and diluted from the original container as indicated in comment 7 above.

Submit a formulators exemption statement and a updated data matrix that includes product chemistry and acute toxicity data. You only listed efficacy data on the matrix.

If you have any questions in regard to this letter please feel free to contact me at 703-305-6100.

Sincerely,

George T. LaRocca Product Manager 13 Insecticide Branch Registration Division (7505C)

チャろ

Enclosure

## PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS. WARNING: Causes eye irritation, Do not get in eyes. Avoid contact with skin or clothing. Harmful if swallowed, Avoid breath ing vapor. Wash thoroughly after handling. For space spraying indoors wear NIOSH or MSHA approved respiratory protection device. Do not allow people or pets to enter treated areas until spray mist or fog has dissipated and treated surfaces are dry. FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IFINEYES-Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact tenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or rinctor for treatment arhites

IF ON SKIN OR CLOTHING: Take off contaminated clothing Rinse skin immediately with plenty of water for 15 to 20 minutes Call a poison control center or doctor for treatment advice.

IF INHALED. Move person to fresh air. If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible. Call a poison control center or dector for further treatment advice

Have this product container or label with you when calling a poison control center, or doctor, or when coing for treatment, You may also cell 1-800-858-7378 for emergency medical ireatment advise, \*\*NOTE TO PHYSICIAN: Solvent presents aspiration hazard. Gastric Lavage is indicated if taken internally.

#### **ENVIRONMENTAL HAZARDS**

This product is extremely toxic to fish. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.<u>BEE WARNING</u>. This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow drift when bees are actively visiting the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity. PHYSICAL and CHEMICAL HAZARDS

Do not use or store near heat or open flame. Flashpoint minimum of 105°F (41°C)

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manne inconsistent with its labeling.

### STORAGE AND DISPOSAL

STORAGE: Store this product in its original container in a dry cool, secured area out of reach of children. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal PESTICIDE DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the container be used, carefully following label directions and precautions. Pesticide wastes are toxic. Improper disposal of excess pesticide, or rinsate is a violation of Federal law. If

these wastes cannot be disposed of by use, according to label instructions, they must be disposed of at, or by, an Approved Weste Disposal Facility, or contact your State Pesticide or Envi ronmental Control Agency, or the Hazardous Waste represen tative at the nearest EPA regional office for guidance. IF EMPTY: Do not reuse this container. Place in trash or offe

IF PARTLY FILLED: Call your local waste agency or 1-800 CLEANUP for disposal instructions. Never place unused procuct down any indoor or outdoor drain. SPILL: In case of spill absorb liquid with absorbent material such as soil or clay etc. Pface absorbed split in a labeled container and dispose of ac-cording to PESTICIDE DISPOSAL directions above. 07.24001

# CHEMotox PERMA-TOX INSECTICIDE

AN INSECTICIDE OFFERING: SPACE CONTROL OF FLYING INSECTS. RESIDUAL KILL OF CRAWLING INSECTS FOR UP TO 4 WEEKS. INDOORS AND OUTDOORS - FOR TREATMENT OF SPORTING and, MILITARY

CLOTHING, UNIFORMS, GEAR, AND BED -NETS.

# **KEEP OUT OF REACH OF CHILDREN** WARNING

STOP: Read Label Before Use

See side panels for additional Precautionary Statements.

#### ACTIVE INGREDIENT:

*Permethrin CAS[52645-53-1]	13.30%
OTHER INGREDIENTS**:	<u>86.70%</u>
Total	100.00%
"Cis/trans isomers ratio: min. 55%(±)cis and max.	45%(±)trans.

\* \*Contains Petroleum Distillate

EPA REG. NO. 45385-69 EPA EST. NO. 56--OH-01

Net Contents: 2 ounces (60 ml) - 1 pint: 1 quart - 1 Gallon / (3.785 Liters)

# MSN 6840-01-334-2666 Perfected in the USA Military Specification MIL-1-44411(1) DIRECTIONS for use on OUTER CLOTHING & BED NETS

If is a vicibility of Federal law to use this product in a manner inconsistant with its labeling. Do not apply to skin or to the inside surfaces of clothing. Keep away from food, eating densits and water supplies. Thoroughly wash any sens contaminated with this product. Wear chemical resistant playes any write contaminated with this product, treast chemical resistant govers which mixing and using. Treasted garments, uniform, netting and teniage provides protection to covered areas of body from bites of mosquitoes, licits, chiggers, and iterate body fice. APPLICATION PROCEDURE: To avoid initiatation and contact, applica-

APPLICATION PROCEDURE: To evold initiatation and contact, applica-tor must wear a respirator and chemical resistant gloves. Thoroughly clean a quart hand ringer Sprayer by trike insign with clean weter. FOR CLOTHING: Place entire contents of this 2 ounce bottle into 1 quart of clean water in a 1 quart and trigger sprayer. Apply only to the outside surfaces of outer clothing. Do not breat the inside surface. Place each stride of clothing, (sporting, battisk, dress, uniform, jacket and/or insurers) or the seconds. Do not spray mist only the cutside of the clothing fabric for 3 to 6 seconds. Do not spray to wet the labit. A fight spray mest is all that is required. Hang the insafed garment to thoroughly dry. When dry, it is reach to waar, Mark clothing with date of reatment. Do not treat under-wear, freedowr, or the inside surfaces of the outer clothing. FOR BED NETTING : Spread netling the ringer spray mist at a distance of 12 to 15 inches using the ringer spray mister. Spray with a alow sweeping motion to completely cover the netling fabric without tunoff. Allow to thoroughly dry and aerate before use. est-se

WARRANTY STATEMENT Our recommendation for the use of this product are ; based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no quarantee, expressed or implied. is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damages, resulting] from its misuse as such, or, in combination with other materials.

# **DIRECTIONS FOR USE (Continued)**

FOR USE IN NON-FOOD AREAS INDOORS AND OUTSIDE PREMISES OF MOTELS, HOMES, PRESCHOOLS, DAY CAMPS, HOME GREEN-HOUSES, SEED STORAGE HOUSES, FOOD HANDLING ESTABLISH-NUDSES, DEED STURAGE NUDSES, FUOD HANDLING ESTABLISH MENTE, RACKAGING AND PROCESSING PLANTS, COMMERCIAL, IN DUSTRIAL, OFFICES, INSTITUTIONAL AND STORAGE BUILDINGS, DARRIES, MILK STORAGE ROOMS, EGG PROCESSING PLANTS, SU-PERMARKETS, RESTAURANTS, BOTTLINGPLANTS, CANNERIES, FLOUR AND FEED MILLS, BAKERIES, AND HOSPITALS. IN HOSPITALS, PA AND FEED MILLS, BUNERIES, AND HOUSH (ALS, IN HOUSH (ALS, PA-TIENTS SHOULD BE REMOVED PRIOR TO TREATMENT. ALSO FOR USE IN DOG KENNELS AND HORSE STABLES. IN HORSE STABLES NOT FOR APPLICATION IF HORSES WILL BE USED FOR FOOD.

Do not use in food areas of food handling establishments, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed or larility is in operation. Serving areas are areas where prepared foods are served such as clining rooms, bu excluding areas where foods may be prepared or held,

Nonfood areas are areas such as garbage norms, lavatories, floor thains (to severa) entries and vestibules, offices, locker rooms, machine rooms; boller norms, garages, mor closets and storage (after canning or bottling). One gallon of finished spray is sufficient to treat 1580 linear feet (482 m with a 6 inct (1.5.2 cm) spray band. For contact and residual sprays, mix PERMA-TOX with water in the following manner:

FINISHED SPRAY	CHEM-TOX PERMA-TOX
0.25%	2-2/3 02. or 79 mills 1 Gal. or 3.79 L Water
0.25%	2/3 02. or 20 mills 1 Cluart or 0.96 L Water
0.50%	5-1/2 02. or 163 mills 1 Gal. or 3.79 L Water
0.50%	1-1/3 02. or 169 mills 1 Gal. or 3.79 L Water

### INDOOR PEST CONTROL

RESIDUAL SPRAY and contact control of cockroaches (waterbugs), crickets, silverfish, (firebrats), clover mites, millipedes, cantipedes, earwigs, sowbugs (pillbugs), spiders, brown dog licks, ione star licks, anis, fire anis, evencuja cancuja, spolens, orown oog icos, one star ucos, ants, me ants, tobhes moths, carpet beeles, lexs, boelder bugs, scorpions, spiler beele, bed bugs, grasshoppers, house files, mosquitoes, gnats, fuit files, duster files, filmig moths, contused flour beelles, asw-counted grain beelles, stimu spider beeltes, grain mites, red flour beeles, Angoumois grain moths, grain weevis, Indian meal moths, Meditemanean flour moths, lesseer grain bores. cigarette bestles, drug slore beetles, chocciate moths. Spray thoroughly into hiding places such as cracks, crevices, moist areas.

ings around pipes and sinks, under refrigerators, baseboards and stor-

The backs of the second second

to dry Inductive use. <u>GRAIN INSECTS:</u> To control confused four beetles, saw-toothed grain beolies, spider beetles, shiny spider beetles, grain mites, red four beetles, Angoumois grain moth, grain weeval, Indian moai moth, Mediterranean four Angournos grain motir, grain weeve, indian moai moth, Mediterranean flour moths, lessor grain borer, digaratte beetise, drug store beeties, chocolate motins, carpot beeties, - remove infested grain, cereal or four and destroy. Avoid contamination of other food and loodstiffs in the area by removing before spraying. Spray directly on exposed stages and into crack and cravices where grain insects may be hiding, in <u>Jonnes</u>, dispose of all infested products. The shelves and drawers should be dispose of all infested products. The shelves and drawers should be disposed for all outputs drawed on ex to confer all there and the bard with the first the first sectore and the shell be and the shell be all there and the shell be all the shell be a messed products. The sneves and drawers should be created and thor-oughly sprayed as as to contact all pests, and covered with shelf paper after allowing shelves to div. Any utensits that are contacted with spray should be washed before using. Repeat as necessary. SPIDERS: Spray directly on these pests and use repeated applications to the webbing and all cracks and cravices which harbor spikers. ANTS (including FIRE ANTS): To Kill ants, spray tents, nests and points of entry within buildings and homes. Spray directly on ants where possible.

Repeat as necessa

VASPS, HORNETS, YELLOW JACKETS, BEES: Application should be made in the Jate evening when insects are at rest. Spray liberally into hiding and breeding places, especially under attic rafters, contacting as many

And Dreeding pasces, especially under attic finiters, connecting as many insects as possible. <u>FIEAS, BROWN DOS TICKS and LONE STAR TICKS</u>: On DOGS, CATS and PREMISES: Lightly spray mist enimats along back under forelegs on other areas, except scrotum and anus. Part hair to contact skin and insure good coverage. Hold sprayer 2 foot (30 cm) away while spraying. Apply until hair is thoroughly molet. Do not spray directly in oyes mouth or nose membranes. The remains the direct and prove the grave the primal harking. time then is tocoughly mode. Do not spag one cay in eyes mount or noise on animals. To prevent indications information years the animal bedding and other resting areas until the surface is slightly moist contacting as many insects as possible. Treat the floor areas, cracks and creakes in the floor and along baseboards at the rate of 2 seconds per linear foot (30 cm) at a distance of 12 to 15 fr. (30-45 cm). Also spray the entire surface of at a distance of 12 to 15 fr. (30-45 cm). Also spray the entire surface of doo houses.

#### INDOOR PEST CONTROL (Continued)

INDOOR PEST CONTROL (Continued) CARPET DEELISSANDLOTHS: To breat carpets - hold container 14 - 16 inches (36 - 46 cm) from carpet. Apply spray evenly, covering the entrie survoe until sightly moist. Spray underside of carpet when possible. Avoid wetting or susking, To treat gaments, bankets, carpets and other fabrics before storage. Dry clean or wash before and after storage. Apply spray evenly, covering the material on both sides until slightly moist. For best results, place treated materials in plastic bags after drying, and store in dark place. Also spray floor and baseboards, closets, cracks, cravles, infedied areas of abelound or when these invests or somethant infested areas of shelving or wherever these indects are seen or suspected.

SPACE SPRAY: FOR HOUSE FLIES, MOSQUITOES, CLUSTER FLIES, GNATS, ERUIT FLIES and FLYING MOTHS: Close off all doors and windows. Cover or remove all lood, food utensits, cover furniture or betting, Point sprayer upward. Direct spray mist to all parts of the room especially windows and other right sources which may attract these insects. Fill the room with mist then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the area when treatment is completed.

### OUTDOOR PEST CONTROL.

PESTS ON OUTSIDE SURFACES OF BUILDINGS: Use this precontrol ants, clover miles, crickets earwigs, millipedes, and sowbugs (i Common energy, down unlikes, bit deels behaviors, innegated estimation and ballongs (j. ) on outside curricates of buildings. To their prevent interstation of building, different currication of a height of 2 to 3, et (0.5 to 1 m), where pests are active and many find entrange. Also, apply as a residual story barry to utside surfaces of buildings, including porches, screens, window insmes, eaves, palios, parages, refuse dumns, and other areas where these pests congregale or have been seen. Repeat treatments as needed

ANTS, LONE STAR TICKS, CRICKETS, FLEAS AND EARWIGS: For bealthe statistic and the statistic of the second second in subject answer where there are woeds or brushy non-crop areas, spray infested areas thoroughly. For ants, theroughly wet hills and runways. Repeat application as intestation warrants and as reintestation posses.

HORNETS, YELLOW JACKETS, BEES, Application should be made in late evening when insects are at rest. Aim spray at nest openings in ground, bushes, and in cracks and crevices which may harbor nests, saturating nest openings and contacting as many insects as possible

TO CONTROL FIRE ANTS: Dilute 2 tablespoons (30 mi) of this product in 1 gai. (3.79 L) of water. Apply 1 gat (3.79 L) of solution as a gentle rain to each Fite Ant mound using a spinkler cash. Throughly wat the mound and sar-rounding area to a 4 lood dismetier (1.32 m). For best results, apply in cool weather, 65-800F 91 (18.51-270C), or in early moming or late evening hours. Treat new mounds as they appear, Do not distub the mound prior to basis. ment. If possible, treat all mounds in the vicinity, and treat all colonies which may not as yet have constructed a more retuint, and act at courses what may not as yet have constructed a mound. Larger quantities of solution can be prepared for multiple mound treatments. Equipment to deliver the dibuted product as a gentie rain at the rate of 1 gal (3.78 L) per mound is still required. Do not use equipment that produces a pressurized panty as pressurized sprays may disturb Fire Ants and cause migration, reducing effectiveness.

SPACE SPRAY: FOR HOUSE FLIES. MOSQUITOES, BITING FLIES, CLUS-TERFLIES, GNATS, FLYING MOTHS: For outdoor use as an aki in reducing annoyance from these insects, Syrary outside surfaces of screens, doors. window frames or wherever these insects may enter the room. Also treat surfaces around light fixiures on porches, in garages, and other placer these insects alight or congregate. Repeat as necessary.

