PYRONOX NO. 2 INSECTICIDE



FOR RAPID CONTROL OF CRAWLING AND FLYING INSECTS

Use in Food Processing Plants, Canneries, Packing Houses, Mills, Elevators, Granaries, Warehouses, Box Cars, Ships Holds, Breweries, Bottling Plants, Bakeries, Stores, Markets, Theaters, Hospitals, Schools, Hotels, Motels and Restaurants.

Active Ingredients:	
Pyrethrins	0.10%
*Piperonyl Butoxide, technical	
+ Inert Ingredients:	

*Equivalent to 0.8%(butykarbityl)(6-propylpiperonyl)ether and to 0.2% of related compounds + contains 98.90% Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN

5,30,90

CAUTION See side panel for additional precautionary statement

PRECAUTIONARY STATEMENT

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: HARMFUL IF SWALLOWED. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. THIS PRODUCT IS TOXIC TO FISH. KEEP OUT OF LAKES, STREAMS OR PONDS. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES.

FOODS SHOULD BE REMOVED OR COVERED DURING TREATMENT. ALL FOOD SURFACES SHOULD PE COVERED DURING TREATMENT OR THOROUGHLY CLEANED BEFORE USING. WHEN USING THE PRODUCT IN THESE AREAS, APPLY ONLY WHEN THE FACILITY IS NOT IN OPERATION. REMOVE ANT BIRDS, PETS OF FISH AQUARIUMS BEFORE SPRAYING. VACATE PREMISES AFTER TREATMENT AND VENTILATE BEFORE REOCCUPYING. SWEEP UP AND REMOVE DEAD INSECTS. REPEAT TREATMENT WHEN RE-INFESTATION OCCURS. CAUTION: PRE-TEST INSECTICIDE FOR POSSIBLE DISCOLORATION OR SPOTTING OF FABRIC, MATERIAL OR CARPET. TEST FOR POSSIBLE ALLERGIC REACTION.

DIRECTIONS FOR USE

IT IS A VIOLATION OF THE FEDERAL LAW TO USE THIS PRODUCT IN A MANNEP INCONSISTENT WITH ITS LABELING.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for the purpose.

SURFACE SPRAYING: To control exposed and accessible stages of ROACHES, ANTS, SPIDERS, CRICKETS, CHEESE MITES, GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEITLES, RUST RED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, SPIDER BEETLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, MEAL WORMS, GRAIN MITES and CADELLES: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface. To control SILVER FISH, apply spray thoroughly to cracks and crevices throughout the entire building, including the attic. Treat inside of book crases, dresser drawers, cabinets and closets. To control CLOVER MITES indoors, spray floor, baseboards, around windows and doors and other possible points of entry. To prevent reinfestation, spray the outside of the foundation and lower walls. Do not spray growing plants with this product. To control FLEAS indoors, spray infested areas thoroughly with particular attention to cracks, crevices, behind baseboards and moldings and under edges of rugs. Pet bedding should be thoroughly cleaned or destroyed and replaced. Pets should be treated using an approved insecticide to prevent reinfestation. To control BROWN DOG TICKS indoors, apply spray carefully to all observable crevices, behind baseboards, pictures, furniture and draperies and under faces of rugs. Dogs should be treated with an approved insecticide to reduce infestation. To control CARPET BEETLES: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets, and infested areas of shelving. Reprat treatment as needed. To control BEDBUGS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and

REPEAT TREATMENT WHEN RE-INFESTATION OCCURS.

SPACE SPRAYING: Close doors and windows and shut of ventilation systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of FLIES, FRUIT FLIES, GNATS, MOSOUITOES, SKIPPER FLIES, WASPS, HORNFTS, ANGOUMOIS GRAIN MOTHS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, AND TOBACCO MOTHS: directosproy as an upward angle, distributing it uniformly throughout the entire area, at a rate of one gallon to 100,000 cu ft of space

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

FPA REG_NO_45385-73

NET CONTENT:

Not Reviewed. Registrant claims to be in accordance with Spregistration of Posticide Product Guidance for Phase 2 For course, Page 2.12, Active to Inert Change in Status.

EPA EST NO 45385 IL-01 CHEM 177 INC BARRINGTON, 15: 60019 ORDFR CODE 0-321