# **CENOL MILL SPRAY** WITH SBP-1382



A synthetic Pyrethroid for controlling the following flying and crawling insects: HOUSE FLIES, MOSQUITOES, GNATS, FLYING MOTHS, (MILLERS), FRUIT FLIES, VINEGAR FLIES, WASPS, HORNETS, YELLOW JACKETS, COCKROACHES, WATERBUGS, BEDBUGS, FLEAS, TICKS, ANTS, CRICKETS, SILVERFISH, EARWIGS, SPIDERS, GRAIN INSECTS (ADULTS OF CONFUSED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, SHINY SPIDER BEETLES, GRAIN WEEVILS, ANGOUMOIS GRAIN MOTHS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, GRAIN MITES). CARPET BEETLES, SOWBUGS, CENTIPEDES, MILLIPEDES AND CIGARETTE BEETLES.

ACTIVE INGREDIENTS:	
* + < Resmethrin	0.50%
"INERT INGREDIENTS:	99.50%
TOTAL:	100.00 w/w

- \*\* contains aliphatic petroleum distillate
- \*Cis/trans isomers ratio: max. 30%(+)cis and min. 70%(+)trans. +U.S. patent nos. 3,465,007 ar.d 3,542,928
- < Roussel-Bio's SBP-1382 Brand of Resmethrin Insecticide

## KEEP OUT OF REACH OF CHILDREN CAUTION See side panel for additional precautionary statement

#### STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Call a physician or Poison Control Center immediately. IF INHALED: Remove to fresh air. 1F ON SKIN: Remove contaminated clothing and wash affected areas with soap and water. 1F IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists. NOTE TO PHYSICIAN: Contains petroleum distillate. Vomiting may cause aspiration pneumonia.

### PRECAUTIONARY STATEMENT

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Avoid contamination with skin, eyes or ciothing. Avoid breathing mist. Cover or remove exposed foods. Avoid contamination of cooking utensils and food preparation surfaces. Remove pets and cover fish aquariems before spraying.

#### ENVIRONMENTAL HAZARDS

This formulation it highly toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from the ated sites may be hazardous to fish in adjacent waters. Do not spray on turf or plants. Do not contaminate water when disporing of equipment washwaters.

#### PHYSICAL AND CHEMICAL HAZARDS

Extinguish open flame before space spraying. This formulation is combustible. Do not use or store near heat or open flame. CAUTION: PRE-TEST INSECTICIDE FOR POSSIBLE DISCOLORATION OR SPOTTING OF FABRIC, MATERIAL OR CARPET. TEST FOR POSSIBLE ALLERGIC REACTION.

LABELING.

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. STORAGE: Store product in original container in a locked storage area. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: 1 Gal and under wrap container in several layers of news paper and discard in trash. Over 1 Gal. Triple rinse (or equivalent) and offer for receding, reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, or burning. If burned, stay out of smoke.

For use indoors and outside premises of HOTELS, MOTELS, HOMES, FOOD HANDLING ESTABLISHMENTS, PACKAGING AND PROCESSING FLANTS, SEED HOUSES, COMMERCIAL, INDUSTRIAL, OFFICE, INSTITUTIONAL and STORAGE BUILDINGS, MILK-HANDLING AREAS OF MILK STORAGE ROOMS, EGGPRODUCING PLANTS, SUPERMARKETS, RESTAURANTS, BOTTLING PLANTS, CANNERIES, FLOUR and FEED MILLS, BAKERIES, BOATS, SHIP CABINS and HOSPITALS. In HOSPITALS. Remove patients prior to dream ent Ventilate rooms for two hours before returning patients. Also for use in BUSES, AUTOMOBILES, TRUCKS, BURCARS and SHIP CARGO HOLDS. Also for use in DOG KENNELS and HORSE STABLES. In HORSE STABLES STABLES AND INCOME. application if horses will be used for food



Foods should be removed or covert during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. After space spraying in food processing plants, wash all equipment, benches, shelving etc. where exposed food will be handled thoroughly with an effective cleaning compound rinse with potable water. When using the

product in these areas, apply only when facility is not in operation.

INDOORS: SPACE SPRAY: To kill HOUSE FLIES, MOSQUITOES, GNATS, FLYING MOTHS (MILLERS), FRUIT FLIES, VINEGAR FLIES. Close room or area to be treated. Shut off air conditioner and ventilator system. Seal all open vents. Remove fish and pets. Apply undiluted as a very fine mist space spray through manually operated equipment or with electrical and other powered applicators. Avoid staining by holding the spray nozzle at least five feet from objects. Direct the fine spray mist toward the upper corner of the room. Fill the room with a fine mist space spray at the rate of 3 fl.oz per 5,000 cu.ft (89 ml per 142 cu meters). Leave room closed for one-half hour. Ventilate thoroughly before re-entry. Sweep up and destroy insects. CONTACT SPRAY: To kill COCKROACHES, WATERBUGS, SILVERFISH, FLEAS, EARWIGS, SOWBUGS, CARPET BEETLES, CENTIPEDES, MILLIPEDES. Adjust to produce large, wetting type spray particles. Direct spray to contact these insects in cracks and crevices in floors, baseboards, around pipes, behind cabinets, and other harborages. Spray these infested areas until thoroughly wet. Repeat as necessary. To kill SPIDERS: Spray directly on these pests and use repeated applications to the webbing and all cracks and crevices which harbor spiders. To kill ANTS: Spray doors, around window frames, ant trails and hills. To kill BEDBUGS: Apply as a spot treatment to cracks and crevices around baseboards, floorboards, bedboards and walls. First thoroughly clean and air mattresses and springs. Then spot treat only areas that may harbor bedbugs, concentrating on tuffs, folds and seams. Repeat as necessary. For persistent infestations, consult professional applicator. To kill FLEAS and TICKS: Spray directly on insects as well as floors, floor covering, baseboards, cracks and crevices, infested furniture and animal resting a 11s. Hold spray nozzle about 18 inches (46 cm) from areas to be treated. Apply spray until area is moist. Treat pet bedding the some way. Also spray entire inside surface of dog houses. Repeat as necessary. Do not allow pets to contact treated areas until spray has dried. Remove or cover animal feed before spraying. Also treat dogs, cats and horses with a registered flee-control product before they re-enter the treated areas. To kill GRAIN INSECTS: ADULTS of CONFUSED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLE, SHINY SPIDER BEETLES, GRAIN WEEVILS, CIGARETTE BEETLES, ANGOUMOIS GRAIN MOTHS, MEDITERRANEAN FLOUR MOTHS, INDIAN MEAL MOTHS, GRAIN MITES. Remove infested grain, cereal or flour and destroy. Avoid contamination of other food and foodstuffs in the areas by removing before spraying. Spray directly on exposed stages and into cracks and crevices where grain insects may be hiding. In homes, dispose of all infested products. The shelves and drawers should be cleaned with shelf paper after allowing shelves to dry. Any utensils that are contacted with spray should be washed before using. Repeat as necessary. To kill WASPS, HORNETS, YELLOW JACKETS: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places contacting as many insects as possible. Repeat as necessary.

OUTDOORS: To control nuisance pests around buildings, patios and paved areas: Apply as a contact spray. Spray directly on insects. Avoid spraying turf or plants. To kill FLEAS and TICKS: Spray directly on insects as well as floors, floor covering, baseboards, cracks and crevices, infested furniture and animal resting areas. Hold spray nozzle about 18 inches (46 cm) from areas to be treated. Apply spray until area is moist. Treat pet bedding the same way. Also spray entire inside surface of dog houses. Repeat as necessary. Do not allow pets to contact areas until spray has dried. Remove or cover animal feed before spraying. Also, treat dogs, cats and horses with a registered flea-control product before they re-enter the treated area. To kill ANTS: Spray around window frames, and trails and hills. To kill WASPS, HORNETS, YELLOW JACKETS: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places contacting as many insects as possible. Aim spray at nest openings in bushes, under eaves, chimneys, cracks and crevices to saturation in and around nest openings. Contact as many insects as possible. Repeat as necessary. To kill EARWIGS, SPIDERS, CENTIPEDES, and MILLIPEDES: Apply as a contact spray, as for indoor use. To control named flying and crawling insects around premises, spray thoroughly around doors, window frames and similar areas of entry. Also, thoroughly spray garbage and refuse areas,

especially around poultry houses, barns, and similar areas of heavy infestation. Do not spray on turf or plants.

THERMAL APPLICATION: For control of HOUSEFLIES and MOSQUITOES. This formulation is to be used undiluted in

only such portable thermal applicator equipment that can deliver spray particles sizes within the aerosol size range.

INDOORS: For control of HOUSEFLIES, MOSQUITOES and GNATS in warehouses, restaurants and industrial institutions. Remove or cover exposed food handling surfaces. Remove animals and plant life. Shut off ventilation systems, air ducts, air-conditioning equipment and close off room or area to be treated. Do not spray while food processing plant is occupied or in operation. Excinguish all flames and pilot lights. Apply as a dry fog at the rate of 1 1/3 fl o. per 1,000 cu ft. Direct nozzic of thermal applicator toward upper corners of the room and along ceiling. Close off room for at least one-half nour. Do not remain in treated area after treatment. Ventilate thoroughly before re-entry. OUTDOORS: For reduction in annoyance from HOUSEFLIES, MOSQUITOES and GNATS in yards, patios, around swimming pools, picnic areas, camp sites, drive-ins and horse stables, and other small area applications. Hold thermal applicator at least 5 feet from any object, especially plants and shrubbery. Avoid wetting any plant foliage. Apply undiluted at the rate of 3 3/4 gallons per acre (35 l per ha). Fog only when there is no breeze or the wind velocity is less than 3 miles per hour. The ideal time is generally at dusk when wind dies down and the air temperature and ground temperature are approximately the same. Under these conditions the fog will linger in the treated area for a longer time. Apply fog while moving away from treated area. Avoid inhalation of the fog. Do not enter the treated areas until the fog has dissipated. Repeat application as necessary.

WARRANTY STATEMENT: Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed on implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such or in combination with other materials.

EPA REG. NO. 45385-78

EPA FST NO 45385 II -01

CHEM-TOX, INC BARRINGTON, IL. 60010 ORDER CODE, O-335

**NET CONTENT:** 

