

45385-99

05/01/2007

1/4



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

MAY 1 2007

H. R. McLane, Ph.D.
H.R. McLane, Inc.
7210 Red Road, Suite 206
Miami, Florida 33143-5321

Subject: Label Amendment to Revise Directions for Use

CENOL, 0.5% Multipurpose Insecticide
EPA Reg. No. 45385-99
Your Submission Dated August 22, 2006


Dear Dr. McLane:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comments listed below. Two (2) copies of the finished labeling must be submitted prior to releasing the product for shipment. A stamped copy of the label is enclosed for your records.

1. Move the statements beginning with "*Avoid contamination...*" and ending "*...until surfaces are dry*" from the precautionary statements to the directions for use section of the label.
2. Revise "*...pick up of insects such as bedbugs*" to read "*...pick up of listed insects such as bedbugs*" or "*...pick up of bedbugs and other listed insects*".
3. Add "*and other aquatic organisms*" to the first sentence under Environmental Hazards.
4. Move the sentence "*Do not contaminate water, food, or feed by storage and disposal*" directly under the subheading "Storage and Disposal".
5. Delete: "*Repeat treatment as needed*" throughout or identify a specific repeat interval based upon efficacy data of the product.

If you have any questions regarding this action, please contact Olga Odiott of my team at (703) 308-9369.

Sincerely,



George T. LaRocca
Product Manager (13)
Insecticide Branch
Registration Division (7505P)

Enclosure

3/4

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before space spraying or surface applications. Do not use on humans. Vacate room after treatment and ventilate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry.

FIRST AID

IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment.

HOT LINE: You may also call 1-800-858-7378 for emergency medical treatment advice.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high water mark. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

Do not use this product in or on electrical equipment due to the possibility of shock hazard. Do not allow product to freeze.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area out of reach of children and pets. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.

PRODUCT DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the container be used, carefully following label directions and precautions.

CONTAINER DISPOSAL

IF EMPTY: Do not reuse this container. Place in trash or offer for recycling if available. For containers larger than 1 gallon: Triple rinse, or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned stay out of smoke.

IF PARTLY FILLED: Call your local waste agency or 1-800-253-2687 for disposal instructions. Never place unused product down any indoor or outdoor drain. **SPILL:** In case of spill, absorb liquid with absorbent material such as soil or clay etc. sweep up and dispose of material according to **PRODUCT DISPOSAL** above.

For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents). Call the National Pesticide Telecommunications at 1-800-858-7378.

NOTICE: Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith. 08-21-08

CHEMotox™

**CENOL 0.5%
MULTIPURPOSE INSECTICIDE
FOR USE BY HOMEOWNERS, PEST CONTROL
OPERATORS OR COMMERCIAL APPLICATORS**

A READY-TO-USE PERMETHRIN STABLE EMULSION/INSECTICIDE TO KILL INSECTS AS LISTED FOR OUTDOORS, IN HOMES, AND NON-FOOD AREAS OF FOOD/FEED PROCESSING ESTABLISHMENTS, KENNELS, COMMERCIAL BUILDINGS, WAREHOUSES, THEATERS, OFFICE BUILDINGS, SCHOOLS, MOTELS AND HOTELS. • PROTECTS TRAVEL LUGGAGE FROM "PICK UP" OF INSECTS SUCH AS BEDBUGS.

• FOR KILL AND CONTROL OF FLEAS, BROWN DOG TICKS, AND LICE ON PREMISES AND • ON DOGS. FOR CONTROL OF TICKS WHICH MAY CAUSE LYME DISEASE; • CONTROLS FLEAS AND TICKS ON DOGS FOR 14 DAYS. • FOR CONTROL OF PESTS ON LIVESTOCK.

ACTIVE INGREDIENT:

Permethrin [CAS 52645-53-1]..... 0.50%

OTHER INGREDIENTS: 99.50%

TOTAL 100.00%

EPA REG. NO. 45385-99 EPA EST. NO. 45385-OH-1

**CAUTION
KEEP OUT OF REACH OF CHILDREN**
See side panels for additional Precautionary Statements.

**ACCEPTED
with COMMENTS
In EPA Letter Dated
MAY 1 2007**
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
45385-99

Net Contents: 6 Oz. - 8Oz. - 12 Oz. - 1 Qt. - 1 Gal.



**CENOL
INCORPORATED
P.O. Box 472
Twinsburg, OH 44087**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds. "SHAKE WELL BEFORE USING"

Ready-to-use and apply for most uses with the attached sprayer.

For travelers who want to protect their luggage from "picking up" insects such as bedbugs, spray the outer and inner surfaces of luggage with a light spray mist of this product. Allow the luggage to completely dry before use.

INDOOR APPLICATION

Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food / feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed. Cover items that might water stain. Nonfood/feed areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling). Vacate treated areas and ventilate before reoccupying. For maximum effectiveness, a combination of spot, Surface treatment & fogging is recommended. Do not apply to classrooms while in use.

SURFACE SPRAYING

TO KILL and CONTROL COCKROACHES, WATERBUGS, PALMETTO BUGS, SCORPIONS, MILLPEDES, CENTIPEDES, SOWBUGS, PILLBUGS, ANTS, SILVERFISH, FIRE BRATS, SPIDERS, CRICKETS, CLOVER MITES, CHEESE MITES, GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, RUST RED FLOUR BEETLES, DRUGSTORE BEETLES, MEAL WORMS, GRAIN MITES and CADELLES: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry.

TO CONTROL CARPET BEETLES: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

TO CONTROL BED BUGS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floor. Repeat treatment as needed. Allow all treated articles to thoroughly dry before use.

TO CONTROL FLEAS (ADULT AND LARVAE) AND TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (ADULTS AND LARVAE): Thoroughly spray infested areas, pet beds, resting quarters, nearby, cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas.

TO CONTROL CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly, and air for several hours in sun light, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining. Allow all treated articles to dry thoroughly before use.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

SPACE SPRAYING

Close all doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of House Flies, Fruit Flies, Gnats, Mosquitoes, Skipper Flies, Wasps, Hornets, Bees, Black Flies, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moths and Tobacco Moths: Direct spray at an upward angle, distributing it uniformly throughout the entire area at a rate of 1 ounce per 1,000 cubic feet of space. Keep area closed for 2 hours. Ventilate before reoccupying.

INDOOR APPLICATION continued:

INDOOR FOGGING: To kill exposed or accessible stages of the above named* insects, use a rate of 1 quart to 1 gallon for 10,000 to 40,000 cubic feet of space. Contact insects directly when possible. Keep area closed for 2 hours. Thoroughly ventilate before reoccupying.

PEST CONTROL ON LIVESTOCK

To protect cattle (beef and dairy), goats, sheep, hogs and horses from; Horn Flies, Face Flies, House Flies, Mosquitoes and Gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

TO KILL and CONTROL: **STABLE FLIES; HORSE FLIES AND DEER FLIES:** Apply at rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

TO CONTROL BLOOD SUCKING LICE: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

TO CONTROL POULTRY LICE: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

TO CONTROL BEDBUGS AND MITES IN POULTRY HOUSES: Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

ON DOGS: To kill and control Fleas, Ticks, and Lice: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Use protective gloves or mitts when applying product. Repeat application every two weeks, if necessary.

Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant, or nursing animals. Do not use on dogs under 12 weeks of age.

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. **DO NOT USE ON CATS OR KITTENS.**

OUTDOOR APPLICATION

TO CONTROL TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (**PERIMETER TREATMENT**): Use Cenol 0.5% Multipurpose Insecticide as a residual treatment to control Ticks including the Deer Tick (*Ixodes dammini*), Lone Star Tick, (*Amblyomma americanum*), Brown Dog Tick (*Rhipicephalus sanguineus*), American Dog Tick (*Demacenter variabilis*) and Western Black-legged Tick (*Ixodes pacificus*).

For each area listed below, apply until surface is wet (1/2 to 1 gallon for 750-1,000 sq.ft)

Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways, and in crawl spaces where these pests may enter. Treat a band of vegetation and leaf litter 6 to 10 feet wide around the outside of buildings to help control ticks occurring there.

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

FOR CONTROL OF FLOWER GARDEN INSECTS

For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray as well as White Flies, Aphids. Exposed Thrips, Spiders, Red Mites, and Leafminers. May be used on plants as a multipurpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, euonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

EFFECTIVE WHEN USED AS DIRECTED AGAINST WHITEFLIES, APHIDS, SPIDER AND RED MITES, EXPOSED THRIPS, LACE BUGS, JAPANESE BEETLES, PAVEMENT ANTS, CLOVER MITES AND ARMYWORMS When spraying plants, do not operate closer than 18 inches. Use a sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

Dossier Page; 4 of 4