

F. F. F.

LAWN CARE PRODUCTS

ST. AUGUSTINE WEED AND FEED  
FERTILIZER + IRON

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

In case of contact: Eyes, flush with water for 15 minutes and get medical attention if irritation persists.

Skin, wash with plenty of soap and water. See back panel for additional precautions.

ACTIVE INGREDIENTS:

\*Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine)--.75%

Inert ingredients-----99.25%

TOTAL-----100.00%

\*Contains 15.01 LBS of Atrazine per ton.

ACCEPTED

DEC - 3 1982

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 45537-1

Net Wt.

Manufactured by  
FFF LAWN CARE PRODUCTS  
MT. DORA, FLA. 32757

EPA EST NO: 45537-FL-1

(FRONT PANEL)

GUARANTEED ANALYSIS

Controls Such Weeds As:

GERMINATING: Crabgrass, Barnyardgrass, Wiregrass, Paspalum Seedlings(Dallisgrass, Vaseygrass, Bahiagrass). Sedges(such as Cyperus Compressus), and Spurge.  
ESTABLISHED AND GERMINATING: Betony, Blackmedic(yellow Trefoil) Burclover, Chickweed, Dandelion, Oxalis, Florida Pusley, Pennywort(Dollarweed) and Creeping Charlie(Matchweed).

PRECAUTIONARY STATEMENTS:

Hazard to Humans and Domestic Animals.

CAUTION:

Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.  
KEEP OUT OF REACH OF CHILDREN. Do not take internally.

ENVIRONMENTAL HAZARD:

~~Keep out of lakes, streams or ponds.~~  
Avoid application of the root zones of shrubs and trees.  
Avoid application on flower and garden plants.  
Delay Spring plantings for one month after application.  
Clean spreader thoroughly after using.  
Do not use on lawns of Bahia grass, Bermuda Grass, Dichondra, Bentgrass, or carpet grass or where clover is present or desired.  
Apply only in Winter, Spring and Fall. This product is not recommended for use during July, August and September.  
DO NOT OVERAPPLY.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product inconsistent with its labeling.  
Apply between Jan. 1 and March 31 in Spring and after September in the fall at the rate of 32 Lbs. on 5,000 Sq. Ft.  
In Florida and other Southern states, applications may be made after March 31.  
For best results apply before the weeds come up.  
Established weeds are harder to control and may require additional applications in the Spring and Fall. WATER IN THOROUGHLY.

SPREADER SETTING:

Cyclone----- 5      Scotts----- 8½  
Note: Spreader rates vary with walking speed. Check your spreader over a known area.

STORAGE AND DISPOSAL

STORAGE: store in a dry place. Do not store near growing plants or fertilizers, seeds, insecticides or fungicides.  
Keep away from feed or foodstuffs.

DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

(BACK PANEL)