

F.F.F. LAWN CARE PRODUCTS, INC.

TREATS 5,000 SQ. FT.
WEED & FEED
(With Atrazine)
25-4-6
Controls Weeds While
Feeding Your Lawn With A
Quick Greening Long-Lasting
Fertilizer.

FOR USE ON ST. AUGUSTINE, ZOYSIA,
CENTIPEDE AND CARPETGRASS LAWNS ONLY

KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS

Avoid contact with skin, eyes, or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS:

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Not for use on or under trees, shrubs, bedding plants, flowers or garden plants.

ACTIVE INGREDIENT:


Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine).. 1.16%

INERT INGREDIENTS:..... 98.84%

23.70 POUNDS ATRAZINE TOTAL: 100.00%

EPA Reg. No

EPA EST. NO. 45537-FL-1



Picture

KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS

Avoid contact with skin, eyes, or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS:

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Not for use on or under trees, shrubs, bedding plants, flowers or garden plants.

ACTIVE INGREDIENT:

Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine).. 1.16%

INERT INGREDIENTS:..... 98.84%

23.20 POUNDS ATRAZINE TOTAL: 100.00%

EPA Reg. No

EPA EST. NO. 45537-FL-1

ACCEPTED
FEB - 4 1986
Under the Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 45537-4

PICURE

NET WT: 20 LBS. (9.07 kg)

F.F.F. LAWN CARE PRODUCTS, INC.

WEED & FEED
(With Atrazine)

GUARANTEED ANALYSIS 25-4-6

Total Nitrogen (N).....	25%
1.6% Ammoniacal Nitrogen	
22.7% Urea Nitrogen*	
0.7% Water Insoluble Nitrogen	
Available Phosphoric Acid (P ₂ O ₅).....	4%
Soluble Potash (K ₂ O).....	6%
Boron (B).....	0.02%
Copper (Cu).....	0.05%
Iron (Fe).....	3.00%
Manganese (Mn).....	0.05%
Molybdenum (Mo).....	0.0005%
Zinc (Zn).....	0.05%

Primary Plant Food Sources: Ureaformaldehyde, Sulfur Coated Urea, Urea, Diammonium Phosphate and Muriate of Potash.

Secondary Plant Food Sources: Fertilizer Grade Borate, Oxides of Copper, Oxides of Iron, Oxides of Manganese, Fertilizer Grade Molybdate, and Oxides of Zinc.

*Part of the nitrogen in this product has been sulfur coated to provide 5.3% coated slow release nitrogen.

Chlorine (Cl), not more than 4.8%.

WHAT THIS PRODUCT DOES

F.F.F. Weed & Feed W/Atrazine controls growing weeds such as dandelions, black medic, buckhorn, burclover, Carolina geranium, chickweed, purslane, Florida betony, Florida pusley, rabbit-tobacco, henbit, matchweed, oxalis, red sorrel, rescuegrass, shepherdspurse, wild mustard, and many others. This product also contains a high quality, slow-release fertilizer. This product will green your lawn in a few days and will continue feeding for 3-4 months.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Apply in January, February, or early March to prevent germination of crabgrass, bull paspalum, sedges (from seed), and spotted spurge.

Apply anytime broadleaf weeds are growing to get control.

Apply in the fall to control henbit, chickweed and other winter weeds.

If complete control is not obtained, a second application may be necessary 2-3 months after the first. For best results apply with a fertilizer spreader at the settings listed below. Apply to newly sprigged St. Augustine or Zoysia lawns to help prevent competition from other weeds. Apply when grass blades are dry. On hot days (above 80°) water the lawn to rinse particles off the grass.

SPREADER SETTINGS

Weed & Feed With Atrazine 25-4-6 5,000 Sq. Ft.	Scott's Drop (PF1) 5.0	Scott's Rotary (C2) 1	Cyclone (B1) 3.75	Sta- Green (EB84) 6.0	Ortho Rotary (#000) 3.0	Ortho Drop (3000) 4.0	Sears Drop (671,19150) 6.0	Penn- ington (EB84) 6.0
---	---------------------------------	--------------------------------	-------------------------	--------------------------------	----------------------------------	--------------------------------	-------------------------------------	----------------------------------

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product

F.F.F. LAWN CARE PRODUCTS, INC.

WEED & FEED
(With Atrazine)

GUARANTEED ANALYSIS 25-4-6

Total Nitrogen (N).....	25%
1.6% Ammoniacal Nitrogen	
22.7% Urea Nitrogen*	
0.7% Water Insoluble Nitrogen	
Available Phosphoric Acid (P ₂ O ₅).....	4%
Soluble Potash (K ₂ O).....	6%
Boron (B).....	0.02%
Copper (Cu).....	0.05%
Iron (Fe).....	3.00%
Manganese (Mn).....	0.05%
Molybdenum (Mo).....	0.0005%
Zinc (Zn).....	0.05%

Primary Plant Food Sources: Ureaformaldehyde, Sulfur Coated Urea, Urea, Diammonium Phosphate and Muriate of Potash.

Secondary Plant Food Sources: Fertilizer Grade Borate, Oxides of Copper, Oxides of Iron, Oxides of Manganese, Fertilizer Grade Molybdate, and Oxides of Zinc.

*Part of the nitrogen in this product has been sulfur coated to provide 5.3% coated slow release nitrogen.

Chlorine (Cl), not more than 4.8%.

WHAT THIS PRODUCT DOES

F.F.F. Weed & Feed W/Atrazine controls growing weeds such as dandelions, black medic, buckhorn, burclover, Carolina geranium, chickweed, purslane, Florida betony, Florida pusley, rabbit-tobacco, henbit, matchweed, oxalis, red sorrel, rescuegrass, shepherdspurse, wild mustard, and many others. This product also contains a high quality, slow-release fertilizer. This product will green your lawn in a few days and will continue feeding for 3-4 months.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Apply in January, February, or early March to prevent germination of crabgrass, bull paspalum, sedges (from seed), and spotted spurge.

Apply anytime broadleaf weeds are growing to get control.

Apply in the fall to control henbit, chickweed and other winter weeds.

If complete control is not obtained, a second application may be necessary 2-3 months after the first. For best results apply with a fertilizer spreader at the settings listed below. Apply to newly sprigged St. Augustine or Zoysia lawns to help prevent competition from other weeds. Apply when grass blades are dry. On hot days (above 80°) water the lawn to rinse particles off the grass.

SPREADER SETTINGS

Weed & Feed With Atrazine 25-4-6 5,000 Sq. Ft.	Scott's Drop (PF1) 5.0	Scott's Rotary (C2) 1	Cyclone (H1) 3.75	Sta- Green (EB84) 6.0	Ortho Rotary (4000) 3.0	Ortho Drop (3000) 4.0	Sears Drop (671, 19150) 6.0	Penn- ington (EB84) 6.0
---	---------------------------------	--------------------------------	-------------------------	--------------------------------	----------------------------------	--------------------------------	--------------------------------------	----------------------------------

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product

Copper (Cu)	0.05%
Iron (Fe)	3.00%
Manganese (Mn)	0.05%
Molybdenum (Mo)	0.0005%
Zinc (Zn)	0.05%

Primary Plant Food Sources: Ureaformaldehyde, Sulfur Coated Urea, Urea, Diammonium Phosphate and Muriate of Potash.

Secondary Plant Food Sources: Fertilizer Grade Borate, Oxides of Copper, Oxides of Iron, Oxides of Manganese, Fertilizer Grade Molybdate, and Oxides of Zinc.

*Part of the nitrogen in this product has been sulfur coated to provide 5.3% coated slow release nitrogen.

Chlorine (Cl), not more than 4.8%.

WHAT THIS PRODUCT DOES

F.F.F. Weed & Feed W/Atrazine controls growing weeds such as dandelions, black medic, buckhorn, burclover, Carolina geranium, chickweed, purslane, Florida betony, Florida pusley, rabbit-tobacco, henbit, matchweed, oxalis, red sorrel, rescuegrass, shepherdspurse, wild mustard, and many others. This product also contains a high quality, slow-release fertilizer. This product will green your lawn in a few days and will continue feeding for 3-4 months.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Apply in January, February, or early March to prevent germination of crabgrass, bull paspalum, sedges (from seed), and spotted spurge.

Apply anytime broadleaf weeds are growing to get control.

Apply in the fall to control henbit, chickweed and other winter weeds.

If complete control is not obtained, a second application may be necessary 2-3 months after the first. For best results apply with a fertilizer spreader at the settings listed below. Apply to newly sprigged St. Augustine or Zoysia lawns to help prevent competition from other weeds. Apply when grass blades are dry. On hot days (above 80°) water the lawn to rinse particles off the grass.

SPREADER SETTINGS

Weed & Feed With Atrazine	Scott's Drop	Scott's Rotary	Cyclone	Sta-Green	Ortho Rotary	Ortho Drop	Sears Drop	Penn-ington
25-4-6	(PF1)	(C2)	(H1)	(EB84)	(4000)	(3000)	(671,19150)	(EB84)
5,000 Sq. Ft.	5.0	1	3.75	6.0	3.0	4.0	6.0	6.0

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides, or seeds.

PRODUCT DISPOSAL: To dispose of bags which contain unused products, securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

PRECAUTIONS:

Do not apply under the branches (root zone area) of shrubs, trees, flowers, etc. Do not apply to newly seeded Zoysia, centipede, or carpetgrass lawns. Wait until they have required at least 2 mowings. Don't apply to flowers, bedding plants, ornamental trees or shrubs.

CAUTION: We recommend that you keep this product (and all fertilizers) out of the reach of children. Avoid getting on concrete walks or drives, swimming pools, stucco or asbestos siding as the combination of certain fertilizer elements can cause permanent stains. Keep in a dry place.

BACK PANEL

F.F.F. LAWN CARE PRODUCTS, INC.

TREATS 5,000 SQ. FT.

WEED & FEED

(With Atrazine)

25-4-6

Controls Weeds While
Feeding Your Lawn With A
Quick Greening Long-Lasting
Fertilizer.

FOR USE ON ST. AUGUSTINE, ZOYSIA,
CENTIPEDE AND CARPETGRASS LAWNS ONLY

KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS

Avoid contact with skin, eyes, or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS:

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Not for use on or under trees, shrubs, bedding plants, flowers or garden plants.

ACTIVE INGREDIENT:

Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine).. 1.16%

INERT INGREDIENTS:..... 98.84%

23.20 POUNDS ATRAZINE (1.16%) TOTAL: 100.00%

EPA Reg. No

EPA EST. NO. 45537-FL-1

ACCEPTED
FEB - 4 1986
Used as Insecticide.
Fungicide, Herbicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 45537-14

Picture

CUTION

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

Avoid contact with skin, eyes, or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS:

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Not for use on or under trees, shrubs, bedding plants, flowers or garden plants.

ACTIVE INGREDIENT:


Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine).. 1.16%

INERT INGREDIENTS:..... 98.84%

23.20 POUNDS ATRAZINE TOTAL: 100.00%

EPA Reg. No

EPA EST. NO. 45537-FL-1



Picture

NET WT: 20 LBS. (9.07 kg)

EPA REG. NO.

F.F.F. LAWN CARE PRODUCTS, INC.

WEED & FEED
(With Atrazine)

GUARANTEED ANALYSIS 25-4-6

Total Nitrogen (N).....	25%
1.6% Ammoniacal Nitrogen	
22.7% Urea Nitrogen*	
0.7% Water Insoluble Nitrogen	
Available Phosphoric Acid (P ₂ O ₅).....	4%
Soluble Potash (K ₂ O).....	6%
Boron (B).....	0.02%
Copper (Cu).....	0.05%
Iron (Fe).....	3.00%
Manganese (Mn).....	0.05%
Molybdenum (Mo).....	0.0005%
Zinc (Zn).....	0.05%

Primary Plant Food Sources: Ureaformaldehyde, Sulfur Coated Urea, Urea, Diammonium Phosphate and Muriate of Potash.

Secondary Plant Food Sources: Fertilizer Grade Borate, Oxides of Copper, Oxides of Iron, Oxides of Manganese, Fertilizer Grade Molybdate, and Oxides of Zinc.

*Part of the nitrogen in this product has been sulfur coated to provide 5.3% coated slow release nitrogen.

Chlorine (Cl), not more than 4.8%.

WHAT THIS PRODUCT DOES

F.F.F. Weed & Feed W/Atrazine controls growing weeds such as dandelions, black medic, buckhorn, burclover, Carolina geranium, chickweed, purslane, Florida betony, Florida pusley, rabbit-tobacco, henbit, matchweed, oxalis, red sorrel, rescuegrass, shepherdspurse, wild mustard, and many others. This product also contains a high quality, slow-release fertilizer. This product will green your lawn in a few days and will continue feeding for 3-4 months.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Apply in January, February, or early March to prevent germination of crabgrass, bull paspalum, sedges (from seed), and spotted spurge.

Apply anytime broadleaf weeds are growing to get control.

Apply in the fall to control henbit, chickweed and other winter weeds.

If complete control is not obtained, a second application may be necessary 2-3 months after the first. For best results apply with a fertilizer spreader at the settings listed below. Apply to newly sprigged St. Augustine or Zoysia lawns to help prevent competition from other weeds. Apply when grass blades are dry. On hot days (above 80°) water the lawn to rinse particles off the grass.

SPREADER SETTINGS

Weed & Feed With Atrazine 25-4-6 5,000 Sq. Ft.	Scott's Drop (PF1) 5.0	Scott's Rotary (C2) 1	Cyclone (H1) 1.75	Sta- Green (EB84) 6.0	Ortho Rotary (4000) 3.0	Ortho Drop (3000) 4.0	Sears Drop (671, 9150) 6.0	Penn- ington (EB84) 6.0
---	---------------------------------	--------------------------------	-------------------------	--------------------------------	----------------------------------	--------------------------------	-------------------------------------	----------------------------------

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product

Boron (B).....	0.05%
Copper (Cu).....	0.05%
Iron (Fe).....	3.00%
Manganese (Mn).....	0.05%
Molybdenum (Mo).....	0.0005%
Zinc (Zn).....	0.05%

Primary Plant Food Sources: Ureaformaldehyde, Sulfur Coated Urea, Urea, Diammonium Phosphate and Muriate of Potash.

Secondary Plant Food Sources: Fertilizer Grade Borate, Oxides of Copper, Oxides of Iron, Oxides of Manganese, Fertilizer Grade Molybdate, and Oxides of Zinc.

*Part of the nitrogen in this product has been sulfur coated to provide 5.3% coated slow release nitrogen.

Chlorine (Cl), not more than 4.8%.

WHAT THIS PRODUCT DOES

F.F.F. Weed & Feed W/Atrazine controls growing weeds such as dandelions, black medic, buckhorn, burclover, Carolina geranium, chickweed, purslane, Florida betony, Florida pusley, rabbit-tobacco, henbit, matchweed, oxalis, red sorrel, rescuegrass, shepherdspurse, wild mustard, and many others. This product also contains a high quality, slow-release fertilizer. This product will green your lawn in a few days and will continue feeding for 3-4 months.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply in January, February, or early March to prevent germination of crabgrass, bull paspalum, sedges (from seed), and spotted spurge.

Apply anytime broadleaf weeds are growing to get control.

Apply in the fall to control henbit, chickweed and other winter weeds.

If complete control is not obtained, a second application may be necessary 2-3 months after the first. For best results apply with a fertilizer spreader at the settings listed below. Apply to newly sprigged St. Augustine or Zoysia lawns to help prevent competition from other weeds. Apply when grass blades are dry. On hot days (above 80°) water the lawn to rinse particles off the grass.

SPREADER SETTINGS

Weed & Feed With Atrazine 25-4-6 5,000 Sq. Ft.	Scott's Drop (PF1) 5.0	Scott's Rotary (C2) 1	Cyclone (H1) 3.75	Sta-Green (EB84) 6.0	Ortho Rotary (4000) 3.0	Ortho Drop (3000) 4.0	Sears Drop (671, 19150) 6.0	Penn-ington (EB84) 6.0
--	------------------------	-----------------------	-------------------	----------------------	-------------------------	-----------------------	-----------------------------	------------------------

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides, or seeds.

PRODUCT DISPOSAL: To dispose of bags which contain unused products, securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

PRECAUTIONS:

Do not apply under the branches (rooting area) of shrubs, trees, flowers, etc. Do not apply to newly seeded Zoysia, centipede, or carpetgrass lawns. Wait until they have required at least 2 mowings. Don't apply to flowers, bedding plants, ornamental trees or shrubs.

CAUTION: We recommend that you keep this product (and all fertilizers) out of the reach of children. Avoid getting on concrete walks or drives, swimming pools, stucco or asbestos siding as the combination of certain fertilizer elements can cause permanent stains. Keep in a dry place.

BACK PANEL