

EXSPOR BASE LABEL - FRONT PANEL

**Alcide® Brand EXSPOR®
Part 1
Base Concentrate**

4:1:1 Sterilant-Disinfectant

**The 3-Minute System for Controlling Cross-Contamination from Treated Hard, Non-Porous Inanimate Surfaces
For Use with EXSPOR® Activator Concentrate (Part 2)**

Bactericidal - Virucidal - Fungicidal - Tuberculocidal - Pseudomonacidal - Sporocidal

Disinfecting Level - Kills all bacteria including Pseudomonas aeruginosa and Mycobacterium sp., pathogenic fungi and viruses*† on hard, non-porous environmental surfaces in 1 - 3 minutes (SEE DIRECTIONS FOR USE).

Sterilizes laboratory equipment and other non-medical immersible items in 10 hours at 20°C. (See directions for use. See side panel for precautionary information.)


***Kills Influenza Virus Type A2 (Hong Kong); Herpes Simplex Virus Types 1 and 2; Poliovirus Type 1; Coxsackievirus; Adenovirus Type 2; Echovirus Type 2; Vaccinia Virus and Vesicular Stomatitis Virus (VSV).**

†Currently, there is no established test for hepatitis viruses.


Active Ingredients:	Sodium Chlorite	1.52%
Inert Ingredients:		<u>98.48%</u>
TOTAL:		100.00%

**Keep Out of Reach of Children
CAUTION - See Side Panel
For Additional Precautionary Statements**

Contents 1 Gallon (3.785 Liters)



45631-3



EXSPOR ACTIVATOR LABEL - FRONT PANEL

**Alcide® Brand EXSPOR®
Part 2
Activator Concentrate**

**4:1:1 Sterilant-Disinfectant
The 3-Minute System for Infection and Contamination Control**



For Use with EXSPOR® Base Concentrate (Part 1)

Contains 9.5% Organic Acid

Follow directions for mixing, use, storage and disposal on the EXSPOR Base label. Wash measuring container before use. Base Concentrate must be diluted with water prior to adding Activator Concentrate (see "Preparation" section of Base Concentrate label). Improper preparation can result in excessive generation of chlorine dioxide fumes.

**Keep Out of Reach of Children
CAUTION - See Back Panel
For Additional Precautionary Statements**

Contents 1 Gallon (3.785 Liters)



3 4 1

EXSPOR BASE LABEL - SIDE PANEL

CAUTION

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

The following statement refers only to the unmixed EXSPOR Base concentrate:

CAUTION - HAZARDS TO HUMANS

Causes eye irritation. Avoid contact with eyes or clothing. Wash skin thoroughly with soap and water after handling.

FIRST AID: If in eyes, flush eyes with plenty of water. Call a physician if irritation persists.


U.S. Patent No. Re. 31,779

Foreign Patents

- EPA Reg. No. 45631-03
- EPA Est. No. 035-347-CA-001

Alcide Corporation
Redmond, Washington 98052

©1994 Alcide Corporation



4 7 7

EXSPOR ACTIVATOR LABEL - SIDE PANEL

U.S. Patent No. Re. 31,779
Foreign Patents

Alcide Corporation
Redmond, Washington 98052

©1994 Alcide Corporation

ACCEPTED
APR 10 1995
Under the Federal Insecticide, Fungicide, and
Rodenticide Act, as amended for the
pesticide registered under
EPA Reg. No. 45631-3

45631-3

4

5 47

EXSPOR BASE LABEL - BACK PANEL

Part 1

Base Concentrate

Alcide® Brand EXSPOR®

4:1:1 Sterilant-Disinfectant

EXSPOR Sterilant-Disinfectant is recommended for use as a disinfectant in schools, hospitals, medical, dental, industrial and institutional facilities to control human health-related microorganisms on hard, non-porous environmental surfaces and for sterilizing laboratory and other non-medical immersible items.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with the labeling. USE IN A WELL-VENTILATED AREA. AVOID BREATHING FUMES. See enclosed Technical Bulletins for additional use information, large scale/fogging applications and HIV-1 decontamination instructions. Base Concentrate must be diluted with water prior to adding Activator Concentrate (see "Preparation"). Improper preparation can result in excessive generation of chlorine dioxide fumes.

Preparation:

1. Measure 1 part EXSPOR Base Concentrate (Part 1) into a clean, empty, NON-METALLIC container.
2. Add 4 parts cool tap water (at or below 25°C).
3. Add 1 part EXSPOR Activator Concentrate (Part 2) and stir.

The final solution should appear colorless or pale yellow. Variations in local water hardness may result in darker color and/or stronger odor. Deionized or distilled water may be substituted to minimize these effects.

Moderate amounts of organic load, such as soil, will not lower the efficacy of EXSPOR solutions. However, to ensure best results, surfaces should be rinsed and, if possible, equipment dismantled and rinsed free of heavy soil and gross debris. Hollow objects must be flushed and filled. User judgment and standard practice may require rinsing with sterile water prior to drying.

DISINFECTION: Prepare EXSPOR solution as described above. Disinfection is not affected by water hardness up to 400 ppm (calculated as CaCO₃).

STERILIZATION OF LABORATORY EQUIPMENT: Immerse equipment for a minimum of 10 hours at 20°C. to destroy resistant pathogenic spores as represented by Clostridium Sporogenes and Bacillus subtilis. Remove equipment from Exspor solution using sterile technique and rinse thoroughly with sterile water.

Discard used solution after each day's use. Although efficacy at a 30-second contact time has been shown to be adequate against HIV-1, this time would not be sufficient for other organisms.

5

EXSPOR ACTIVATOR LABEL - BACK PANEL

Part 2
Activator Concentrate
Alcide® Brand EXSPOR®
4:1:1 Sterilant-Disinfectant

The following statement refers only to the unmixed EXSPOR Activator Concentrate:

CAUTION - HAZARDS TO HUMANS



Causes eye irritation. Avoid contact with eyes or clothing. Wash skin thoroughly with soap and water after handling.

FIRST AID: If in eyes, flush eyes with plenty of water. Call a physician if irritation persists.

See Part 1 Label - DIRECTIONS FOR USE

Warranty Limitation and Disclaimer

ALL EXPRESS OR IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR PARTICULAR PURPOSE OR OTHERWISE ARE EXPRESSLY EXCLUDED. THERE ARE NO WARRANTIES WHICH EXTEND BEYOND THE DESCRIPTION ON THE FACE HEREOF. In the event this product or its performance shall not conform with the description on the face hereof, Manufacturer's sole obligation shall be to refund to Purchaser the purchase price hereof, or to replace this unit of product with a conforming unit at Seller's option. Purchaser agrees that such refund or replacement shall be Purchaser's sole remedy for claims arising out of the purchase or use of this product. Before using this product, user shall determine the suitability of the product for its intended purpose and user assumes all risk and liability in connection therewith. User shall be responsible for its compliance with all federal, state and local laws and regulations regarding the product and its use, storage, handling and disposal.



Exspor Base Label - Back Panel (cont.)

EXSPOR solution will inactivate bacteria (including *M. tuberculosis*), fungi and viruses in 3 minutes at 20°C.

Soaking is the best method of application. However, the solution may be sprayed or wiped on by any convenient means (mop, cloth, sponge, etc.). The surface should be thoroughly wet with solution. To minimize potential for oxidation or corrosion, a contact time of 3 minutes on metal surfaces or equipment should not be exceeded.

NOTE: EXSPOR solution may be safely used on most plastics, ceramic, and glass. Never soak items in metal vessels. Prolonged or repeated exposure will oxidize metals, including aluminum and stainless steel. Do not use on copper, brass or bronze. Do not apply to surfaces containing residues from bleach, strong acids, or other chemicals, without first rinsing surface thoroughly with water.

STORAGE AND DISPOSAL

Store unmixed concentrates in a cool, dark, dry place in the original containers. Always replace covers. Keep upright. Thoroughly wash empty concentrate containers with water and discard in a safe place. Do not reuse containers. Used and outdated EXSPOR solution may be flushed down the drain. Prepared EXSPOR solution should be stored in a cool, dark place.

See Part 2 Activator Concentrate Label for Warranty Limitation and Disclaimer

