



FICAM [®] WW

76% WETTABLE POWDER INSECTICIDE

FOR USE ONLY BY PROFESSIONAL APPLICATORS

ACTIVE INGREDIENT:

Bendiocarb: (2,2-dimethyl-
1,3-benzodioxol-4-yl methylcarbamate†)

INERT INGREDIENTS:

Percent by Weight

76%

24%

Total

100%

†Protected by U.S. Patent No. 3,736,338

EPA REGISTRATION NO. 45639-1

PRECAUCION AL USARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicaca ampliamente

KEEP OUT OF REACH OF CHILDREN

WARNING

HARMFUL OR FATAL IF SWALLOWED

IF SWALLOWED — DRINK 1 OR 2 GLASSES OF WATER AND INDUCE VOMITING BY TOUCHING BACK OF THROAT WITH FINGER. DO NOT INDUCE VOMITING OR GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. GET MEDICAL ATTENTION.

IF INHALED — KEEP PATIENT STRICTLY AT REST AND TRANSFER TO HOSPITAL IMMEDIATELY TO GET MEDICAL ATTENTION.

IF ON SKIN — WASH WITH SOAP AND WARM WATER.

IF IN EYES — RINSE THOROUGHLY WITH CLEAN WATER.

NOTE TO PHYSICIAN: CONTAINS ANTICHOLINESTERASE CARBAMATE INSECTICIDE. IN CASES OF OVER-EXPOSURE, ATROPINE SULFATE — 2 MG. (EQUIVALENT TO 1/30 GRAIN ADULT DOSE) SHOULD BE GIVEN AND REPEATED AS NECESSARY UNTIL FULLY ATROPINIZED.

Manufactured by



TM

BFC Chemicals, Inc.
4311 Lancaster Pike, P.O. Box 2867
Wilmington, DE 19805

CANNgraphics August 13, 1982

456.37-1

IMPORTANT. Read the entire Directions for Use, Precautions and Disclaimer of Warranties before using this product.

GENERAL INFORMATION. FICAM W is a wettable powder containing 76% by weight active ingredient which has given effective knockdown and residual control of those insects listed below.

FICAM W does not damage paints, plastics, fabrics or other surfaces where water alone causes no damage.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Thoroughly wash out sprayer and strainer with water and detergent before using FICAM W.


For use by Professional Applicators for residual pest control in and on buildings and structures and their immediate surroundings and on modes of transport. Permitted areas of use include, but are not limited to, stores, warehouses, industrial buildings, houses, apartment buildings, mausoleums, greenhouses, laboratories, schools, nursing homes, hospitals, restaurants, hotels and food manufacturing, processing and servicing establishments and on vessels, aircraft, rail cars, buses, trucks and trailers.

FICAM W is intended for dilution with water for application with hand or power-operated sprayers. Do not use strainers finer than 100 mesh size. When using hand sprayers, fill with the desired volume of water, add FICAM W, close and shake before use to insure proper suspension of the wettable powder. Shake or re-agitate sprayer before use if spraying is interrupted.

Dosage Chart: For Conventional Spray Volumes

Use the scoop provided for measuring from the poly jar.

Each level scoopful contains 0.4 oz FICAM W, which equals one packet.

Amount of FICAM W	Amount of Water	Concentration of Spray Active Ingredient (ai)
1 packet or 1 scoop (0.4 oz) per 	Two gallons of water 	0.125%
1 packet or 1 scoop (0.4 oz) per 	Gallon of Water 	0.25%
2 packets or 2 scoops (0.8 oz) per 	Gallon of water 	0.50%
1 pound  per	40 gallons of water	0.25%

INDOOR USE

Cockroaches (including both non-resistant and organophosphate and organochlorine resistant strains), **Ants**, **Crickets**, **Silverfish** and **Firebrats**: Mix one packet or one scoop of FICAM W per gallon of water (0.25% ai) and apply as a coarse low pressure spot or crack and crevice spray to areas where these pests hide, such as baseboards, storage areas, closets, around water pipes, doors and windows, behind

and under refrigerators, cabinets, sinks and stoves, the underside of shelves, drawers and similar areas. For ants, apply to ant trails around doors and windows and other places where ants enter premises.

Following cleanout applications or where insect infestation is low and where sites are inspected and treated at monthly intervals or less, these insects can be controlled with sprays of one packet or one scoop of FICAM W per two gallons of water (0.125% ai).

Carpenter Ants: Mix one packet or one scoop of FICAM W per gallon of water (0.25% ai) and apply to ant trails, around doors and windows and other places where ants enter premises. Where possible, apply FICAM W directly to ant nest or infested wood.

Termites: Mix one packet or one scoop of FICAM W per gallon of water (0.25% ai) and apply to voids or channels in damaged wooden members of a structure, or to cracks, spaces or bearing joints between wooden members of a structure or between such members and the foundations, in locations vulnerable to termite attack such as crawl spaces. Drilling may be necessary prior to applying FICAM W to voids, cracks or joint spaces in vulnerable locations and to channeled sections of wood but it must be recognized that such drilling and treating of a timber in place may not result in complete coverage of infested and vulnerable voids or galleries. All treatment holes drilled in construction elements in commonly occupied areas of structures must be securely plugged.

The purpose of such applications of FICAM W is to kill workers or winged reproductive forms which may be present in the treated channels and spaces at the time of treatment and to impart a temporary resistance at these places to termite attack which may aid, in some part, in the prevention of immediate reinfestation by termites. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but merely a supplement, particularly where use of other insecticides may pose an odor problem.

Do not apply solution until location of heat or air conditioning ducts, vents, water and sewer lines and electrical conduits are known and identified. Extreme caution must be taken to avoid contamination of these structural elements and airways.

FICAM W at 0.25% ai has consistently provided control of all of the above insects for 30 and often for 60 days.

Brown Dog Ticks and Fleas In Buildings: Mix one packet or one scoop of FICAM W per gallon of water (0.25% ai) and thoroughly apply spray to infested areas such as pet beds and resting quarters, nearby cracks and crevices, rugs and floor coverings, and under their edges, between and under cushions of upholstered furniture and other areas where these pests may be present. Entire rugs may be treated. Apply the spray evenly (approx. 1 gal/800 ft.²) to rugs and furniture. Old bedding of pets should be removed and replaced with clean fresh bedding after treatment of pet areas. Additionally, insure that pets are treated with an EPA approved product registered for such use. Do not treat pets or their bedding with this product. Do not permit children or pets near treated surfaces until spray has dried completely.

Japanese Beetles — Aircraft: Mix one packet or one scoop of FICAM W per gallon of water (0.25% ai). Apply to areas where beetles may enter or be found such as carpets, under seats, storage areas, closets, around doors, behind and under galley equipment, in cargo holds and wheel wells.

Spiders (Including Black Widow Spiders): Mix one packet or one scoop of FICAM W per gallon of water (0.25% ai) and apply to infested baseboards, window and door frames, corners, pipes, storage localities, attics and under eaves. Make spot applications to other areas over which these pests may crawl.

Wasps and Bees: Mix one packet or one scoop of FICAM W per gallon of water (0.25% ai) and thoroughly spray nest and entrance and surrounding areas where insects alight. It is generally advisable to spray the nests in the evening when the insects are less active and have returned to the nest. For best results, check nest carefully one or two days after spraying to insure complete kill, then remove and destroy nest to prevent emergence of newly hatched insects.

Sowbugs, Pillbugs, Ground Beetles, Earwigs, Millipedes, Centipedes and Scorpions: Mix one packet or one scoop of FICAM W per gallon of water (0.25% ai) and apply around doors and windows, and other places where these pests may enter premises. Spray baseboards, storage areas and other locations where these pests are found.

45157-

Pantry Pests - Grain Beetles (Rusty, Merchant and Saw-toothed), Flour Beetles (Red and Confused), Drugstore Beetles, Rice Weevils, Lesser Grain Borers, Spider Beetles and Tobacco Moths: Treat exposed stages. Mix one packet or one scoop of FICAM W per gallon of water (0.25% ai). Remove all foodstuffs, shelving paper and utensils from area to be treated. Apply FICAM W to cupboards and shelves and allow to dry. Cover with shelving paper prior to replacement of food and utensils. Infested foodstuffs should be destroyed.

Carpet Beetles (Black, Furniture and Varied): Mix one packet or one scoop of FICAM W per gallon of water (0.25% ai). Apply as spot applications on and under edges of floor coverings, under rugs and furniture, and in closets or other localities where these insects are found. Entire rugs may be treated.

Note: Where it is desired to obtain a longer period of control of all of the above pests in areas where access is difficult or where deposits will not be quickly removed by cleaning, apply FICAM W at a rate of two packets or two scoops per gallon of water (0.50% ai). This rate may also be used to quickly reduce heavy infestations of cockroaches.

Webbing Clothes Moths: Mix two packets or two scoops of FICAM W per gallon of water (0.50% ai) and apply spray as spot application in closets and other storage areas where webbing clothes moths are found. This treatment is an adjunct treatment and will not control moth larvae already on the clothes. Do not apply FICAM W to clothing.

To prevent recurrence of pest infestations, repeat applications as required.

Interior Plants: Refer to the ornamental pests recommendations in the Outdoor Use section.

Food Handling Establishments — Places other than private residences in which food is held, processed, prepared or served.

Food Areas — Application limited to spot or crack and crevice treatments only. For spot treatments, no individual spot will exceed two (2) square feet.

Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas when food is exposed and facility is in operation are also considered a food area.

Prepare a spray solution as directed above and apply in small amounts directly into cracks and crevices using equipment capable of delivering a pinstream of insecticide in points between different elements of construction, between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs and bases where cockroaches, ants, silverfish, firebrats, spiders and crickets hide. Spot treatment using coarse spray may also be made to areas where any of the above pests hide or inhabit such as baseboards, storage areas, closets, around water pipes, doors and windows, behind and under refrigerators, cabinets, sinks, stoves and other equipment, the underside of shelves, drawers and similar areas.

Make applications using low pressure to avoid possible atomization or splashing of the spray. Avoid contamination of food or food processing surfaces.

Non-Food Areas — FICAM W may be used as a crack and crevice treatment or as a spot application in non-food areas. All areas where insects hide or through which insects may enter should be treated.

This product is authorized for use in non-food areas of Federally Inspected Meat and Poultry Plants.

OUTDOOR USE

Perimeter Treatment — Residual spray for control of ants, bees, brown dog ticks, centipedes, cockroaches, crickets, earwigs, firebrats, fleas, ground beetles, millipedes, pillbugs, scorpions, silverfish, sowbugs, spiders (including Black Widow Spiders), and wasps. Mix one packet or one scoop of FICAM W per gallon of water (0.25% ai) and apply as a residual spray or with a paint brush to surfaces of buildings, porches, screens, window frames, eaves, patios, garages and other areas where these pests congregate.

To help prevent infestations of buildings by the above pests, treat a band of soil and vegetation 6 to 10 feet wide around and adjacent to the

building. Also treat the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. Mix one packet or one scoop of FICAM W per gallon of water (0.25% ai) and apply as a coarse spray to thoroughly and uniformly wet the band area.

Yard Treatment — For the control of nuisance pest populations of ants, crickets, spiders, ground beetles, sowbugs and pillbugs, mix one packet or one scoop of FICAM W in FIVE gallons of water (0.05% ai) and apply as a coarse spray to thoroughly wet the treated areas.

Ornamental Pests: Use FICAM W to control specified pests on non-edible ornamentals such as Ageratum, Asarina, Azalea, Baby's breath, Begonia, Boxweed, Calendula, Cedar, Celosia, Chrysanthemum, Coleus, Daffodil, Dahlia, Dianthus, Euonymus, Ferns, Ficus, Flowering almonds, Flowering crab apples, Four-o'clock, Fuchsia, Gazania, Geranium, Holly, Hydrangea, Hypoestes, Impatiens, Iris, Ivy, Japanese yew, Juniper, Kalandra, Lilac, Lily of Valley, Marigold, Mimosa, Nasturtium, Nicotiana, Oleander, Pachysandra, Pansy, Petunia, Pine, Portulaca, Pyracantha, Rhododendron, Rose, Salvia, Schefflera, Shasta daisy, Snapdragon, Spirea, Taxus, Verbena, Vinca and Zinnia.

This product is highly toxic to bees exposed to direct treatment or to residues on plants.

Do not use on Cardoon, since plant injury may occur.

IMPORTANT: FICAM W has demonstrated excellent plant safety; however, all species and varieties have not been tested. When treating large numbers of plants of a single variety, treat a few plants and observe prior to full-scale application.

Apply FICAM W to point of run-off using recommended rates indicated below:

GYPSY MOTH

Egg Mass Treatment — Mix 4 oz. of FICAM W per 100 gallons of water and apply to egg masses until saturation. Applications can be made up to 3 weeks before gypsy moth hatch.

Larvae Treatment:

Residual Foliar and Contact Spray: Mix 3 oz. of FICAM W per 100 gallons of water and apply to the whole tree (wood and leaves). This will control all larvae stages. If required for optimum control in heavily infested areas, applications can be made every two weeks.

Supplementary Residual Trunk Spray: When large larvae (1-2½", 4-6 instars) are mobile and seeking resting places and a foliage spray is impractical, mix 2.5 lbs. (.25% a.i.) of FICAM W per 100 gallons of water and apply to tree trunks until fully saturated, making sure to treat cracks and crevices. A harborage (burlap or tape bands) is recommended to keep the larvae on the treated surface. If required for optimum control in heavily infested areas, applications can be made every two weeks.

DOSAGES AND DIRECTIONS FOR USE ON ORNAMENTAL PLANTS

PEST	AMOUNT OF FICAM W ¹ per 100 gallons of water
Chrysanthemum aphid, Eastern tent caterpillar, Green apple aphid, Hemispherical scale, Florida wax scale	3 oz
Brown soft scale, Spirea aphid, Tea scale, Citrus mealybug, Hawthorne lacebug, Fall webworm, Euonymus scale	6 oz
Japanese beetle, Thrips, Azalea lacebug, Azalea caterpillar, Taxus mealybug (adults), Elm leaf beetle, Willow leaf beetle (larvae), Poplar tent maker, Yellow-necked caterpillar, Spring cankerworm, Bagworm, Sycamore lacebug, Oleander caterpillar, Oleander aphid	11 oz
Mimosa webworm, Obscure root weevil (adults)	1 lb.
Greenhouse whitefly, Citrus blackfly (larvae), Black vine weevil, Pine spittlebug, Azalea caterpillar, Bronze birch borer ²	1 lb. 5 oz
Nantucket pine tip moth ³	2 lbs
Pales pine weevil, Northern pine weevil, Azalea leafminer	2 lbs. 10 oz

¹When measuring small amounts of product, use the 1 oz. scoop provided.

²Bronze Birch Borer: Use 3 sprays, spaced two weeks apart.

³Spray at the time of moth flight for each generation. Consult local Cooperative Extension authorities regarding time of moth flights in your area.

Ornamental Tree Pests: Mix FICAM W as recommended in the table below. Two gallons of mixture are recommended for mounds 8 inches or larger in diameter. Less mixture can be used on smaller mounds. However, thorough penetration of insect galleries must be achieved. Apply a small amount of material around the perimeter of the mound. Apply the remaining volume directly on the mound so as to penetrate insect galleries. A second treatment may be necessary if complete control is not obtained.

PEST	DOSEAGE	APPLICATION PROCEDURE
White grubs Japanese beetle Black tun grass stem borer European choker Northern masked choker Green June beetle Annual white grub Mole crickets	1/2 oz. per 1000 sq. ft.	Apply when pests are evident. Use adequate quantity of water to thoroughly moisten grass and thatch.
Chinch bugs Sod webworm Leafhoppers	0.5-1 oz. per 1000 sq. ft.	Apply when pests are evident. Use adequate quantity of water to thoroughly moisten grass and thatch.

Fire Ants: Mix FICAM W as recommended in the table below. Two gallons of mixture are recommended for mounds 8 inches or larger in diameter. Less mixture can be used on smaller mounds. However, thorough penetration of insect galleries must be achieved. Apply a small amount of material around the perimeter of the mound. Apply the remaining volume directly on the mound so as to penetrate insect galleries. A second treatment may be necessary if complete control is not obtained.

Do not disturb the mound prior to application. If possible, make applications after a recent rainfall. Applications may not be as effective during hot, dry periods. If possible, treat all colonies in the vicinity. Also, treat colonies which have not yet constructed a mound.

Dosage Table

AMOUNT OF FICAM W	AMOUNT OF WATER
2-3 teaspoons	2 gallons
1 pkg. or 1 scoop (0.4 oz.)	8 gallons

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

USE PRECAUTIONS

Harmful or fatal if swallowed. May be absorbed through the skin. Avoid contact with eyes, skin and clothing. If contact occurs, wash skin and clothing immediately with soap and warm water, and eyes with clean water.

Avoid breathing and working in spray mist or dust.

Wash hands and exposed skin before eating, drinking or smoking and after spraying or handling. Wash all contaminated clothing thoroughly before re-use.

Do not apply to plants that are being sprayed.

Do not use on treated surfaces. Do not use on treated surfaces.

Do not use on treated surfaces. Do not use on treated surfaces.

Do not spray where electrical shock circuits might occur.

Do not use on or contaminate fruit, vegetables, or other food crops.

Do not store near food or food products and do not contaminate these. Do not use on food preparation surfaces, dishes, kitchen utensils and food containers with either concentrate or spray.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water. Keep out of lawn and garden fish pools. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or to residues on plants.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures.

CONTAINER DISPOSAL: Packet — After insuring that both used packets and outer containers are empty, crush and dispose of in trash container.

Poly Jar — Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by state and local authorities.

DISCLAIMER OF WARRANTIES: THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OR MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of BFC Chemicals, Inc. is authorized to make any warranties beyond those contained herein or modify the warranties contained herein. UNDER NO CIRCUMSTANCES SHALL BFC CHEMICALS, INC. BE LIABLE FOR ANY INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM FAILURE TO USE THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS.

FOR FURTHER INFORMATION CONTACT ONE OF THE FOLLOWING OFFICES

BFC CHEMICALS, INC.
Suite 385
340 Interstate North Pkwy.
Atlanta, GA 30339
(404) 955-9441

BFC CHEMICALS, INC.
Suite 149
5100 North Sixth
Fresno, CA 93710
(209) 227-2911

BFC CHEMICALS, INC.
Airworld Center 320
10920 Ambassador Drive
Kansas City, MO 64153
(816) 891-0603

BFC CHEMICALS, INC.
4311 Lancaster Pike
PO Box 2867
Wilmington, DE 19805
(302) 575-7850

OCT 1 1982

RFC Chemicals, Inc.
P. O. Box 2867
4311 Lancaster Pike
Wilmington, DE 19805

Attention: Dr. K. D. Ihde

Gentlemen:

Subject: Ficam W Insecticide
Addition to use directions for gypsy moth control
EPA Registration No. 45639-1
Application Date: August 20, 1982

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,

Jay S. Ellenberger
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

TS-767:ELLENBERGER:DCR-24752:WANG-0400C:lisa:Raven:479-2013: 9/27/82