

POISON FIRST-AID TREATMENT

EMERGENCY

REMOVE PATIENT FROM TORAK EXPOSURE

IF SWALLOWED, INDUCE VOMITING
BY GAGGING WITH FINGER OR
WITH SALTY OR SOAPY WATER

NEVER GIVE ANYTHING BY MOUTH
TO AN UNCONSCIOUS PERSON

IF ON THE SKIN, REMOVE CONTAMINATED
CLOTHING AND IMMEDIATELY
WASH THE SKIN WITH SOAP AND WATER

CALL A PHYSICIAN IMMEDIATELY
IN ALL CASES OF SUSPECTED POISONING

If symptoms and signs of poisoning develop, i.e., headache, nervousness, blurred vision, nausea, cramps, pain in chest, sweating, constricted pupils, tearing, salivation, cyanosis, and muscle twitching, give 2 atropine tablets 1/100 grain each. Have the patient lie down and keep quiet.

TREATMENT BY PHYSICIAN

Give 2 to 4 mg (1/30 to 1/15 grain) atropine sulfate intravenously. Repeat at 5 to 10-minute intervals until signs of atropinization appear (dry, flushed skin and tachycardia). If excessive secretions occur, keep patient fully atropinized by giving atropine every hour up to 25 to 50 mg in one day. If patient fails to respond satisfactorily to atropine, give 1 gram of 2-PAM intravenously.

Artificial respiration and oxygen administration may be necessary. Decontaminate skin, stomach, and eyes as indicated. A mild degree of atropinization should be maintained for 24-48 hours, and the patient should be observed continuously for 48 hours and treated symptomatically.

DIRECTIONS FOR USE:

Apples (for use in states bordering on or east of the Mississippi River): For control of codling moth, plum curculio, and red-banded leaf roller, use 1 lb of Torak 50% Wettable Powder per 100 gal of water for full coverage sprays, starting with the petal fall spray and continuing in cover sprays as needed. For concentrate sprays, the volume of spray applied per acre should be reduced proportionally. Usually suppresses the European red mite in early season and midseason when used in a seasonal program. Do not use more than 5 lbs of product per acre per application at this rate.

For control of apple maggot and when the plum curculio is a serious problem, use 1.5 lbs of Torak 50% Wettable Powder per 100 gal of water in the petal fall spray and first cover spray. Continue in

cover sprays at the 1.0-lb-per-gal rate as needed to maintain control. Do not use more than 4.5 lbs of product per acre per application at this rate.

If additional control is needed after 6 sprays have been applied (petal fall + 5 covers), another pesticide (suitably registered) must be used. DO NOT apply more than 6 sprays per season.

Cracking of Stayman variety may be increased in some localities.

DO NOT apply within 60 days of harvest.

Consult your State Agricultural Experiment Station for the proper timing of sprays in your area.

ENVIRONMENTAL CAUTIONS:

This product is toxic to fish and wildlife. Keep out of lakes, ponds, and streams.

Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not apply to any area not specified on the label.

Keep out of tidal marshes and estuaries; shrimp and crab may be killed.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

USAGE CAUTIONS:

Observe the preharvest interval. Do not pick sooner than 60 days after the last spray. Workers should bathe or shower after each working day and should wash hands before eating. Workers should wear clothing that will minimize skin contact with foliage or other treated surfaces. Clean clothing should be worn each day.

During application keep unprotected persons away from areas being treated, or where there may be drift.

To avoid illegal residues do not apply within 60 days of harvest.

Do not combine with Bordeaux mixture.

Do not graze livestock in treated areas.

Do not feed treated cover crops to livestock or poultry.

Do not contaminate feed or foodstuffs.

Not for use or storage in or around the home environment.

Keep out of reach of children.

45639-44