

SPECIMEN LABEL

ACCEPTED

JUN 11 1991

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 45639-61

Mitac® WWP

50% WETTABLE POWDER

Insecticide for Pears

ACTIVE INGREDIENT:

Amtraz (N-(2,4-dimethylphenyl)-N-[(2,4-dimethylphenyl)methyl]-N-methylmethanamide)

Percent by Weight

INERT INGREDIENTS

50
50
Total 100

This product contains the toxic inert ingredient paraformaldehyde.

EPA Reg. No. 45639-61

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicada ampliamente.

KEEP OUT OF REACH OF CHILDREN

WARNING — AVISO

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with clean water. Get medical attention if irritation persists.

IF SWALLOWED: Drink promptly a large quantity of milk or water, whichever is available. Do not induce vomiting. If vomiting occurs, do not repeat. Avoid alcohol.

IF ON SKIN: Wash hands with soap and water immediately. If on clothing, remove clothing and wash it separately. Wash exposed skin with soap and plenty of water.

ANTIDOTE: None known.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

NOR-AM

NOR-AM CHEMICAL COMPANY
A Subsidiary of

3500 S. Riverside Blvd., P.O. Box 7195
Wilmington, DE 19803

NET CONTENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

WARNING

CAUSES EYE IRRITATION. HARMFUL IF SWALLOWED. REPEATED SKIN CONTACT MAY CAUSE AN ALLERGIC REACTION. DO NOT GET IN EYES. WEAR GOGGLES OR FACE SHIELD. WASH THOROUGHLY AFTER HANDLING. AVOID BREATHING DUST OR SPRAY MIST.

PROTECTIVE CLOTHING & EQUIPMENT

During mixing/loading or application wear a protective suit which has long sleeves and long pants. Wear chemical resistant gloves; a hat, boots and goggles or face shield. A helmet with visor may be substituted for the hat and goggles during aerial application. Mixer/loaders should also wear a chemical resistant apron when handling the concentrated product. Wash thoroughly with soap and water after handling and before eating, urinating, or smoking. Remove and wash clothing before reuse. Clothing should be laundered separately from household articles. Replace gloves frequently. Clothing which has been drenched and used gloves should be disposed of in accordance with state or local regulations. Instead of clothing and equipment specified above, the applicator can use an enclosed tractor cab or cockpit with properly filtered air supply.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to fish in adjacent sites. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this pesticide only as specified on this label.

Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep closed. Store in a cool, dry place.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DO NOT REUSE EMPTY CONTAINER.

IN CASE OF FIRE, LEAKY, OR DAMAGED CONTAINERS OR OTHER EMERGENCY, REPORT AT ONCE BY TOLL-FREE TELEPHONE TO: 800-424-9300.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

REENTRY STATEMENT: Do not reenter treated areas for 24 hours without protective clothing.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure.) When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

WARNING: Area treated with MITAC® WP on (date of application). Do not enter without appropriate protective clothing for 24 hours. In case of accidental exposure see Statement of Practical Treatment on the label.

USE ON PEARS: MITAC WP is an insecticide for use on pears. When applied as directed, MITAC WP will control pear psylla. MITAC WP can be applied by air or ground equipment, as dilute or concentrate sprays. The rate of MITAC WP per acre is based on thorough coverage with a dilute spray of 400 gallons from a conventional sprayer.

When using concentrate sprayers, the amount of water is reduced but the amount of MITAC WP per acre should remain the same as when using dilute sprays.

COMPATIBILITY: MITAC WP is compatible with most commonly used pear insecticides, miticides and fungicides; however, do not combine MITAC WP with detergents, sulfur, Bordeaux mixture, NAA or other highly alkaline materials as a tank mix.

CAUTION: DO NOT GRAZE TREATED AREAS

PEAR PSYLLA: Apply 1½ to 3 pounds MITAC WP per acre. Where full dilute coverage requires less than 400 gallons per acre, use 6 to 12 ounces per 100 gallons.

For prebloom control of overwintering adults, apply MITAC WP after eggs have moved into the nymphal psylla product, but prior to heavy egg laying. For best results, MITAC WP should be applied when daily maximum temperatures exceed 50°F. Consult your local University Extension Service for further information concerning best timing for control of overwintering adult psylla. If weather or orchard conditions do not permit spraying before heavy egg laying, spray after the majority of the first generation eggs have hatched.

For postbloom control, apply MITAC WP as needed. Best control will be obtained when the majority of pear psylla are in the adult or young nymphal stages of development.

Rate should not exceed 3 pounds of MITAC WP per acre per application. Do not apply within 7 days of harvest. Do not apply more than 6 pounds per acre of MITAC WP during the use season. Do not make more than two applications of MITAC WP per season.

IMPORTANT: READ BEFORE USE


By using this product, user accepts the following conditions, warranty, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, because of extreme weather and soil conditions, manner of use and other factors beyond NOR-AM Chemical Company's control, it is impossible for NOR-AM to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible.

DISCLAIMER OF WARRANTIES: THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of NOR-AM Chemical Company is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. UNDER NO CIRCUMSTANCES SHALL NOR-AM CHEMICAL COMPANY BE LIABLE FOR ANY INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM FAILURE TO USE THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS.

LIMITATIONS OF LIABILITY: In no event shall NOR-AM Chemical Company's liability for damages arising out of the use of this product exceed the purchase price of the product used.

© NOR-AM Chemical Company, 1985



3509 Silverside Road, P.O. Box 7495
Wilmington, DE 19803