SPECIMEN LABEL



ACCEPTED

JUL 1 4 1987

Under the .
Fungicities of the posticide registered under EPA Reg. No. 45631-66

# SYNERGIZED PYRETHRINS

## WETTABLE POWDER INSECTICIDE

Not for Sale to or Use by Homeowners.

#### **ACTIVE INGREDIENTS:**

†Bendiocarb: (2.2 dimethyl-1.3 benzodioxol-4-yl-methylcarbamate)

Pyrethrins

(

\*Piperonyl Butoxide Technical

**INERTINGREDIENTS:** 

Percent by Weight

3.06 7.66

59.83

Total

100.00

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que el etiqueta haya sido explicado ampliamente.

EPA Reg. No. 45639-66

## KEEP OUT OF REACH OF CHILDREN WARNING

## STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED — CALL A PHYSICIAN OR POISON CONTROL CENTER. GIVE 1 OR 2 GLASSES OF WATER AND INDUCE VOMITING BY TOUCHING FINGER TO BACK OF THROAT OR ADMINISTER IPECAC SYRUP USP ADULT DOSE 20 ML. CHILDREN 10 ML. BO, NOT INDUCE VOMITING OR GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS OR CONVULSING PERSON. GET MEDICAL ATTENTION

IF INHALED - KEEP PATIENT STRICTLY AT REST AND TRANSFER TO HOSPITAL IMMEDIATELY TO GET MEDICAL ATTENTION

IF ON SKIN - WASH WITH SOAP AND WARM WATER FOR 15 MINUTES.

IF IN EYES → FLUSH WITH PLENTY OF WATER FOR 15 MINUTES. GET MEDICAL ATTENTION

NOTE TO PHYSICIAN: CONTAINS ANTICHOLINESTERASE CARBAMATE INSECTICIDE IN CASES OF OVEREXPOSURE ATROPINE SULFATE 2
MG IERUIVALENT TO 1:30 GRAIN ADULT DOSE SHOULD BE GIVEN AND REPEATED AS NECESSARY UNTIL FULLY ATROPINIZED PRALIODXIME IP2SI IS
INEFFECTIVE, MORPHINE CONTRAINDICATED CONTROL CONVULSIONS WITH DIAZEPAM 10:20 MG. INTRAVENOUSLY.

## **MNOR-AM**

3509 Silverside Road, P.O. Box 7494 Wilmington, DE 19803 **NET CONTENTS:** 

And the second of a second of the

IMPORTANT: Read the entire Directions for Use, Precautions and Disclaimer of Warranties before using this product.

## PRECAUTIONARY STATEMENTS **HAZARDS TO HUMANS** AND DOMESTIC ANIMALS WARNING

Harmful or fatal if swallowed, May cause eye irritation. May be absorbed through the skin. Avoid contact with eyes, skin and clothing. If contact occurs, wash skin and clothing immediately with soap and warm water, and eyes with clean water.

Avoid breathing and working in spray mist or dust.

Wash hands and exposed skin before eating, drinking or smoking and after spraying or handling. Wash all contaminated clothing thoroughly before re-

Do not apply to humans, their clothing or bedding

Do not permit children or pets to contact treated surfaces until spray has dried

With proper spraying, Ficam\* Plus\* does not result in a visible deposit. However, if unevenly applied, or sprayed to run-off, a visible deposit may occur on some dark surfaces. Do not apply this product in conduits, motor housing, junction and switch boxes or other electrical equipment because of possible shock hazard. If there is a need to treat such areas. use oil-based sprays or Ficam\* D. Do not use on or contaminate fruit, vegetables or other feed or food crops. Do not store near feed or food products and do not contaminate these or food preparation surfaces, dishes, kitchen utensils and food containers with either concentrate or spray.

**ENVIRONMENTAL HAZARDS** 

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Keeptout of lawn and garden dish pools. Do not contaminate water 'by bleaning of equipment or disposal of wastes.

'This product is highly toxic to bees exposed to direct treatment or to residues on plants.

## "STORAGE AND DISPOSAL

Storage: Store ih a cool, dry place. Pesticide Disposal: Do not contaminate water, food or feed by storage or disposal.

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Container Disposal: Packet—After insuring both used packets and outer containers are empty, crush and dispose of in trash container.

**GENERAL INFORMATION:** Ficam Plus is a specially formulated wettable powder which gives knockdown and residual control of those pests listed below. It is especially suited for control of cockroaches.

Ficam Plus does not damage paint, plastics, fabrics, or other surfaces where water alone causes no damage.

**DIRECTIONS FOR USE** 

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Thoroughly wash out sprayer and strainer with water and detergent before using Ficam Plus. For use by service persons for residual pest control in and on buildings and structures and their immediate surroundings and on modes of transport. Permitted areas of use include, but are not limited to, stores, warehouses, industrial buildings, houses, apartment buildings, mausoleums, greenhouses, laboratories, schools, nursing homes, hospitals, restaurants, hotels and food manufacturing, processing and servicing establishments and on vessels, aircraft, rail cars, buses, trucks and trailers.

Applications should be made to hiding places, runways, and other places where pests are found.

Pay particular attention to: cracks and crevices; around water and sewer pipes; under and behind cabinets, refrigerators and sinks; around window and door frames; along baseboards; and in attics and crawl spaces.

Ficam Plus is intended for dilution with water for application with hand or power-operated sprayers. Do not use strainers finer than 50-mesh size. When using hand sprayers, fill with the desired volume of water, add Ficam Plus, close tightly, and invert sprayer several times before use to insure proper suspension of the wettable powder. Re-invert sprayer before use if spraying is interrupted.

Repeat treatments as necessary to maintain adequate control.

**Dosages for Conventional Spray** Volumes:

1. One packet Ficam Plus mixed in one gallon of water yields 0.25% ai. bendiocarb + 0.026% ai pyrethrins + 0.065% ai piperonyl butoxide.

2. Two packets of Ficam Plus mixed in one gallon of water yields 0.5% ai bendiocarb + 0.052% ai pyrethrins + 0.13% ai piperonyl butoxide.

INDOOR USE

Cockroaches (including both nonresistant and organochilorine resistant strains), Ants, Crickets, Sliverfish and Firebrats: Mix one packet of Ficam Plus per gallon of water and apply as a coarse low pressure spot or crack and crevice spray to areas where these pests hide, such as baseboards, storage areas, closets, around water pipes, doors and windows, behind and under refrigerators. cabinets, sinks and stoves, the underside of shelves, drawers and similar areas. For ants, apply to ant trails, around doors and windows and other places where ants enter premises.

Carpenter Ants: Mix one packet of Ficam Plus per gallon of water and apply to a in trails, around doors and windows and other places where ants enter premises. Where possible, apply Ficam Plus directly to ant nest or infested wood.

Termites: Mix one packet of Ficam Plus per gallon of water and apply to voids or channels in damaged wooden members of a structure, or to cracks, spaces or bearing joints between wooden members of a structure or between such members. and the foundations, in locations vulnerable to termite attack such as crawl spaces. Drilling may be necessary prior to applying Ficam Plus to voids, cracks or joint spaces in vulnerable locations and to channeled sections of wood but it must be recognized that such drilling and treating of a timber in place may not result in complete coverage of infested and vulnerable voids or galleries. All treatment holes drilled in construction elements in commonly occupied areas of structures must be securely plugged. The purpose of such applications of Ficam Plus is to kill workers or winged reproductive forms which may be present in the treated channels and spaces at the time of treatment and to impart a temporary resistance at these places to termite attack which may aid, in some part, in the prevention of immediate reinfestation by termites. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but merely a supplement, particularly where use

of other insecticides may pose an odor problem.

Do not apply solution until location of heat or air conditioning ducts, vents, water and sewer lines and electrical conduits are known and identified. When located, applications may be made around these structures However, caution must be taken to avoid contamination of these structural elements and airways.

Brown Dog Ticks and Fleas in **Buildings:** Mix one packet of Ficam Plus per gallon of water and thoroughly apply spray to intested areas such as pet beds and resting quarters, nearby cracks and crevices, rugs and floor coverings, and under their edges, between and under cushions of uphoistered furniture and other areas where these pests may be present. Entire rugs may be treated. Apply the spray evenly (approx. 1 gal/800 ft.2) to rugs and furniture. Old bedding of pets should be removed and replaced with clean fresh bedding after treatment of pet areas. Additionally, insure that pets are treated with an EPA approved product registered for such use. Do not treat pets or their bedding with this product. Do not permit children or pets to contact treated surfaces until spray has dried completely.

Spiders (including Black Widow Spiders): Mix one packet of Ficam Plus per gallon of water and apply to infested baseboards, window and door frames, corners, pipes, storage localities, attics and under eaves. Make spot applications to other areas over which these pests may crawl.

Wasps and Bees: Mix one packet of Ficam Plus per gallon of water and thoroughly spray nest and entrance and surrounding areas where insects alight. It is generally advisable to spray the nests in the evening when the insects are less active and have returned to the nest. For best results, check nest carefully one or two days after spraying to insure complete kill, then remove and destroy nest to prevent emergence of newly hatched insects.

Sowbugs, Pilibugs, Ground Beetles, Earwigs, Millipedes, Centipedes and Scorpions: Mix one packet of Ficam Plus per gallon of water and apply around doors and windows, and other places where these pests may enter premises. Spray baseboards, storage areas and other locations where these pests are found.

Pantry Pests - Grain Beetles (Rusty, Merchant and Sawtoothed), Flour Beetles (Red and Confused), Drugstore Beetles, Rice Weevils, Lesser Grain Borers, Spider Beetles and Tobacco Moths: Treat exposed stages. Mix one packet of Ficam Plus per gallon of water. Remove all foodstuffs, shelving paper and utensils from area to be treated. Apply Ficam Plus to cupboards and shelves and allow to dry. Cover with shelving paper prior to replacement of food and utensils. Infested foodstuffs should be destroyed.

Carpet Beetles (Black, Furniture and Varied): Mix one packet of Ficam Plus per gallon of water. Apply as spot applications on and under edges of floor coverings, under rugs and furniture, and in closets or other localities where these insects are found. Entire rugs may be treated.

Note: Where it is desired to obtain a longer period of control of all of the above pests in areas where access is difficult or where deposits will not be quickly removed by cleaning, apply Ficam Plus at a rate of two packets per gallon of water. This rate may also be used to quickly reduce heavy infestations of cockroaches.

Webbing Clothes Moths: Mix two packets of Ficam Plus per gallon of water and apply spray as spot application in closets and other storage areas where webbing clothes moths are found. This treatment is an adjunct treatment and will not control moth larvae already on the clothes. Do not apply Ficam Plus to clothing.

Food Handling Establishments — Places other than private residences in which food is held, processed, prepared or served.

Food Areas — Application limited to spot or crack and crevice treatments only. For spot treatments, no individual spot will exceed two (2) square feet.

Includes areas for receiving, storage, packing, (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas when food is exposed and facility is in operation are also considered a food area.

Prepare a spray solution as directed above and apply in small amounts directly into cracks and crevices using equipment capable of delivering a pinstream of insecticide in points between different elements of construction, between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs and bases where cockroaches, ants, silverfish, firebrals, spiders and crickets hide. Spot treatment using coarse spray may also be made to areas where any of the above pests hide or inhabit such as baseboards, storage areas, closets, around water pipes, doors and windows, behind and under refrigerators, cabinets, sinks. stoves and other equipment, the underside of shelves, drawers and similar areas.

Make applications using low pressure to avoid possible atomization or splashing of the spray. Avoid contamination of food or food processing surfaces.

Non-Food Areas — Ficam Plus may be used as a crack and crevice treatment or as a spot application in non-food areas. All areas where insects hide or through which insects may enter should be treated.

This product is authorized for use in non-food areas of Federally Inspected Meat and Poultry Plants.

### **OUTDOOR USE**

Perimeter Treatment — Residual spray for control of ants, bees, brown dog ticks, centipedes, cockroaches, crickets, earwigs, firebrats, fleas, ground beetles, millipedes, pillbugs, scorpions, silverfish, sowbugs, spiders (including Black Widow spiders), and wasps. Mix one packet of Flcam Plus per gallon of water and apply as a residual spray or with a paint brush to surfaces of buildings, porches, screens, window frames, eaves, palios, garages and other areas where these pests congregate.

To help prevent infestations of buildings by the above pests, treat a band of soil 6 to 10 (eet white around and adjacent to the building. Also, treat the building foundation to a height of 2 to 3 feet white epests were active and may find entrance. Mixtona packet of Ficam Plus per gallon of water and apply as a coarse spray to thoroughly and uniformly wet the band area.

Yard Treatment - For the control of nuisance pest populations of ants. crickets, spiders, ground beetles, ticks, sowbugs, and pillbugs, mix one packet of Ficam Plus in five gallons of water and apply as a coarse spray to thoroughly wet the treated areas.

INPORTANT: REAB BEFORE USE

By using this product, user accepts the following conditions, warranty, disclaimer of warrantes and firindations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, because of externe weather and soil conditions, manner of use and other factors beyond HOR-AM Chemical Company's control, it is impossible for NOR-AM to elemical company's control, it is impossible for NOR-AM to elemical company's control, it is impossible for NOR-AM to elemical company is outloaded as a result, crop injury or ineffectioners is always possible.

DISCLAIMER OF WARRANTIES: THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILLITY OR OF FITNESS FOR A PARTICULAR PUPPOSE OR OTHERWISE, WHICH EXTERD BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of NOR-AM Chemical Company is authorized to make any warrantes beyond those contained herein or to modify the warrantes contained herein. UNDER NO GIRCUMSTANCES SHALL NOR-AM CHEMICAL COMPANY BE LIABLE FOR ANY INCIDENTAL OR CONSCIUNTIAL DAMAGES RESULTING FROM FAILURE TO USE THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS.

LIMITATIONS OF LIABLITY: In no event shall NOR-AM Chemical Company's habiting of challists: (in no event shall HOR-AM Chemical Company's habiting for damages arising out of the use of this product exceed the purchase power of the product used. © NOR-AM Chemical Company, 1985



## ETTABLE POWDER INSECTICID

Not for Sale to or Use by Homeowners.

ACTIVE INGREDIENTS: 4 56

Bendiocarb: (2,2 dimethyl-1,3 benzodioxol-4-yl-methylcarbamate)

vrethrins

\*Piperonyl Butoxide, Technical

**INERT INGREDIENTS:** 

Percent by We

**Total** 

\*Equivalent to 6-1248° a (buty'carbityl) (6 propylpiperonyl) ather and 1,5312 🙀 related compounds

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que el etigi haya sido explicado ampliamente.

EPA Reg. No. 45639-66

## H OF CHILDREN KEEP OUT

IF SWALLOWED - CALL A PHYSICIAN OR POISON CONTROL CENTER. GIVE 1 OR 2 GLASSES OF WATER AND INDUCE VOMITING BY TOUCHING FINGER TO BACK OF THROAT OR ADMINISTER IPECAC SYBUP USP. ADULT DOSE 20 ML, CHILDREN 10 ML. DO NOT INDUCE VOMITING OR GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS OR CONVULSING PERSON. GET MEDICAL ATTENTION.

IF INHALED — KEEP PATIENT STRICTLY AT REST AND TRANSFER TO HOSPITAL IMMEDIATELY TO GET MEDICAL ATTENTION.

IF ON SKIN - WASH WITH SOAP AND WARM WATER FOR 15 MINUTES

IF IN EYES - FLUSH WITH PLENTY DF WATER FOR 15 MINUTES GET MEDICAL ATTENTION.

NOTE TO PHYSICIAN: CONTAINS ANTICHOLINESTERASE CARBAMATE INSECTICIDE. IN CASES OP-OVEREXPOSURE, ATROPINE SULFATE: 2 MG (EQUIVALENT TO 1/30 GRAIN ADULT DOSE) SHOULD BE GIVEN AND REPEATED AS NECESSARY UNTIL FULLY ATROPINIZED. PRAILID/DXIM'S IPZ INEFFECTIVE; MORPHINE CONTRAINDICATED. CONTROL CONVULSIONS WITH DIAZEPAM 10:20 MG. INTRAVENOUSLY.

3509 Silverside Road, P.O. Box 7495 Wilmington, DE 19803

**NET CONTENTS:** 

†Protected by U.S.Patent No. 3.7 (6.138)

IMPORTANT: Read the Entire Directions for User Prenautions and Diside memor Alertanties perfore living the princip.

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

Harmful or fatal if cwallowed May cause eye ordanch. May be absorbed through the skin Ascidic ordant with eyes, sein and cictiers. If contact 30 curs wast, skin and cotters from ediate by with soap and water and eyes with clean water.

Avoid breathing in a working in opinal mist or dust

Wash hands and exposed skin before eating, drinking or smoking at diafter spraying or handing. Wash all contaminated plothing thoroughly before reuse.

Do not apply to humans, their onthing or bedding

Do not permit children or pets to contact treated surfaces until spray has dried completely

With proper spraying, Ficam\* Plus\* does not result in a visible deposit However, if uneveniv applied, or sprayed to run-off, a visible deposit may occur on some dark surfaces Do not apply this product in conduits. motor housing, junction and switch. boxes or other electrical equipment because of possible shock nazard. It there is a need to treat such areas. use oil-based sprays or Ficam: D Do not use on or contaminate fruit, vegetables or other feed or food crops Do not store near feed or food products and do not contaminate these or food preparation surfaces, dishes, kitchen utensils and food containers with either concentrate or spray

### **ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds, and sother wildlife. Do not apply directly to water. Keepfort of lawn and garden fish pools. Do not contaminate water by bleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or to residues on plants

## STORAGE AND DISPOSAL

Storage: Store in a cool, dry place Pesticide Ulsposal: Do not contaminate water, food or feed by storage or disposal.

Wastes resulting from the use of this product may be disposed of on site or

at an approved wastershops at facility Container Disposal: Packet After incurring both used packets and outer centuries are empty, urgan and depole of intrase container.

GENERAL INFORMATION: Ficam Plus is a specially formulated wettable powder, which gives kneckdown and residual control of those pests listed below it is especially suited for control of cockroaches.

Figam Plus does not damage paint, prastic - fabrics or others creaces where water alone causes no damage.

### **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling

Thoroughly wash out sprayer and strainer with water and detergent before using Ficam Plus For use by service persons for residual pest control in and on buildings and structures and their immediate surroundings and on modes of transport. Permitted areas of use include, but are not limited to, stores warehouses, industrial buildings, houses, apartment buildings, mausoleums, greenhouses, laboratories, schools, nursing homes, hospitals. restaurants, hotels and food manufacturing, processing and servicing establishments and on vessels, aircraft, rail cars, buses, trucks and trailers

Applications should be made to hiding places, runways, and other places where pests are found. Pay particular attention to cracks and crevices, around water and sewer pipes, under and behind cabinets, refrigerators and sinks, around window and door frames, along baseboards, and in attics and crawl spaces.

Ficam Plus is intended for dilution with water for application with hand or power-operated sprayers. Do not use strainers finer than 50-mesh size. When using hand sprayers, fill with the desired volume of water, add Ficam Plus, close tightly, and invert sprayer several times before use to insure proper suspension of the wettable powder. Re-invert sprayer before use if spraying is interrupted.

## Repeat treatments as necessary to maintain adequate control. Dosages for Conventi mal Spray Volumes:

 One packet Figam Plus mixed in one gallon of water yields 0.25% at bendiocarb + 0.026% at pyrethrins + 0.065% at piperonyl butoxide. 2. Two packets of Figam Plus mixed in one galian of water verido 6.5 car bendiocarb + 2.50 car professions + 0.15 car percey but one

#### **INDOOR USE**

Cockroaches (including both nonresistant and organochlorine resistant strains), Ants, Crickets, Silverfish and Firebrats: Mo one packet of Ficam Plus per dation of water and apply as a coan ellow premare spet or crack and crevice spray thateas where these pests hide, such as baseboards, storage areas, closure around water pipes, doors, and wit dows, behind and under refrigerator cabinets, sinks and stoves, the underside of shelves, drawers and similar areas. For ants, apply to ant trails, around doors and windows and other places where ants enter premises

Carpenter Ants: Mix one packet of Ficam Plus per gallon of water and apply to ant trails, around doors and windows and other piaces where ants enter premises. Where pous-tile apply Ficam Plus directly to ant nest or infested wood.

Termites: Mix one packet of Ficarra Plus per gation of water and apply to voids or channels in damaged. wooden members of a structure, or to cracks, spaces or bearing joints between wooden members of a structure or between such members. and the foundations, in locations vulnerable to termite attack such as crawl spaces. Drilling may be necessary prior to applying Ficam-Plus to voids, cracks or joint spaces in vulnerable locations and to channeled sections of wood but it must be recognized that such drilling and treating of a timber in place may not result in complete coverage of infested and vulnerable voids or galleries. All treatment holes drilled. in construction elements in commonly occupied areas of structures must be securely plugged The purpose of such applications of Ficam Plus is to kill workers or winged reproductive forms which may be present in the treated channels and spaces at the time of treatment and to impart a temporary resistance at these places to termiteattack which may aid, in some part, in the prevention of immediate reinfestation by termites. Such applications are not substitute for mechanical alteration, soil treatment or foundation treatment but merely a supplement, particularly where use

of other insecticide in a local habitide problem.

Do not apply solution of the atlant of the atlant and sewer the expression of the atlant and sewer the expression of the atlant and sewer the expression of the expression of

Brown Dog Ticks and Fleas in Buildings: Morone parket of Feach Plus perigation of water or to thoroughly apply spray to the test dead such as petibed in 10% this quarters inearby or a kill of a recibes, fugs and fine and under their edge at his which and under cushions of uphorstered furniture and other areas where these pests may be prefernt. Entire rugs may be treated Applette bray eveniy (approx\_1 gat 860 ft and ruge and furniture. Old bedding of cet should be removed and replaced with clean fresh begding after treatment of pet areas. Ad affor any insure that pets are treated with an EPA approved product registered for such use. Do not treat certific them pedding with this product. Be not permit children or pets to contact treated surfaces until spray transcried completely

Spiders (including Black Widow Spiders): Mix one packet of Finant. Plus per gallon of water and apply to infested baseboards, window and door frames, corners, pipes istorage localities, attics and under eaves. Make spot applications to other areas over which these pests may crawl

Wasps and Bees: Mix one packet of Ficam Plus per gallon of water and thoroughly spray nest and entrance and surrounding areas where insects alight. It is generally advisable to apray the nests in the evening when the insects are less active and have returned to the nest. For best results check nest carefully one or two days after spraying to insure complete kill then remove and destroy nest to prevent emergence of newly hatched insects.

Sowbugs, Pillbugs, Ground Beetles, Earwigs, Millipedes, Centipedes and Scorpions: Mix one packet of Ficam Plus per gallon of water and apply around doors and windows, and other places where these pests may enter premises. Spray baseboards, surage areas and other locations where these pests are found.

Pantry Pests - Grain Beetles (Rusty, Merchant and Sawtoothed), Flour Beetles (Red and Confused), Drugstore Beetles. Rice Weevils, Lesser Grain Borers. Spider Beetles and Tobacco **Moths:** Freat exposed stade - N > one packet of Figam Plus per disk to of water. Remove all foodstuff shelving paper and utendes from area to be treated. Apply Figam Fig. to cupboards and shelves and a → ∴ to dry. Cover with shelving paper prior to replacement of food and stensis. Infested foodstuffs in that be destroyed

Carpet Beetles (Black, Furniture and Varied): Mix one packet of Ficam Plus per gallon of water. Appay as spot applications on and under edges of floor coverings, under rugs and furniture, and in closets or other localities where these insects are found. Entire rugs may be treated

Note: Where it is desired to obtain a longer period of control of all of the above pests in areas where access is difficult or where deposits will not be quickly removed by cleaning apply Ficam Plus at a rate of two packets per gallon of water. This rate may also be used to quickly reduce heavy infestations of cockroacher.

Webbing Clothes Moths: Mix two packets of Ficam Plus per galion of water and apply spray as spot application in closets and other storage areas where webbing clothes moths are found. This treatment is an adjunct treatment and will not control moth larvae already on the clothes. Do not apply Ficam Plus to clothing.

Food Handling Establishments — Places other than private residences in which food is held, processed, prepared or served

Food Areas — Application limited to spot or crack and crevice treatments only. For spot treatments, no individual spot will exceed two (2) square feet.

Includes areas for receiving storage packing, (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas when food is exposed and facility is in operation are also considered a food area.

Prepare a spray solution is a treatest above and apply in imalians (un) directly into cracks and in the using equipment car able of der vering a pinstream of the end of the in points between different comment of construction, between a pupilier t and floors, openings earning in and nollow spaces in will equipment legs and baller is their cockroaches, ants, sweet in firebrats, spiders and of check in the Spot treatment using Coat or 1711. may also be made to are a line of any of the above pests hipsing inhabit such as baseboard in the p areas, closets, around #ate (14) doors and windows (Pf. 1.3.3) under refrigerators, cation to line stoves and other equipment the underside of shelves, drawer, and similar areas

Make applications using low pressure to avoid possible atomization or splashing of the spray. Avoid contamination of food processing surface.

Non-Food Areas — Figure Property be used as a crack and creative to treatment or as a spot applicative to non-food areas. All areas where insects hide or through which in a tomay enter should be treated.

This product is authorized for use in non-food areas of Federally Inspected Meat and Poultry Plants.

## **OUTDOOR USE**

Perimeter Treatment — Residual spray for control of ants, bees brown dog ticks, centipedes, chakroaches crickets, earwigs, firebrats, fleas ground beetles, millipedes, pillbugs scorpions, silverfish, sowbugs, spiders (including Black Widow spiders), and wasps. Mix one packet of Ficam Plus per gallon of water and apply as a residual spray or with a paint brush to surfaces of buildings, poiches screens, window frames, eaves patios, garages and other areas where these pests congregate.

To help prevent infestations of buildings by the above pasts, treat a band of soil 6 to 10 feet wide around and adjacent to the building. Also treat the building foundation to a height of 2 to 3 feet where pests the active and may find entrance. Missioner packet of Ficam Plus per dallon of water and apply as a coasse spray to thoroughly and uniformly wet the band area.

Yard Treatment — For the control of nuisance pest populations of ants, crickets, spiders, ground beetles ticks, sowbugs, and pillbugs, mix onepacket of Ficam Plus in five gallons of water and apply as a coarse spray to thoroughly wet the treated areas

#### IMPORTANT READ BEFORE USE

IMPORTANT READ BEFORE USE

TO CONDITIONS TO A CONTROL OF THE PROPERTY OF THE P

• In Learns Into Lucio LIMITATIONS OF LIABILITY In not exercise at MVE AM Internal Limitation Supplied to the BOPS at Society of the Lucion to Supplied except the politicise or centified providing sent.

NOR AM THE CONTRACTOR

