12 12 120 160

SPECIABEL



ACCEPTED

SYNERGIZED: PYRETHRINS

WETTABLE POWDER INSECTICIDE

RECOMMENDED FOR USE ONLY BY PEST CONTROL OPERATORS

ACTIVE INGREDIENTS:

(

†Bendiocarb: (2,2 dimethyl-1,3 benzodloxol-4-yl-methylcarbamate)

Pyrethrins
*Piperonyl Butoxide, Technical
INERT INGREDIENTS:

Percent by Weight

29.45% 29.06

7.68

Total =

00.00F

*Equivalent to 6 1248% (butylcarbityl) (6-propylpiperonyl) ether and 1 5312% related compounds

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que el etiqueta haya sido explicado ampliamente.

EPA Reg. No. 45639-66

KEEP OUT OF REACH OF CHILDREN WARNING TATEMENT OF PRACTICAL TREATMEN

IF SWALLOWED - CALL A PHYSICIAN OR POISON CONTROL CENTER. GIVE 1 OR 2 GLASSES OF WATER AND INDUCE VOMITING BY TOUCHING FINGER TO BACK OF THROAT OR ADMINISTER IPECAC SYRUP USP. ADULT DOSE 20 ML, CHILDREN 10 ML, DO NOT INDUCE VOMITING OR GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS OR CONVULSING PERSON. GET MEDICAL ATTENTION.

IF INHALED - KEEP PATIENT STRICTLY AT REST AND TRANSFER TO HOSPITAL IMMEDIATELY TO GET MEDICAL ATTENTION.

IF ON SKIN - WASH WITH SOAP AND WARM WATER FOR 15 MINUTES.

IF IN EYES - FLUSH WITH PLENTY OF WATER FOR 15 MINUTES. GET MEDICAL ATTENTION.

NOTE TO PHYSICIAN: CONTAINS ANTICHOLINESTERASE CARBAMATE INSECTICIDE. IN CASES OF OVEREXPOSURE, ATROPINE SULFATE-2 MG.(EDUIVALENT TO 1/30 GRAIN ADULT DOSD SHOULD BE GIVEN AND REPEATED AS NECESSARY UNTIL FULLY ATROPINIZED PRALIDOXIME (P2S) IS INEFFECTIVE; MORPHINE CONTRAINDICATED. CONTROL CONVULSIONS WITH DIAZEPAM 10-20 MG. INTRAVENOUSLY

WNOR-AND NOR AM CHEMICAL COMPANY

3509 Silverside Road, P.Q. Box 7495 Wilmington, DE 19803

NET CONTENTS:

†Protected by U.S. Palent No. 3 736 338

IMPORTANT: Read the entire Directory of the Frecautions and Discourse of Warranties before using the product.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

Practice of later if swallowed. May be about the through the skin. Avoid contact with the Hyes, skin, and each of the fitting intriduction, with so about a water, and eyes with them water.

De at the eyes or mouth with hard the local sick of any way. STOP WORK and get help right away. See First Aid (Fractical Treatment) section.

Avoid breathing and working in spray must on dust

When applying in outdoor sites keep all unprotected persons, children, and animals away from treated area or where there is danger of drift.

Wash harids and exposed skin before eating, drinking or smoking and after spraying or handling. Wash all contaminated clothing thoroughly before recise.

Do not apply to humans, their clothing or bedding

Do not permit children or pets to contact treated surfaces until spray has dried completely

With proper spraying, Ficam* Plus* does not result in a visible deposit. However, if unevenly applied, or sprayed to run-off, a visible deposit may occur on some dark surfaces. Do not apply this product in conduits. motor housing, junction and switch boxes or other electrical equipment because of possible shock hazard. If there is a need to treat such areas. use oil-based sprays or Ficam* D Do not use on or contaminate fruit. vegetables or other feed or food crops. Do not store near feed or food products and do not contaminate these or food preparation surfaces, dishes, kitchen utensils and food containers with either concentrate or spray.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of waste.

This project is highly toxic to bees exposed to direct treatment on idealing appropriate allow it to drift to the control the treatment area.

STORAGE AND DISPOSAL

Storage: Store in a cool, dry place **Pesticide Disposal:** Do not contaminate water food or feed by storage in disposal.

Whether the uting from the use of this product may be disposed of on site or at an approved waste disposal facility **Container Disposal:** Packet—After insuring both used packets and outer containers are empty, crush and

dispose of in trash container

GENERAL INFORMATION: Ficam

Plus is a specially formulated wettable powder which gives knockdown and residual control of those pests listed

of cockroaches Ficam Plus does not damage paint, plastics, fabrics, or other surfaces where water alone causes no damage

below it is especially suited for control

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Thoroughly wash out sprayer and strainer with water and detergent before using Ficam Plus For use by service persons for residual pest control in and on buildings and structures and their immediate surroundings and on modes of transport. Permitted areas of use include, but are not limited to, stores, warehouses, industrial buildings, houses, apartment buildings, mausoleums, greenhouses, laboratories, schools, nursing homes, hospitals, restaurants, hotels and food manufacturing, processing and servicing establishments and on vessels, aircraft, rail cars, buses, trucks and

Applications should be made to hiding places for ways and other places where pests are found. Pay particular attention to cracks and crevices, around water and sewer pipes, under and behind cabinets, refrigerators and sinks, around window and door frames, along base boards, and in attics and crawl spaces.

Ficam Plus is intended for dilution with water for application with hand or power-operated sprayers. Do not use strainers finer than 50-mesh size

When using hand sprayers, fill with the desired volume of water, add Fidam-Plus, close tightly, and invert sprayer several times before use to insure or per suspension of the wettable bowder. Re-invert sprayer before use if spraying is interrupted.

Repeat treatments as necessary to maintain adequate control.

Dosages for Conventional Spray Volumes:

- One packet Fidam Plus mixed in are gallion of water yields 0.25 as bendiocarb + 0.026°, ai pyrethrins + 0.065° ai piperonyl butoxide
- 2 Two packets of Ficam Plus mixed mone gation of water yields 0.5 Labendiocarb + 0.052% ai pyrethrins + 0.13% ai piperonyl butoxide

INDOOR USE

Cockroaches (including both nonresistant and organochlorine resistant strains), Ants, Crickets, Silverfish and Firebrats: Mix one packet of Ficam Plus per gallon of water and apply as a coarse low pressure spot or crack and crevice spray to areas where these pests hide, such as baseboards, storage areas, closets. around water pipes, doors and windows, behind and under refrigerators, cabinets, sinks and stoves, the under side of shelves, drawers and similar areas. For ants, apply to ant trails, around doors and windows and other places where ants enter premises

Carpenter Ants: Mix one packet of Ficam Plus per gallon of water and apply to ant trails, around doors and windows and other places where ants enter premises. Where pc sible, apply Ficam Plus directly to ant nest or infested wood.

Termites: Mix one packet of Ficam Plus per gallon of water and apply to voids or channels in damaged wooden members of a structure, or to cracks, spaces or bearing joints between wooden members of a structure or between such members and the foundations, in locations vulnerable to termite attack such as crawl spaces. Drilling may be necessary prior to applying Ficam Plus to voids, cracks or joint spaces in vulnerable locations and to channeled sections of wood but it must be recognized that such drilling and treating of a timber in place may not result in complete coverage of infested and vulnerable voids or galleries. All treatment holes drilled in construction elements in commonly occupied areas of

Litrary are amost be securely paugged The purpose of such applications of fix at if it is to kill workers or with the controductive forms, which ma, be present in the treated manner i land spallers at the time the control of the tracket attention of the Promise in the Phaces to formite arrow and home, and in some part on a place to not immediate re invitation his termites. Supt at ν , at ν parent of a substitute for menture, a teración, sor treatment on the later in treatment but merely a Experient participanty where disstill the clinibectic designay poser at Same of the same

Direction is a condition and leads an of near or an conditioning ducts over the warm and sever times and electrical conducts are known and identified.

the register applications may be made on these structures on the selectructures of the taken to all its order matters of these of the taken to the selection of these of the taken to the selection of these of the taken to the selection of these of the taken to take the taken to the taken to the taken to take the taken to taken to take the taken to take the taken to take the t

Brown Dog Ticks and Fleas in Buildings: Mix one packet of Figam Plus perigation of water and thero , any apply spray to infested areas such as pet beds and resting gualities, hearby cracks and crevious rugs and floor coverings and under their edges, between and under cushions of upholstered furniture and other areas wherethese pests may be present. En: ++ rugs may be treated. Apply the suray eventy (approx 1 gal/800 ft) to rugs and furniture. Old bedding of pets. should be removed and replaced with clear, fresh bedding after treatment of pet areas. Additionally, insure that pets are treated with an EPA approved product registered for such use. Do not treat pets or their bedding with this product. Do not permit children or pets to contact treated surfaces until spray has dried completely

Spiders (including Black Widow Spiders): Mix one packet of Ficam Plus per gallon of water and apply to infested baseboards, window and door frames corners, pipes, storage localities attics and under eaves Make spot applications to other areas over which these pests may crawl

Wasps and Bees: Mix one packet of Ficam Plus per gallon of water and thoroughly spray nest and entrance and surrounding areas where insects alight. It is generally advisable to spray the nests in the evening when the insects are less active and have

returned to the nest. For best results checking at carefully one or two days after spraying to insure complete kill their remove and destroy nest to tracer to hergence of newly hatched in a street.

Sowbugs, Pillbugs, Ground Beetles, Earwigs, Millipedes, Centipedes and Scorpions: Mix one packet of Enam Prus per gallon of water and apply around doors and windows, and other places where these pests may enter premises. Spray baseboards for any areas and other locations where these pests are found.

Pantry Pests - Grain Beetles (Rusty, Merchant and Sawtoothed), Flour Beetles (Red and Confused), Drugstore Beetles, Rice Weevils, Lesser Grain Borers. Spider Seetles and Tobacco Moths: Treat exposed stages. Mix one packet of Figam Plus per gailon of water. Remove all foodstuffs. sherting paper and utensils from area to be treated. Apply Figam Plus to cuppoards and shelves and allow to dry. Cover with shelving paper prior to replacement of food and stensils. Infested foodstuffs should DH destroyed

and varied): Mix one packet of Floam Pics per gallon of water. Apply a spet applications on and under edges of floor coverings, under rugs and furniture, and in closets or other localities where these insects are found. Entire rugs may be treated. Note: Where it is desired to obtain a longer period of control of all of the above pests in areas where access is difficult or where deposits will not be quickly removed by cleaning, apply Floam Plus at a rate of two

packets per gallon of water. This rate

may also be used to quickly reduce

Carr et Beetles (Black, Furniture

Webbing Clothes Moths: Mix two packets of Ficam Plus per gallon of water and apply spray as spot application in closets and other storage areas where webbing clothes moths are found. This treatment is an adjunct treatment and will not control moth larvae already on the clothes. Do not apply Ficam Plus to clothing.

Food Handling Establishments — Places other than private residences in which food is held, processed, prepared or served

Food Areas — Application limited to spot or crack and crevice treatments only. For spot treatments, no individual spot will exceed two (2) square feet.

Includes are all for reserving istorage. parking to anning bottling, wrapping boxing) preparing, edible waste storage and enclosed processing systems (milis dairies ledible oils syrups). Serving areas, when tood is exposed and facility is \$10pe ation are also considered a Good area Prepare a spréy solution ansurecteu spoke and apply in small amount greatly into a racks and create usang égu pinent dapat kilor delikering a pinstream of inserticude in points between different element. of construction, between equipment and floors, openings leading to voids and hollow spaces in Walls, equipment legs and bases where cockroaches, ants, stive fish, firebrats, spiders and crickets hide Spot treatment using coarse spray may also be made to aleas where any of the above pests hide or inhabit such as baseboards, storage areas, closets, around water pipes, doors and windows, behind and under refrigerators, cabinets, sinks stoves and other equipment, the underside of shelves, drawers and similar areas

Make applications using low pressure to avoid possible atomization or splashing of the spray. Avoid contamination of food or food processing surfaces.

Non-Food Areas — Ficam Plus may be used as a crack and crevice treatment or as a spot application in non-food areas. All areas where insects hide or through which insects may enter should be treated.

This product is authorized for use in non-food areas of Federally Inspected Meat and Poultry Plants.

OUTDOOR USE

Perimeter Treatment — Residual spray for control of ants, bees, brown dog ticks, centipedes, cockroaches, crickets, earwigs, firebrats, fleas, ground beetles, millipedes, pillbugs, scorpions, silverfish, sowbugs, spiders (including Black Widow spiders), and wasps. Mix one packet of Ficam Plus per gallon of water and apply as a residual spray or with a paint brush to surfaces of buildings, porches, screens, window frames, eaves, patios, garages and other areas where these pests congregate.

1

To help prevent infestations of buildings by the above pests, treat a band of soil 6 to 10 feet wide around arid adjacent to the building. Also treat the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. Mix one packet of Ficam Plus per gallon of water and apply as a coarse spray to thoroughly and uniformly wet the band area

Yard Treatment - For the control of nuisance pest population; of ants crickets, spiders, ground paeties ticks, sowbligs, and fallbugs, ma one packet of Ficam Plus in tige oalions of water and apply as a coarse spray to thoroughly wet the treafed areas

IMPORTANT READ BEFORE USE

By using this produce user accepts the following conditions, war antidisclaimer of warranties and himitations of labelity
CONDITIONS. The directions also use of this product alle behaved to be
adequate and should be followed Sarefully. However, because if
extreme weather and soluconditions, manner of use and other rayto's
beyond NJR AM Chemical Company's control it is imprissible for
NJR AM to eliminate alt risks associated with the use of this indust
As a result crop injury or ineffectiveness is always possible.

DISCLAIMER OF WARRANTIES. THERE ARE NO WARRANTES.
EXPRESS OR IMPLIED. OF MERCHANTIABLITY OR OF EINESS IN A
PARTICULAR PURPOSE OR OTHERWISE, WHICH EXITED 15 VON.
THE STATEMENTS MADE ON THIS LABE. No agent of NIR AM
Chemical Conidany is authorized to make any warranties beyond muse
contained herein or to modify the warranties runnaned herein or to warranties warranties runnaned herein or to warranties warranties runnaned herein or to warranties runnaned herein A THICABEL DIRECTIONS LIMITATION OF CLABILITY IN CLEARNING NOR AMILITY N

• TRIAM Themical Sympatry, 1985.

Replaces 517-SL-FP-2 5M-Rep.3/88 Revisions Include. Warning Statements **Environmental Hazards Discard Previous Labels**

517-SL-FP-10M-Rev.6/88