

RESTRICTED-USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification

TAKTIC

EMULSIFIABLE CONCENTRATE

EPA REG. NO. \_\_\_\_\_

FOR THE CONTROL OF MANGE AND LICE ON SWINE

SUPPLEMENTAL LABELING

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT

Remove feed from pen and cover drinking bowls. Remove and destroy bedding. Hose out feces and excess feed. Use within 6 hours of mixing.

TREATMENT

Spray the animals until the mixture runs off, paying particular attention to treat the jowl, legs, inside the ears and underside of the body. Apply treatments as stated below, making sure to treat each animal.

CONTROL PROGRAM


Mix 2 fl. oz of TAKTIC EC with 3 gallons of water. A sprayer with a coarse nozzle, delivering the solution at a minimum of 200 psi is recommended. Spray the walls, floor, and any fittings in the pen. At the same time, treat animals whether visibly affected or not. Repeat the treatment 7-10 days after the first treatment has been made.

MAINTENANCE PROGRAM

Begin the maintenance program only after the double treatments have been made.

Mix 2 fl. oz of TAKTIC and 3 gallons of water. Apply with a conventional type sprayer and treat animals as recommended below.

1. Spray boars every 3 months.
2. Spray or dip (see below) pigs at weaning or before moving to the fattening/growing houses.
3. Isolate and spray any new or additional animals with the two initial treatments before mixing with the other animals.



# NOR-AM Chemical Company

(RESTRICTED USE PESTICIDE - CONTINUED)

## ERADICATION PROGRAM

Mix 2 fl. ounces with 3 gallons of water. A sprayer with a coarse nozzle, delivering the solution at a minimum of 200 psi is recommended. Spray the walls, floor, and any fittings in the pen. Spray animals (at the same time) whether visibly affected or not. Repeat the treatment 7-10 days after the first treatment has been made.

## MAINTENANCE PROGRAM

Begin the maintenance program only after the double treatments have been made. Mix 2 fluid ounces with 3 gallons of water. Apply with a conventional type sprayer and treat animals as recommended below.

1. Spray boars every 2 or 3 months.
2. Spray sows in their stalls before being transferred to pens.
3. Spray sows in their farrowing pens before being transferred back to sow stalls.
4. Dip (see below) or spray young pigs at weaning or before moving to fattening houses.
5. Isolate and spray any new or additional animals with the two initial treatments before mixing with the other animals.

## FOR DIPPING PIGLETS

Dip piglets (weaners) in a suitable size container. Dip 30 animals and then replace dip wash. Use dip wash to treat premises.

DO NOT apply to animals within **2** days of slaughter.

DO NOT open container until ready to use.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE TAKTIC E.C. CONTAINER LABEL ARE TO BE FOLLOWED.

THIS LABELING MUST BE IN POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

DISCLAIMER OF WARRANTIES: THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of NOR-AM Chemical Company is authorized to make any warranties beyond those contained herein or modify the warranties contained herein. UNDER NO CIRCUMSTANCES SHALL NOR-AM CHEMICAL COMPANY BE LIABLE FOR ANY INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM FAILURE TO USE THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS.

Sold by:

NOR-AM Chemical Company  
Wilmington, DE 19803 U.S.A.

