

Reg no: 45639-70

PM-12

193

# SPECIMEN LABEL

ACCEPTED

DEC 13 1988

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registration number  
EPA Reg. No. 45639-70

# Taktic<sup>®</sup>

EMULSIFIABLE CONCENTRATE



**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

In case of spillage, cover with an absorbent such as soda ash, lime, clay or sawdust. Sweep up and dispose of as directed under Storage and Disposal.

In case of fire or other emergency, report at once by toll free telephone to 800-424-9300.

must be functional and at the correct angles to thoroughly soak the animals within a few seconds. After filling with clean water and adding the appropriate amount of TAKTIC E.C., circulate the finished solution through the spray dip machine to ensure a homogeneous mixture. Applications should be made as instructed by USDA, APHIS, VS bulletins. Two treatments 7-10 days apart are required to control cattle scabies.



**SWINE  
MANGE MITES AND LICE:  
IMPORTANT**

Remove feed from pen and cover drinking bowls  
Remove and destroy bedding. Hose out feces and  
excess feed. Use TAKTIC solutions within 6 hours of  
mixing.

**TREATMENT**

Spray the animals until the mixture runs off, paying  
particular attention to treat the lower legs, inside the ears  
and underside of the body. Apply treatments as stated  
below, making sure to treat each animal.

**CONTROL PROGRAM**

Mix **ONE QUART CAN** (760 ml) of TAKTIC E.C. in 50  
gallons water or **ONE PINT CAN** (380 ml) in 25 gallons  
water. For smaller quantities of spray, mix 1 fl. ounce  
with 2 gallons water. A sprayer with a coarse nozzle,  
delivering the solution at 70 to 150 psi is  
recommended. Spray the walls, floor, and any fittings in  
the pen. Spray animals (at the same time) whether visibly  
affected or not. Repeat the treatment 7-10 days after the  
treatment has been made.

**MAINTENANCE PROGRAM**

Begin the maintenance program only after the double  
treatments have been made. Mix as described above.  
Apply with a conventional type sprayer and treat animals  
as recommended below.

1. Spray sows in their stalls before being transferred to  
farrowing pens.
2. Dip (see below) or spray young pigs at weaning or  
before moving to fattening houses.
3. Spray sows in their farrowing pens before being  
transferred back to sow stalls.
4. Isolate and spray any new or additional animals with  
the two initial treatments before mixing with the  
other animals.
5. Spray boars every 2 or 3 months.

**FOR DIPPING PIGLETS**

Dip piglets (weaners) in a suitable size container. Dip 30  
animals and then replace dip wash. Use dip wash to  
treat premises.

**RESTRICTIONS:**

**NO POST TREATMENT SLAUGHTER INTERVAL ON  
BEEF AND DAIRY CATTLE.**

**CAN BE USED ON LACTATING DAIRY ANIMALS.**

**NO WITHHOLDING PERIOD FOR MILK FOLLOWING  
APPLICATION.**

**DO NOT APPLY TO SWINE WITHIN ONE DAY OF  
SLAUGHTER.**

**DO NOT OPEN CONTAINER UNTIL READY FOR USE.**

**NOT FOR USE ON HORSES. MAY CAUSE FATAL  
COLON IMPACTION.**

