

# SPECIMEN LABEL

**ACCEPTED**

JUL 2 1992

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 45639-75

FOR AGRICULTURAL USE ONLY

# Carzol<sup>®</sup> SP

## MITICIDE/INSECTICIDE

**ACTIVE INGREDIENT:**

Formetanate hydrochloride\*

**Percent by Weight**

92%

**INERT INGREDIENTS:**

8%

**TOTAL**

100%

\*N, N dimethyl N [3 [(dimethylamino)carbonyl]oxy]phenyl] methanimidamide monohydrochloride

EPA Reg. No. 45639-74

EPA Est. No. 33689-WG-1

**PRECAUCION AL USUARIO:** Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicada ampliamente.

**KEEP OUT OF REACH OF CHILDREN**



**DANGER — POISON  
PELIGRO — VENENO**



### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED**

IF SWALLOWED, do not induce vomiting. Do not take any medicine unless directed by a physician.

**IF ON SKIN:**

IF ON SKIN, wash with plenty of water.

IF IN EYES, flush with water for 15 minutes.

**NOTICE TO PHYSICIAN**

Carzol SP is a form of Formetanate Hydrochloride. Advise your physician of this if you are taking any other medicine.

SEE REAR AND SIDE PANELS FOR  
ADDITIONAL PRECAUTIONS AND DIRECTIONS FOR USE  
DO NOT REUSE THIS BOX

**NOR-AM**

NOR-AM CHEMICAL COMPANY  
Agriculture Division

3509 University Road, P.O. Box 7446  
Wilmington, DE 19881

**NET CONTENTS:**

1.0 LITER (33.8 FL. OZ.) (33.8 FL. OZ.)

# PRECAUTIONARY STATEMENTS DANGER

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Fatal if swallowed. Do not breathe spray mist. Can cause irreversible eye damage. May cause skin sensitization. Do not get in eyes, on skin or on clothing. Wear goggles, face shield or safety glasses. Wear a pesticide respirator approved jointly by the Mining Enforcement and Safety Administration (formerly U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part II. Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash before reuse.

Do not apply this product in such a manner as to directly or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to wildlife. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Cover or incorporate spills.

This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treatment area. Do not allow animals to graze in orchard areas treated with CARZOL<sup>®</sup> SP.

## STORAGE AND DISPOSAL

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Completely empty box into application equipment. Then dispose of empty box in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**IN CASE OF FIRE, LEAKY OR DAMAGED CONTAINERS OR OTHER EMERGENCY, REPORT AT ONCE BY TOLL-FREE TELEPHONE TO: 800-424-9300.**

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

#### REENTRY STATEMENT:

Do not enter treated areas for 24 hours unless protective clothing is worn.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure.) When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

**DANGER: Area treated with CARZOL<sup>®</sup> SP on (date of application). Do not enter without appropriate protective clothing for 24 hours or until leaves are completely dry. In case of accidental exposure see Statement of Practical Treatment on the label.**

CARZOL SP is completely soluble in water. Preparation of a premix of slurry before adding to the spray tank is not necessary. Add the desired amount of CARZOL SP directly to the spray tank 1/2 full of water with the agitator running. Then add the remainder of the water.

NOTE: CARZOL SP is not stable in alkaline water. Do not prepare the spray mixture until just before application. Do not prepare more spray mixture than will be used within 4 hours.

### USE PRECAUTIONS

To avoid excessive residues, observe the following limitations:

- ▷ CITRUS: Do not apply more than 5 lbs. CARZOL SP per acre in a 12 month period.  
**Arizona and California:** Allow 14 days between applications.  
**Florida and Texas:** Allow 30 days between applications.  
NOTE: If brown soft scale is present in the grove, a suitable scallicide should be applied after the CARZOL SP application.
- ▷ APPLES, PEARS: Do not apply more than 4 lbs. CARZOL SP per acre in one crop season.
- ▷ PEACHES: Do not apply more than 2 1/2 lbs. CARZOL SP per acre in one crop season.
- ▷ NECTARINES: Do not apply more than 4 lbs. CARZOL SP per acre in one crop season.
- ▷ PLUMS, PRUNES: Do not apply more than 3 lbs. CARZOL SP per acre in one crop season.
- ▷ ALFALFA (Seed Crop Only): Use this product only on pure stands of alfalfa to be harvested for seed. Do not allow spray to drift onto adjacent crops. Do not spray while bees are visiting the treatment area. Do not allow livestock to graze treated fields. Do not use treated crops for livestock feed. Crop residues are not to be fed to livestock or baled for such use. Fields treated with CARZOL SP must be cleared after harvest by burning or by disking into the soil. CARZOL SP does not control aphids. If aphids appear, add an aphicide to the spray. Do not apply more than 2 lbs. CARZOL SP per acre per season. Do not make more than 2 applications of CARZOL SP per season. Allow 21 days before harvest.
- ▷ Do not apply this product through any type of irrigation system.

**BEST AVAILABLE COPY**

To avoid excessive residue observe the pre-harvest intervals shown in the table.  
For other use limitations see side panel under Use Precautions.

CROP	PEST	APPLICATION
ORANGES, LEMONS, TANGERINES, TANGELOS, AND GRAPEFRUIT (Arizona only)	THRIPS	Apply 1/4 to 1 1/4 lbs. of CARZOL SP per acre as outside coverage spray when thrips appear. For aerial applications, use a minimum of 10 gallons of water per acre. Apply as needed up to a limit of 5 lbs. CARZOL SP per acre per 12 month period. Allow 7 days before harvest.
ORANGES, LEMONS, TANGERINES, TANGELOS, AND GRAPEFRUIT (California Only)	THRIPS	Apply 1/4 to 1 1/4 lbs. of CARZOL SP per acre as outside coverage spray when thrips appear. For aerial application, use a minimum of 10 gallons of water per acre. Apply no more than twice per season. Allow 7 days before harvest.
ORANGES, LEMONS, TANGERINES, AND GRAPEFRUIT (Florida)	APOPKA WEEVIL, CITRUS ROOT WEEVIL (BLUE-GREEN WEEVIL), FULLER'S ROSE BEETLE, LITTLE LEAF NOTCHER	Apply 1 lb. CARZOL SP as a concentrate spray of 100 gallons per acre or less. Repeat application as needed. Allow 7 days before harvest.
ORANGES, LEMONS, TANGERINES, AND GRAPEFRUIT (Florida)	RUST MITE	Apply 1/2-1 lb. CARZOL SP per acre. Repeat application as needed. Allow 7 days before harvest.
ORANGES, LEMONS, TANGERINES, AND GRAPEFRUIT (Texas)	RUST MITE	Use 2-4 oz. CARZOL SP per 100 gallons of water when mites appear. For full cover dilute spray, apply 500 gallons of water per acre. Repeat application as needed. Allow 7 days before harvest.
APPLES	TENTIFORM LEAF MINER	Use 4-8 oz. CARZOL SP per 100 gallons of water at pink and petal fall. For full cover dilute spray, apply 400 to 800 gallons of water per acre depending on the size of the trees. For concentrate spray, apply 1-4 lbs. CARZOL SP per acre in at least 50 gallons of water. Allow 7 days before harvest.
APPLES	WHITE APPLE LEAFHOPPER	Use 2-4 oz. CARZOL SP per 100 gallons when leafhoppers appear. For full cover dilute spray, apply 400 gallons of water per acre. For concentrate spray, apply 1/2-1 lb. CARZOL SP per acre in at least 50 gallons of water. Repeat application as needed. Allow 7 days before harvest.
APPLES	CAMPYLOMMA	Use 4 oz. CARZOL SP per 100 gallons of water at pre-bloom/bloom and repeat as necessary through petal fall. For full cover dilute spray, apply 400 gallons of water per acre. For concentrate spray or dilute application of less than 400 gallons, apply 1 lb. CARZOL SP per acre in a minimum of 50 gallons of water per acre. Apply at night or while bees are not working in blooms. Allow 7 days before harvest.
APPLES AND PEARS	EUROPEAN RED MITE, TWO SPOTTED MITE	Use 4-8 oz. CARZOL SP per 100 gallons of water when mites appear. For full cover dilute spray, apply 400 to 800 gallons of water per acre depending on the size of the trees and severity of the mite infestation. For concentrate spray, apply 1-4 lbs. CARZOL SP per acre in at least 50 gallons of water. Repeat application as needed. Allow 7 days before harvest.
APPLES (Also see below for Arizona only)	THRIPS	Use 4-5 oz. CARZOL SP per 100 gallons of water at bloom and repeat as necessary through petal fall. For full cover dilute spray, apply 400 gallons of water per acre. For concentrate spray or dilute application of less than 400 gallons, apply 1 to 1 1/4 lbs. CARZOL SP per acre in a minimum of 50 gallons of water per acre. Apply at night or while bees are not working in blooms. Allow 7 days before harvest.
PEARS	PEAR RUST MITE	Use 4 oz. CARZOL SP per 100 gallons of water. Apply post bloom to harvest at first indication of rust mite infestation. For full cover dilute spray, apply 400-800 gallons of water per acre depending on the size of the trees. For concentrate spray, apply 1-2 lbs. CARZOL SP per acre in at least 50 gallons of water. Repeat application as needed. Allow 7 days before harvest.
PEACHES AND NECTARINES	EUROPEAN RED MITE, McDANIEL MITE, TWO-SPOTTED MITE	Use 4 oz. CARZOL SP per 100 gallons of water when mites appear. For full cover dilute spray, apply 400 to 500 gallons of water per acre depending on the size of the trees. For concentrate spray, apply 1-1 1/4 lbs. CARZOL SP per acre in at least 50 gallons of water. Repeat application as needed. Allow 21 days before harvest for peaches, 14 days for nectarines.
PEACHES	LYGUS BUGS, STINK BUGS	Use 4 oz. CARZOL SP per 100 gallons of water at first sign of infestation. For full cover dilute spray, apply 400-500 gallons of water per acre. For concentrate spray, apply 1-1 1/4 lbs. CARZOL SP per acre. Repeat application as needed. Allow 21 days before harvest.
NECTARINES	THRIPS	Use 4-8 oz. CARZOL SP per 100 gallons of water when thrips infest blossoms through petal fall. For full cover dilute spray, apply 400 to 500 gallons of water per acre depending on the size of the trees. For concentrate or aerial spray, apply 1-2 lbs. CARZOL SP per acre. Apply at night or while bees are not working in blooms. Repeat application as needed. Allow 14 days before harvest.
PLUMS AND PRUNES	EUROPEAN RED MITE, McDANIEL MITE, TWO-SPOTTED MITE	Use 4 oz. CARZOL SP per 100 gallons of water when mites appear. For full cover dilute spray apply 400 to 800 gallons of water per acre depending on the size of the trees. For concentrate spray, apply 1-1 1/4 lbs. CARZOL SP per acre in at least 50 gallons of water. Repeat application as needed. Allow 7 days before harvest.
ALFALFA (seed crop only in California, Oregon, Washington, Idaho, and Nevada. Use pre-bloom only on alfalfa in all areas except California)	LYGUS BUGS	Use 1/2-1 lb. CARZOL SP per acre. Apply when lygus bugs reach damaging numbers. Consult your State Agricultural Experiment Station or State Extension Service Specialist for proper timing.
	CONSPERSE STINK BUGS (California only)	Use 1 lb. CARZOL SP per acre for knockdown of a heavy nymphal population. A lygus control program will control stink bugs. When applied by air use a minimum of 5 gallons per acre. When applied by ground use a minimum of 20 gallons per acre.
	TWO-SPOTTED MITE	Use 1/2-1 lb. CARZOL SP per acre. Apply when mites appear. When applied by air use a minimum of 10 gallons of spray per acre. When applied by ground use a minimum of 20 gallons per acre.

**IMPORTANT: READ BEFORE USE**

By using this product, user or buyer accepts the following conditions, warranty, disclaimer of warranties and limitations of liability.


**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However, because of extreme weather and soil conditions, manner of use and other factors beyond NOR-AM Chemical Company's control, it is impossible for NOR-AM to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OR MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of NOR-AM Chemical Company is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. UNDER NO CIRCUMSTANCES SHALL NOR-AM CHEMICAL COMPANY BE LIABLE FOR ANY SPECIAL, INDIRECT, OR CONSEQUENTIAL DAMAGES RESULTING FROM USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES, OR DAMAGE RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT NOR-AM'S ELECTION, THE REPLACEMENT OF PRODUCT.

© NOR-AM Chemical Company, 1991

**BEST AVAILABLE COPY**



**NOR-AM**

NOR-AM CHEMICAL COMPANY

A Schering-Boehringer Company

AGRIBUSINESS DIVISION

3509 Silverside Road, P.O. Box 7495

Wilmington, DE 19803

 Printed on Recycled Paper

5-29-92, OPP 161138

PM19

45639-74

184

# SPECIMEN LABEL

**BEST AVAILABLE COPY**

**ACCEPTED**

**JUL 2 1992**

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 45639-74

FOR AGRICULTURAL USE ONLY

# Carzol<sup>®</sup> SP

## MITICIDE/INSECTICIDE

ACTIVE INGREDIENT	Percent by Weight
Formetanilol	95.0
INERT INGREDIENTS	5.0
	<b>TOTAL</b>
	100.0

EPA Reg. No. 45639-74

EPA Est. No. 33689-WG-1

**PRECAUCION AL USUARIO:** Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicada ampliamente.

**KEEP OUT OF REACH OF CHILDREN**

 **DANGER — POISON** 

 **PELIGRO — VENENO** 

**STATEMENT OF PRACTICAL TREATMENT**

IF SWALLOWED:

IF ON SKIN:

IF IN EYES:

NOTICE TO PHYSICIAN:

SEE REAR AND SIDE PANELS FOR  
 ADDITIONAL PRECAUTIONS AND FIRST AID INSTRUCTIONS  
 IF NECESSARY, CONTACT:

**NOR-AM**

NET CONTENTS

# PRECAUTIONARY STATEMENTS DANGER

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Fatal if swallowed. Do not breathe spray mist. Can cause irreversible eye damage. May cause skin sensitization. Do not get in eyes, on skin or on clothing. Wear goggles, face shield or safety glasses. Wear a pesticide respirator approved jointly by the Mining Enforcement and Safety Administration (formerly U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part II. Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash before reuse.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to wildlife. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Cover or incorporate spills.

This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treatment area. Do not allow animals to graze in orchard areas treated with CARZOL<sup>®</sup> SP.

## STORAGE AND DISPOSAL

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Completely empty box into application equipment. Then dispose of empty box in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**IN CASE OF FIRE, LEAKY OR DAMAGED CONTAINERS OR OTHER EMERGENCY, REPORT AT ONCE BY TOLL-FREE TELEPHONE TO: 800-424-9300.**

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

#### REENTRY STATEMENT:

Do not enter treated areas for 24 hours unless protective clothing is worn.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure.) When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

**DANGER: Area treated with CARZOL<sup>®</sup> SP on (date of application). Do not enter without appropriate protective clothing for 24 hours or until leaves are completely dry. In case of accidental exposure see Statement of Practical Treatment on the label.**

CARZOL SP is completely soluble in water. Preparation of a premix of slurry before adding to the spray tank is not necessary. Add the desired amount of CARZOL SP directly to the spray tank 1/4 full of water with the agitator running. Then add the remainder of the water.


NOTE: CARZOL SP is not stable in alkaline water. Do not prepare the spray mixture until just before application. Do not prepare more spray mixture than will be used within 4 hours.

### USE PRECAUTIONS

To avoid excessive residues, observe the following limitations:

- ☐ CITRUS: Do not apply more than 5 lbs. CARZOL SP per acre in a 12 month period.  
**Arizona and California:** Allow 14 days between applications.  
**Florida and Texas:** Allow 30 days between applications.  
NOTE: If brown soft scale is present in the grove, a suitable scallicide should be applied after the CARZOL SP application.
- ☐ APPLES, PEARS: Do not apply more than 4 lbs. CARZOL SP per acre in one crop season.
- ☐ PEACHES: Do not apply more than 2 1/2 lbs. CARZOL SP per acre in one crop season.
- ☐ NECTARINES: Do not apply more than 4 lbs. CARZOL SP per acre in one crop season.
- ☐ PLUMS, PRUNES: Do not apply more than 3 lbs. CARZOL SP per acre in one crop season.
- ☐ ALFALFA (Seed Crop Only): Use this product only on pure stands of alfalfa to be harvested for seed. Do not allow spray to drift onto adjacent crops. Do not spray while bees are visiting the treatment area. Do not allow livestock to graze treated fields. Do not use treated crops for livestock feed. Crop residues are not to be fed to livestock or baled for such use. Fields treated with CARZOL SP must be cleared after harvest by burning or by disking into the soil. CARZOL SP does not control aphids. If aphids appear, add an aphicide to the spray. Do not apply more than 2 lbs. CARZOL SP per acre per season. Do not make more than 2 applications of CARZOL SP per season. Allow 21 days before harvest.
- ☐ Do not apply this product through any type of irrigation system.

**BEST AVAILABLE COPY**



To avoid excessive residues observe the pre-harvest intervals shown in the table.  
For other use limitations see side panel under Use Precautions.

CROP	PEST	APPLICATION
ORANGES, LEMONS, TANGELINES, TANGELOS, AND GRAPEFRUIT (Arizona only)	THRIPS	Apply 1/4 to 1 1/4 lbs. of CARZOL SP per acre as outside coverage spray when thrips appear. For aerial applications, use a minimum of 10 gallons of water per acre. Apply as needed up to a limit of 5 lbs. CARZOL SP per acre per 12 month period. Allow 7 days before harvest.
ORANGES, LEMONS, TANGELINES, TANGELOS, AND GRAPEFRUIT (California Only)	THRIPS	Apply 1/4 to 1 1/4 lbs. of CARZOL SP per acre as outside coverage spray when thrips appear. For aerial application, use a minimum of 10 gallons of water per acre. Apply no more than twice per season. Allow 7 days before harvest.
ORANGES, LEMONS, TANGELINES, AND GRAPEFRUIT (Florida)	APOPKA WEEVIL, CITRUS ROOT WEEVIL (BLUE-GREEN WEEVIL), FULLER'S ROSE BEETLE, LITTLE LEAF NOTCHER	Apply 1 lb. CARZOL SP as a concentrate spray of 100 gallons per acre, or less. Repeat application as needed. Allow 7 days before harvest.
ORANGES, LEMONS, TANGELINES, AND GRAPEFRUIT (Florida)	RUST MITE	Apply 1/4-1 lb. CARZOL SP per acre. Repeat application as needed. Allow 7 days before harvest.
ORANGES, LEMONS, TANGELINES, AND GRAPEFRUIT (Tennessee)	RUST MITE	Use 2-4 oz. CARZOL SP per 100 gallons of water when mites appear. For full cover dilute spray, apply 500 gallons of water per acre. Repeat application as needed. Allow 7 days before harvest.
APPLES	TENTIFORM LEAF MINER	Use 4-8 oz. CARZOL SP per 100 gallons of water at pink and petal fall. For full cover dilute spray, apply 400 to 800 gallons of water per acre depending on the size of the trees. For concentrate spray, apply 1-4 lbs. CARZOL SP per acre in at least 50 gallons of water. Allow 7 days before harvest.
APPLES	WHITE APPLE LEAFHOPPER	Use 2-4 oz. CARZOL SP per 100 gallons when leafhoppers appear. For full cover dilute spray, apply 400 gallons of water per acre. For concentrate spray, apply 1/4-1 lb. CARZOL SP per acre in at least 50 gallons of water. Repeat application as needed. Allow 7 days before harvest.
APPLES	CAMPYLOMMA	Use 4 oz. CARZOL SP per 100 gallons of water at pre-bloom/bloom and repeat as necessary through petal fall. For full cover dilute spray, apply 400 gallons of water per acre. For concentrate spray or dilute application of less than 400 gallons, apply 1 lb. CARZOL SP per acre in a minimum of 50 gallons of water per acre. Apply at night or while bees are not working in blooms. Allow 7 days before harvest.
APPLES AND PEARS	EUROPEAN RED MITE, TWO SPOTTED MITE	Use 4-8 oz. CARZOL SP per 100 gallons of water when mites appear. For full cover dilute spray, apply 400 to 800 gallons of water per acre depending on the size of the trees and severity of the mite infestation. For concentrate spray, apply 1-4 lbs. CARZOL SP per acre in at least 50 gallons of water. Repeat application as needed. Allow 7 days before harvest.
APPLES (Also see below for Arizona only)	THRIPS	Use 4-6 oz. CARZOL SP per 100 gallons of water at bloom and repeat as necessary through petal fall. For full cover dilute spray, apply 400 gallons of water per acre. For concentrate spray or dilute application of less than 400 gallons, apply 1 to 1 1/4 lbs. CARZOL SP per acre in a minimum of 50 gallons of water per acre. Apply at night or while bees are not working in blooms. Allow 7 days before harvest.
PEARS	PEAR RUST MITE	Use 4 oz. CARZOL SP per 100 gallons of water. Apply post bloom to harvest at first indication of rust mite infestation. For full cover dilute spray, apply 400-800 gallons of water per acre depending on the size of the trees. For concentrate spray, apply 1-2 lbs. CARZOL SP per acre in at least 50 gallons of water. Repeat application as needed. Allow 7 days before harvest.
PEACHES AND NECTARINES	EUROPEAN RED MITE, McDANIEL MITE, TWO-SPOTTED MITE	Use 4 oz. CARZOL SP per 100 gallons of water when mites appear. For full cover dilute spray, apply 400 to 800 gallons of water per acre depending on the size of the trees. For concentrate spray, apply 1-1 1/4 lbs. CARZOL SP per acre in at least 50 gallons of water. Repeat application as needed. Allow 21 days before harvest for peaches, 14 days for nectarines.
PEACHES	LYGUS BUGS, STINK BUGS	Use 4 oz. CARZOL SP per 100 gallons of water at first sign of infestation. For full cover dilute spray, apply 400-800 gallons of water per acre. For concentrate spray, apply 1-1 1/4 lbs. CARZOL SP per acre. Repeat application as needed. Allow 21 days before harvest.
NECTARINES	THRIPS	Use 4-8 oz. CARZOL SP per 100 gallons of water when thrips infect blossoms through petal fall. For full cover dilute spray, apply 400 to 800 gallons of water per acre depending on the size of the trees. For concentrate or aerial spray, apply 1-2 lbs. CARZOL SP per acre. Apply at night or while bees are not working in blossoms. Repeat application as needed. Allow 14 days before harvest.
PLUMS AND PRUNES	EUROPEAN RED MITE, McDANIEL MITE, TWO-SPOTTED MITE	Use 4 oz. CARZOL SP per 100 gallons of water when mites appear. For full cover dilute spray apply 400 to 800 gallons of water per acre depending on the size of the trees. For concentrate spray, apply 1-1 1/4 lbs. CARZOL SP per acre in at least 50 gallons of water. Repeat application as needed. Allow 7 days before harvest.
ALFALFA (seed crop only in California, Oregon, Washington, Idaho, and Nevada. Use pre-bloom only on alfalfa in all areas except California)	LYGUS BUGS	Use 1/4-1 lb. CARZOL SP per acre. Apply when lygus bugs reach damaging numbers. Contact your State Agricultural Experiment Station or State Extension Service Specialist for proper timing.
	CONSPICUOUS STINK BUGS (California only)	Use 1 lb. CARZOL SP per acre for breakdown of a heavy nymphal population. A lygus control program will control stink bugs. When applied by air use a minimum of 5 gallons per acre. When applied by ground use a minimum of 20 gallons per acre.
	TWO-SPOTTED MITE	Use 1/4-1 lb. CARZOL SP per acre. Apply when mites appear. When applied by air use a minimum of 10 gallons of spray per acre. When applied by ground use a minimum of 20 gallons per acre.

**IMPORTANT: READ BEFORE USE**


By using this product, user or buyer accepts the following conditions, warranty, disclaimer of warranties and limitations of liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However, because of extreme weather and soil conditions, manner of use and other factors beyond NOR-AM Chemical Company's control, it is impossible for NOR-AM to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OR MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of NOR-AM Chemical Company is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. UNDER NO CIRCUMSTANCES SHALL NOR-AM CHEMICAL COMPANY BE LIABLE FOR ANY SPECIAL, INDIRECT, OR CONSEQUENTIAL DAMAGES RESULTING FROM USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES, OR DAMAGE RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT NOR-AM'S ELECTION, THE REPLACEMENT OF PRODUCT.

© NOR-AM Chemical Company, 1991



**NOR-AM**  
NOR-AM CHEMICAL COMPANY  
A Schering-Plough Company  
AGRI-BUSINESS DIVISION  
3500 Silverside Road, P.O. Box 7496  
Wilmington, DE 19803