

Reg # 45639-146

PM-19

193

SPECIMEN LABEL

ACCEPTED

APR -7 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 45639-146

Ovasyn[®]

INSECTICIDE / MITICIDE

dae.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

In case of spillage, cover with an absorbent such as soda ash, lime, clay, or sawdust. Sweep up and bury. Wash area thoroughly with detergent and water.


NOTICE TO DRUM HANDLERS:

In Case of Fire, Leaky, or Damaged Containers, or Other Emergency, Report at Once by Toll-Free Telephone to: 800-424-9300.

DO NOT REUSE EMPTY CONTAINER.

dent exposure, as described under Precautionary Statements on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

WARNING: Area treated with OVASYN® on (date of application). Do not enter without appropriate protective clothing for 24 hours or until sprays have dried. In case of accidental exposure see Statement of Practical Treatment on the label.



BEST AVAILABLE COPY

GENERAL INFORMATION

OVASYN is an insecticide/miticide with ovicidal and synergistic activity on bollworm, tobacco budworm, pink bollworm, whitefly and mites on cotton. OVASYN may be applied alone for control of two-spotted spider mite, strawberry mite, pacific mite, and carmine mite. For insecticidal control, OVASYN can be applied alone or tank-mixed with other insecticides registered for use on cotton.

OVASYN is not phytotoxic to young cotton.

APPLICATION

Equipment: Apply by air or ground, using low drift nozzles. When applying by air, use 1-5 gallons of water per acre. For applications made with ground equipment, use 10-50 gallons of water per acre. Use of the higher volume of water will assure better coverage for mite and whitefly control in larger, more dense cotton.

Mite Control: Apply OVASYN as a single treatment of 2 3/4 to 5 pints per acre, or two sequential applications of 2 3/4 pints per acre. DO NOT exceed 5 1/2 pints per acre per season. Treatments should be applied when mite populations begin to build. OVASYN may be applied from the time cotton plants are 4-6 inches tall and up until boll opening. DO NOT apply after cotton bolls are open.

Insect Control (Ovicide/Synergist): In fields where bollworm, tobacco budworm, pink bollworm or other insect pests have reached economic levels, apply OVASYN at 3/4 to 1 1/2 pints per acre as needed or on an appropriate schedule. To control whitefly, apply 1 1/2 to 5 1/2 pints per acre in sufficient water to ensure complete coverage. DO NOT exceed 5 1/2 pints OVASYN per acre per season. DO NOT apply after cotton bolls are open. Use other registered cotton insecticides in a tank-mix at rates recommended on the labels for those products and observe all limitations and precautions which appear on those labels. Do not tank-mix OVASYN with Cyprex¹, Ferbam², bordeaux mixture or other highly alkaline materials.

TABLE OF APPLICATION RATES

Desired Rate/AC. (lbs. AI)	Acres Treated Per Gallon	Pints Per Acre
.125	= 12 AC.	= .66
.17	= 9 AC.	= .89
.25	= 6 AC.	= 1.33
.50	= 3 AC.	= 2.66
.75	= 2 AC.	= 4.0
1.0	= 1.5 AC.	= 5.33

¹Trademark of American Cyanamid
²Trademark of FMC Corp.

ULV Applications: OVASYN may be applied in vegetable, cottonseed or similar oils. After determination of the required volume of OVASYN, add sufficient oil to apply 1 to 2 quarts total liquid per acre. If tank-mixing with additional cotton insecticides which are registered for ULV applications, their volume must be accounted for in the total 1 to 2 quarts per acre.

RESTRICTIONS AND LIMITATIONS

- DO NOT** apply more than 5 1/2 pints per acre OVASYN per growing season.
- DO NOT** apply OVASYN after cotton bolls are open.
- DO NOT** apply shortly before rainfall.
- DO NOT** allow prepared solutions to stand overnight.
- DO NOT** feed foliage from treated cotton plants to livestock.

IMPORTANT: READ BEFORE USE


By using this product, user or buyer accepts the following conditions, warranty, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, because of extreme weather and soil conditions, manner of use and other factors beyond NOR-AM Chemical Company's control, it is impossible for NOR-AM to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OR MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of NOR-AM Chemical Company is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. UNDER NO CIRCUMSTANCES SHALL NOR-AM CHEMICAL COMPANY BE LIABLE FOR ANY SPECIAL, INDIRECT, OR CONSEQUENTIAL DAMAGES RESULTING FROM USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES, OR DAMAGE RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT NOR-AM'S ELECTION, THE REPLACEMENT OF PRODUCT.

© NOR-AM Chemical Company, 1991



BEST AVAILABLE COPY