

PM 25 45639-159

1046

US ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (75-767)
WASHINGTON, DC 20460

EPA REGISTRATION NO.
45639-159

DATE OF ISSUANCE

APR - 5 1994

TERM OF ISSUANCE

NOTICE OF PESTICIDE: REGISTRATION
 REREISTRATION

(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

NAME OF PESTICIDE PRODUCT

NA 301 ...
Herbicide

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

MOR-AM Chemical Co.
Little Falls Centre One
2711 Centerville Rd.
Wilmington, DE 19868

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(B) provided that you:

1. Submit/deliver all data required for your product's reregistration of your product under FIFRA Section 3(c)(5) and sec. 4 when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for shipment:

a. Add the phrase "EPA Registration No. 45639-159".

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the 4-77 enclosure for a further description of final printed labeling.

If these conditions are not complied with, this registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes your acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

BEST AVAILABLE COPY

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

Robert H. ...

DATE

APR 5 1994

SPECIMEN LABEL

FOR AGRICULTURAL USE ONLY

ACCEPTED

APR - 5 1995

The Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under Reg. No. 45639-159

NA 308

SUGAR BEET HERBICIDE

Postemergence Herbicide for Control of Weeds in Sugar Beets

ACTIVE INGREDIENTS:

Phenmedipham*	70%
Desmedipham**	70%
Ethofumesate***	70%
INERT INGREDIENTS:	79.0%
TOTAL:	100.0%

Contains 1.8 lbs. active ingredient per gallon.
 This product contains the toxic inert ingredient isophorone.
 *3-methoxycarbonylamino-phenyl-3-methylcarbamilate
 **Ethyl m-hydroxycarbamate carbamate (ester)
 ***2-ethoxy-2,3-dihydro-3,3-dimethyl-5-benzofuranyl methanesulfonate

EPA Reg. No. 45639-

EPA Est. No. 407-MN-1

KEEP OUT OF REACH OF CHILDREN DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Give a glass of water. If vomiting occurs, do not give more water. Do not give any food. If a person has swallowed a large quantity of water, do not attempt to give anything else to drink. Seek medical attention. See note to physician below.

NOTE TO PHYSICIAN: This product contains arsenate and other organophosphorus compounds which are important to the treatment of certain types of poisoning. If a patient has had a previous poisoning, should be treated with special care with an antidote if available. It is recommended that the physician be notified of the product name and the material damage by contact may contain the name of the manufacturer.

IF INHALED: Move the patient to fresh air. If breathing is difficult, give oxygen. If breathing is stopped, give artificial respiration. See note to physician below.

IF IN EYES: Flush eyes with water for at least 15 minutes. If irritation persists, consult a physician.

IF ON SKIN: Wash with soap and water. If irritation persists, consult a physician.

THIS ECHO SYSTEM™ CONTAINER IS THE SOLE PROPERTY OF NOR-AM CHEMICAL COMPANY.
 SEE OTHER PANELS FOR ADDITIONAL PRECAUTIONS

NOR-AM
 NOR-AM CHEMICAL COMPANY
 A Schering-Plough Company

AGRI-BUSINESS DIVISION
 Little Fall's Centre One
 2711 Centerville Road
 Wilmington, DE 19808

NET CONTENTS:

U.S. Patent No. 3,404,976
 U.S. Patent No. 3,692,820
 U.S. Patent No. 3,689,507

BEST AVAILABLE COPY

3076

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
(and Domestic Animals)
DANGER**

PRECAUTIONARY STATEMENTS
CORROSIVE. CAUSES IRREVERSIBLE EYE DAMAGE. DO NOT GET IN EYES OR ON CLOTHING. HARMFUL IF SWALLOWED OR INHALED. AVOID BREATHING VAPOR OR SPRAY MIST.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category B on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- > Long-sleeved shirt & long pants
- > Shoes plus socks
- > Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

THIS PESTICIDE IS TOXIC TO FISH AND AQUATIC ORGANISMS. FOR TERRESTRIAL USES, DO NOT APPLY DIRECTLY TO WATER, OR TO AREAS WHERE SURFACE WATER IS PRESENT, OR TO INTERTIDAL AREAS BELOW THE MEAN HIGH WATER MARK. DRIFT AND RUNOFF FROM TREATED AREAS MAY BE HAZARDOUS TO FISH AND AQUATIC ORGANISMS IN ADJACENT AQUATIC SITES. DO NOT CONTAMINATE WATER THROUGH DISPOSAL OF EQUIPMENT WASHWATERS.

PHYSICAL OR CHEMICAL HAZARDS
DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep closed. Store in a cool, dry place. Do not use or store near heat or open flame. Protect NA 308 from freezing temperatures.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: ECHO-SYSTEM SVR Return Procedure: Return the ECHO-SYSTEM SVR container clean (outside only) and empty to the place of business from which the NA 308 was purchased.

IN CASE OF FIRE, LEAKY OR DAMAGED CONTAINERS, OR OTHER EMERGENCY, REPORT AT ONCE BY TOLL-FREE TELEPHONE TO: 800-424-9300.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read the entire Directions for Use before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- > Coveralls
- > Water-proof gloves
- > Shoes plus socks
- > Protective eyewear

BEST AVAILABLE COPY

When used as directed, NA 308 is selective in beets past the two-true leaf stage. Beets may be severely injured if treated before the two-true leaf stage. The stage of growth of the weeds is very important for satisfactory control. For best results, spray when the weeds are at the two-true leaf stage. Best results are obtained when the weeds are actively growing and are not under water or heat stress.

Mixing the Spray: Make sure the Sprayer is CLEAN. NA 308 contains sufficient wetting agents for optimum coverage. Do not add additional wetting agents or other spray adjuvants. Add sufficient water to fill the lines. Then add the desired amount of NA 308 and the remaining quantity of water with the bypass agitator running. Bypass agitation is sufficient. Mechanical agitation is not necessary. Only use freshly prepared spray emulsions.

Always spray immediately after preparing the spray solution. Prepare only enough spray solution to last less than four hours.

GENERAL INFORMATION

NA 308 is a selective postemergence herbicide for use in sugar beets (including beets grown for seeds) for the control of the following weeds.

Annual bluegrass	<i>Poa annua</i>
Annual sowthistle	<i>Sonchus oleraceus</i>
Black nightshade	<i>Solanum nigrum</i>
Hairy nightshade	<i>Solanum sarrachoides</i>
Canary grass	<i>Phalaris canariensis</i>
Coast fiddleneck	<i>Amsinckia intermedia</i>
Common chickweed	<i>Stellaria media</i>
Common lambsquarters	<i>Chenopodium album</i>
Common ragweed	<i>Ambrosia artemisiifolia</i>
Green foxtail	<i>Setaria viridis</i>
Groundcherry	<i>Physalis lanceifolia</i>
Kochia*	<i>Kochia scoparia</i>
Ladysthumb	<i>Polygonum persicaria</i>
London rocket	<i>Sisymbrium irio</i>
Nettleleaf goosefoot	<i>Chenopodium murale</i>
Pennsylvania smartweed ..	<i>Polygonum pennsylvanicum</i>
Pigeon grass (yellow foxtail)	<i>Setaria glauca</i>
Prostrate pigweed	<i>Amaranthus gracizans</i>
Purslane	<i>Portulaca oleraceus</i>
Redroot pigweed	<i>Amaranthus retroflexus</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Wild buckwheat	<i>Polygonum convolvulus</i>
Wild mustard	<i>Brassica kaber</i>

* Spray kochia while in the rosette stage, less than 1" in diameter. For best results, spray when the weeds are at the two-leaf stage.

USE PRECAUTIONS: DO NOT APPLY NA 308 TO SUGAR BEETS LATER THAN 75 DAYS PRIOR TO HARVEST.

DO NOT EXCEED A TOTAL OF 13.0 PINTS NA 308 PER ACRE PER SEASON.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

4096
NA 308 MAY CAUSE BEET INJURY IF THE CROP IS UNDER STRESS FROM ONE OR MORE OF THE FOLLOWING CONDITIONS:

- ▲ Rapid climatic changes from cool, overcast days, to hot (85° or over) bright days. When the air temperature is, or is likely to be, above 80°F on the day of spraying, application should be made in the evening when the temperature is lower.
- ▲ Windy conditions or drought.
- ▲ Use of a preplant or preemergence herbicide or other chemicals.
- ▲ Insect or disease injury.
- ▲ Close cultivation.

If stress conditions are present delay application in order to give plants a chance to recover.

If extreme weather conditions are of short duration, delay spraying until the end of such a period. If NA 308 applications must be made on days with extreme temperature and/or brightness, delay spraying until evening.

DO NOT OVERTREAT: The use of higher than recommended rates may cause beet injury.

Do not spray while dew is present.

Rainfall within 6 hours of spraying may reduce weed kill. Do not allow spray drift to contact adjacent crops which may be injured by spray drift.

IMPORTANT: NA 308 may cause temporary growth retardation and/or chlorosis or tipburn on sugar beets. Beets usually resume normal growth within 10 days.

RATE OF APPLICATION

By Ground: Apply NA 308 at the rate of 3.25 to 5.0 pints in 20 to 50 gallons water broadcast basis. For band application, see dosage chart.

By Air: Apply NA 308 at the rate of 3.25 to 5.0 pints per acre using 5 to 15 gallons of spray per acre.

Apply the 3.25 to 5.0 pint rates only to sugar beets past the two-true leaf stage. Use the 5.0 pint rate only on well established beets which are not under stress.

In order to avoid phytotoxic spray drift to non-target crops during application of NA 308, the following buffer zones should be observed:

Cotton, Potatoes, Sunflowers,

Sorghum, Wheat 50 feet

Blackeye beans, Cabbage, Flax 100 feet

Lettuce, Rape, Tomatoes 300 feet

DO NOT APPLY WHEN WIND SPEED IS OVER 10 MILES PER HOUR. AVOID APPLICATIONS WHEN CONDITIONS FAVOR DRIFT.

To avoid excessive phytotoxicity to fall-planted sugar beets in Arizona and south of the Tehachapi Mountains in California when temperatures are above 85° F, apply NA 308 at the rate of 3.25 pints per acre (broadcast equivalent). Evening applications are recommended.

For further information, contact your County Agricultural Agent, Farm Advisor or NOH-AM Chemical Company.

BEST AVAILABLE COPY

50/6

REPEAT APPLICATION OF NA 308: For control of later germinating weeds, make a second application of NA 308. Use 3.25 to 5.0 pints of NA 308. Allow at least 7 days between first and second applications. Apply when beets have at least 4 leaves. For best results, use the higher rate and spray when weeds are at the two-true leaf stage. Apply lower rates when the beets are under stress as explained on the previous page, the General Information section.

SPLIT (LOW RATE) APPLICATIONS: Split (low rate) applications of NA 308 may be applied to sugar beets to control early germinating weeds. The first spray must be applied when the earliest emerging weeds have reached cotyledon size. See Chart 1, below, for broadcast rates. See Chart 2, below, for equivalent band rates. For band applications, apply in 5-10 gallons of water per acre. Any weeds which are not completely controlled by the first treatment will usually be checked and controlled by repeat applications. The repeat application should be made 5 to 7 days after the preceding application or when another flush of weeds germinates. If the second application is delayed, conventional treatment as described above will be necessary.

NA 308 broadens and enhances the control of troublesome weeds, including the following:

ANNUAL BROADLEAF WEEDS

- Annual sowthistle *Sonchus oleraceus*
- Black nightshade *Solanum nigrum*
- Hairy nightshade *Solanum sarrachoides*
- Coast fiddleneck *Amsinckia intermedia*
- Common chickweed *Stellaria media*
- Common lambsquarters *Chenopodium album*

- Common ragweed *Ambrosia artemisiifolia*
- Groundcherry *Physalis lanceifolia*
- Kochia* *Kochia scoparia*
- Ladysthumb *Polygonum persicaria*
- London rocket *Sisymbrium irio*
- Nettleleaf goosefoot *Chenopodium murale*
- Pennsylvania smartweed .. *Polygonum pennsylvanicum*
- Prostrate pigweed *Amaranthus gracizans*
- Purslane *Portulaca oleraceus*
- Redroot pigweed *Amaranthus retroflexus*
- Shepherdspurse *Capsella bursa-pastoris*
- Wild buckwheat *Polygonum convolvulus*
- Wild mustard *Brassica kaber*

* Spray Kochia while in the rosette stage, less than one inch in diameter.

ANNUAL GRASS WEEDS

- Annual bluegrass *Poa annua*
- Canarygrass *Phalaris canariensis*
- Green foxtail *Setaria viridis*
- Pigeon grass (Yellow foxtail) *Setaria glauca*

See Chart 1, below, for broadcast rates. See Chart 2 for equivalent band rates.

**CHART 1
DOSAGE CHART FOR BROADCAST APPLICATION**

Sugar Beet Stage	Pints/Acre Broadcast
	NA 308
Cotyledon	1.75
2 leaf	2.25
4 leaf	3.25

**CHART 2
DOSAGE CHART FOR BAND APPLICATION**

Broadcast Equivalent	Band Width	Band Rate — Row Spacing		
		22"	28"	30"
1.75 pints/acre	5"	6.4 fl. oz.	5.0 fl. oz.	4.7 fl. oz.
	7"	8.9 fl. oz.	7.0 fl. oz.	6.5 fl. oz.
2.25 pints/acre	5"	8.2 fl. oz.	6.4 fl. oz.	6.0 fl. oz.
	7"	11.5 fl. oz.	9.0 fl. oz.	8.4 fl. oz.
3.25 pints/acre	5"	11.8 fl. oz.	9.3 fl. oz.	8.7 fl. oz.
	7"	16.5 fl. oz.	13.0 fl. oz.	12.1 fl. oz.
5.0 pints/acre	5"	18.2 fl. oz.	14.3 fl. oz.	13.3 fl. oz.
	7"	25.5 fl. oz.	20.0 fl. oz.	18.7 fl. oz.

BEST AVAILABLE COPY

6046

IMPORTANT: READ BEFORE USE

By using this product, user or buyer accepts the following conditions, warranty, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, because of extreme weather and soil conditions, manner of use and other factors beyond NOR-AM Chemical Company's control, it is impossible for NOR-AM to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of NOR-AM Chemical Company is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. NOR-AM disclaims any liability whatsoever for incidental or consequential damages, including, but not limited to, liability arising out of breach of contract, express or implied warranty (including warranties of merchantability and fitness for a particular purpose), tort, negligence, strict liability or otherwise.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT NOR-AM'S ELECTION, THE REPLACEMENT OF PRODUCT.

© NOR-AM Chemical Company, 1993

BEST AVAILABLE COPY



AGRI-BUSINESS DIVISION
Little Falls Centre One
2711 Centerville Road
Wilmington, DE 19806

